

PHARMACY POLICY STATEMENT Kentucky Medicaid	
DRUG NAME	Cimzia (certolizumab pegol)
BILLING CODE	Must use valid NDC code
BENEFIT TYPE	Pharmacy
SITE OF SERVICE ALLOWED	Home
COVERAGE REQUIREMENTS	Prior Authorization Required (Non-Preferred Product) Alternative preferred products include Enbrel & Humira (if appropriate for indication) QUANTITY LIMIT – 1200 per 28 days
LIST OF DIAGNOSES CONSIDERED <b>NOT</b> MEDICALLY NECESSARY	<a href="#">Click Here</a>

Cimzia (certolizumab pegol) is a **non-preferred** product and will only be considered for coverage under the **pharmacy** benefit when the following criteria are met:

Members must be clinically diagnosed with one of the following disease states and meet their individual criteria as stated.

### ANKYLOSING SPONDYLITIS (AS)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Must have a documented negative TB test (i.e. tuberculosis skin test (PPD), an interferon-release assay (IGRA), or a chest x-ray) within 6 months prior to starting therapy; AND
3. Medication must be prescribed by a rheumatologist; AND
4. Member has had back pain for 3 months or more that began before the age of 45; AND
5. Current imaging results show an inflammation of one or both of the sacroiliac joints; AND
6. Member shows at least **one** of the following signs or symptoms of Spondyloarthritis:
  - a) Arthritis;
  - b) Elevated serum C-reactive protein;
  - c) Inflammation at the tendon, ligament or joint capsule insertions;
  - d) Positive HLA-B27 test;
  - e) Limited chest expansion;
  - f) Morning stiffness for 1 hour or more; AND
7. Member meets at least **one** of the following scenarios:
  - a) Member has Axial (spinal) disease;
  - b) Member has peripheral arthritis without axial involvement and has tried and failed treatment with methotrexate or sulfasalazine. Treatment failure requires at least 30 days of therapy without an adequate response; AND
8. Member has tried and failed to respond to treatment with at least 2 prescription NSAIDs taken at the maximum recommended dosages. Treatment failure requires at least 30 days of therapy without an adequate response; AND
9. Member has tried and failed treatment with **both** Enbrel and Humira. Treatment failure requires at least 30 days of therapy (for each drug) without an adequate response.
10. **Dosage allowed:** Inject 400 mg subcutaneously once a week at weeks 0, 2, and 4 and then 200 mg every other week or 400 mg every four weeks.

***If member meets all the requirements listed above, the medication will be approved for 12 months.***

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease.

***If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.***

### **CROHN'S DISEASE (CD)**

For **initial** authorization:

1. Member is 18 years of age or older with moderate to severe active CD; AND
2. Must have a documented negative TB test (i.e. tuberculosis skin test (PPD), an interferon-release assay (IGRA), or a chest x-ray) within 6 months prior to starting therapy; AND
3. Medication must be prescribed by a gastroenterologist; AND
4. Member has tried and failed at least 30 days of treatment with Humira; AND
5. Member has had a documented trial and inadequate response to a 30-day trial of 6-mercaptopurine, azathioprine, methotrexate or corticosteroids; OR
6. Member has severe disease, as indicated by at least **one** of the following:
  - a) Esophageal or gastroduodenal disease;
  - b) Extensive small-bowel disease involving more than 100 cm;
  - c) History of colonic resection;
  - d) History of two or more small-bowel resections;
  - e) Perianal or rectal disease.
7. **Dosage allowed:** Inject 400 mg subcutaneously once a week at weeks 0, 2, and 4 and then 400 mg every four weeks.

***If member meets all the requirements listed above, the medication will be approved for 12 months.***

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease.

***If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.***

### **PSORIATIC ARTHRITIS (PsA)**

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Must have a documented negative TB test (i.e. tuberculosis skin test (PPD), an interferon-release assay (IGRA), or a chest x-ray) within 6 months prior to starting therapy; AND
3. Medication must be prescribed by a rheumatologist or dermatologist; AND
4. Member has tried and failed treatment with **both** Enbrel and Humira. Treatment failure requires at least 30 days of therapy (for each drug) without an adequate response; AND
5. Member meets at least **one** of the following scenarios:

- a) Member has predominantly axial disease (i.e. sacroiliitis or spondylitis) as indicated by radiographic evidence;
  - b) Member has shown symptoms of predominantly axial disease (i.e. sacroiliitis or spondylitis) for more than 3 months (i.e. limited spinal range of motion, spinal morning stiffness for more than 30 minutes) AND has tried and failed to respond to treatment with at least 2 prescription NSAIDs taken at the maximum recommended dosages. Treatment failure requires at least 30 days of therapy without an adequate response;
  - c) Member has predominately non-axial disease and has tried and failed to respond to treatment with at least 30-day trial of methotrexate and 30-day trial NSAID.
6. **Dosage allowed:** Inject 400 mg subcutaneously once a week at weeks 0, 2, and 4 and then 200 mg every other week or 400 mg every four weeks.

***If member meets all the requirements listed above, the medication will be approved for 12 months.***

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease.

***If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.***

## RHEUMATOID ARTHRITIS (RA)

For **initial** authorization:

1. Member must be 18 years of age or older with moderate to severe active RA; AND
2. Must have a documented negative TB test (i.e. tuberculosis skin test (PPD), an interferon-release assay (IGRA), or a chest x-ray) within 6 months prior to starting therapy; AND
3. Medication must be prescribed by a rheumatologist; AND
4. Member must have tried and failed treatment with at least **two** non-biologic DMARDS (i.e. methotrexate, hydroxychloroquine, sulfasalazine, azathioprine, cyclosporine and leflunomide) or must have documented contraindication to all non-biologic DMARDS. Treatment trial duration with each non-biologic DMARD agent must have been at least 30 days; AND
5. Member must have tried and failed treatment with **both** Enbrel and Humira. Treatment failure requires at least 30 days of therapy (for each drug) without an adequate response.
6. **Dosage allowed:** Inject 400 mg subcutaneously once a week at weeks 0, 2, and 4 and then 200 mg every other week thereafter.

***If member meets all the requirements listed above, the medication will be approved for 12 months.***

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease.

***If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.***



**CareSource considers Cimzia (certolizumab pegol) not medically necessary for the treatment of the following disease states based on a lack of robust clinical controlled trials showing superior efficacy compared to currently available treatments:**

- Active infections
- Asthma
- Cellulitis
- Dissecting scalp cellulitis
- For use in combination with other TNF-inhibitors (i.e. Kineret, Enbrel, Remicade)
- Giant-cell arteritis
- Guttate psoriasis
- Infectious uveitis
- Lupus perino
- Osteoarthritis
- Recurrent pregnancy loss
- Relapsing polychondritis
- Sarcoidosis
- Sciatica
- Spondyloarthritis (other than ankylosing spondylitis)
- Takayasu's arteritis
- Vogt-Koyanagi

DATE	ACTION/DESCRIPTION
05/15/2017	New policy for Cimzia created. Policies SRx-0041 and SRx-0042 achieved. New diagnosis of AS with criteria was added. For diagnosis of CD: TNF inhibitor Humira and corticosteroids trials were added. For PsA: TNF inhibitors Humira and Enbrel were listed as required trials. For RA: non-biologic DMARDS were listed, and TNF inhibitors Humira and Enbrel were listed as required trials. Trials length changed to 30 days for each drug trial due to KY MCD regulations. List of diagnosis considered not medically necessary was added.

References:

1. Cimzia [prescribing information]. Smyrna, GA: UCB, Inc.; Revised October 2013.
2. US Food and Drug Administration Drug Safety Data. [http://www.accessdata.fda.gov/drugsatfda\\_docs/label/2014/125261s114lbl.pdf](http://www.accessdata.fda.gov/drugsatfda_docs/label/2014/125261s114lbl.pdf) (October 14, 2014).
3. Menter A, Gottlieb A, Feldman SR, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 1. Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. J Am Acad Dermatol. 2008 May;58(5):826-50.
4. Gottlieb A, Korman NJ, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. J Am Acad Dermatol. 2008 May;58(5):851-64.
5. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: section 4. Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. J Am Acad Dermatol. 2009 Sep;61(3):451-85. Epub 2009 Jun 3.
6. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis Section 6. Guidelines of care for the treatment of psoriasis and psoriatic arthritis: Case-based presentations and evidence-based conclusions. J Am Acad Dermatol. 2011 Feb 7. [Epub ahead of print].

7. National Institutes of Health, Clinicaltrials.gov. [cited 9/16/2014]; Available from: <http://www.clinicaltrials.gov>.
8. Singh, J., et al., (2012). 2012 Update of the 2008 American College of Rheumatology Recommendations for the Use of Disease-Modifying Anti-rheumatic Drugs and Biologic Agents in the Treatment of Rheumatoid Arthritis. *Arthritis Care & Research*, 64(5), 625-639.
9. Rheumatoid Arthritis. *Arthritis & Rheumatology (Arthritis Care & Research)*, 64:5, 625-639, May 2012.
10. Wolters Kluwer. Facts & Comparisons. [www.factsandcomparisons.com](http://www.factsandcomparisons.com). 2014.
11. *Gastroenterology*. 2013 Dec;145(6):1459-63.
12. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring of therapeutic agents for the treatment of arthritis and systemic features. *Arthritis Care Res. (Hoboken)*. 2011 Apr;63(4):465-82.
13. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 Recommendations for the Use of Non- biologic and Biologic Disease-Modifying Anti-rheumatic Drugs in Rheumatoid Arthritis. *Arthritis Care & Research. Arthritis Rheum* 2008;59(6):762-84.
14. Lichtenstein GR, Hanauer SB, Sandborn WJ, Practice Parameters Committee of American College of Gastroenterology. Management of Crohn's disease in adults. *American Journal of Gastroenterology* 2009;104(2):465-83; quiz 464, 484. DOI: 10.1038/ajg.2008.168. (Reaffirmed 2014 Oct)
15. Wolters Kluwer. Facts & Comparisons. [www.factsandcomparisons.com](http://www.factsandcomparisons.com), 2015.
16. Sandborn, W., Binion, D., Persley, K., Atreja, A., & Kosinski, L. (2014). AGA Institute Guidelines for the Identification, Assessment and Initial Medical Treatment in Crohn's Disease: Clinical Decision Support Tool. AGA Institute. Retrieved August 14, 2015, from [www.gastro.org/IBDcarepathway](http://www.gastro.org/IBDcarepathway).
17. Gottlieb A, Korman NJ, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. *J Am Acad Dermatol*. 2008 May;58(5):851-64.

Effective date: 05/15/2017

Revised date: 05/08/2017