



AAT DEFICIENCY

Products Affected

- ARALAST NP
- GLASSIA

- PROLASTIN-C INTRAVENOUS **SOLUTION**
- ZEMAIRA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of severe congenital A1-PI deficiency who have clinically evident emphysema, weight, A1-PI phenotype, A1-PI baseline level
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a pulmonologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ABILIFY IM

Products Affected

• ABILIFY MAINTENA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and other treatments tried and outcome.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a psychiatrist or mental health specialist.
Coverage Duration	5 years
Other Criteria	Patient must a have a reason aripiprazole oral cannot be used.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ACTIMMUNE

Products Affected

• ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies, and result of prior therapy.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Must be prescribed by a hematologist or oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ACTINIC KERATOSIS

Products Affected

• diclofenac sodium topical gel 3 %

DA C '	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Prior use of 5% fluorouracil topical and 5% imiquimod topical, unless contraindicated.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ADAPALENE

Products Affected

- adapalene topical cream
- adapalene topical gel 0.3 %
- adapalene topical gel with pump
- adapalene topical solution
- adapalene topical swab

PA Criteria	Criteria Details
Exclusion Criteria	Not approved when used to treat photo aging.
Required Medical Information	Diagnosis, previous treatments, and response therapy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Must have failure, intolerance, or contraindication to tretinoin
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ADBRY

Products Affected

• ADBRY SUBCUTANEOUS SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used and result of prior therapy. If continuation, response to tralokinumab with documented reduction in number of acute exacerbations.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information
Prescriber Restrictions	Prescribing limited to an allergist, immunologist, pulmonologist, otolaryngologist or dermatologist.
Coverage Duration	3 years
Other Criteria	For atopic dermatitis: trial of at least one topical corticosteroid (fluticasone, fluocinonide, desonide) - and - one at least one topical immunomodulator (tacrolimus, pimecrolimus).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ADEMPAS

Products Affected

• ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Confirmation of diagnosis, documentation of response to any prior therapies
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by pulmonologist or cardiologist.
Coverage Duration	3 year
Other Criteria	For WHO Group 1 diagnosis, patient must have a history of taking or contraindication to sildenafil (Revatio).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ADLARITY

Products Affected

ADLARITY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous treatments, and response therapy
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Must have a intolerance, contraindication, or medical reason the tablets are not acceptable.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





AFINITOR

Products Affected

• everolimus (antineoplastic)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR THERAPIES, AND TREATMENT RESPONSE
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST OR NEUROLOGIST
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

AIMOVIG

Products Affected

• AIMOVIG AUTOINJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried, and outcome. Member has been evaluated for and does not have medication overuse headache.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Patient must have an inadequate response, contraindication, or intolerance to two different chronic migraine prevention drugs. The two prerequisite drugs must be from different classes such as anticonvulsants (topiramate/valproate), beta blockers (propranolol, metoprolol), and antidepressants (nortriptyline/venlafaxine).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

AJOVY

Products Affected

• AJOVY AUTOINJECTOR

• AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried, and outcome. Member has been evaluated for and does not have medication overuse headache.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Patient must have an inadequate response, contraindication, or intolerance to two different chronic migraine prevention drugs. The two prerequisite drugs must be from different classes such as anticonvulsants (topiramate/valproate), beta blockers (propranolol, metoprolol), and antidepressants (nortriptyline/venlafaxine).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





AKEEGA

Products Affected

AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. Member must have metastasis from malignant tumor of prostate, Castration-resistant, deleterious or suspected deleterious BRCA-mutated.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	prescribed by an Oncologist or Urologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ALK POSITIVE TYROSINE KINASE INHIBITORS

Products Affected

- ALECENSA
- XALKORI ORAL CAPSULE
- XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG
- ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other treatments tried and outcome
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective. 12/01/2024

ALUNBRIG

Products Affected

• ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Metastatic non-small cell lung cancer (NSCLC): must be ALK-positive, as detected by an approved test.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	5 years
Other Criteria	Approve for continuation of prior therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





AMIFAMPRIDINE

Products Affected

• FIRDAPSE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Must have a documented diagnosis of Lamber-Eaton with electrodiagnostic studies including repetitive nerve stimulation and anti-P/Q-type voltagegated calcium channel (VGCC) antibody testing to confirm the diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a neurologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ANALEPTIC

Products Affected

• armodafinil

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Excessive sleepiness due to SWSD defined as the patient is working at least 5 overnight shifts per month. As adjunctive/augmentation treatment for depression in the adult if the patient is concurrently receiving other medication therapy for depression.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ANTIDEPRESSANTS

Products Affected

- AUVELITY
- FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)
- FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR
- TRINTELLIX
- vilazodone

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR THERAPIES TRIED AND FAILED
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	For treatment of major depressive disorder (MDD), must have tried two generic antidepressants from different classes: SSRIs, SNRIs, TCAs, NOREPINEPHRINE-DOPAMINE REUPTAKE INHIBITORS, or NORADRENERGIC and SPECIFIC SEROTONERGIC ANTIDEPRESSANTS. Examples include: sertraline, citalopram, escitalopram, fluoxetine, paroxetine, venlafaxine, duloxetine, bupropion, amitriptyline, doxepin, etc.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Part B Prerequisite	No





Date Effective: 12/01/2024

ANTIPSYCHOTICS

Products Affected

- asenapine maleate
- CAPLYTA
- FANAPT
- REXULTI ORAL TABLET
- SECUADO
- VERSACLOZ
- VRAYLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and treatement history.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	For BIPOLAR DISORDER or SCHIZOPHRENIA, documentation of diagnosis, and treatment failure with two atypical anti-psychotics: ZIPRASIDONE, RISPERIDONE, QUETIAPINE, OLANZAPINE, CLOZAPINE, ARIPIPRAZOLE) or rationale as to why alternatives are not suitable. For treatment of major depressive disorder (MDD), must have tried generic antidepressants from at least two different classes: SSRIs, SNRIs, TCAs, NOREPINEPHRINE-DOPAMINE REUPTAKE INHIBITORS, or NORADRENERGIC and SPECIFIC SEROTONERGIC ANTIDEPRESSANTS.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Part B Prerequisite	No





Date Effective: 12/01/2024

APREPITANT

Products Affected

• aprepitant

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For use with highly and moderately-emetogenic chemotherapy, provide the chemotherapy regimen including drug, dose, and frequency. Ondansetron is preferred for post-operative nausea/vomiting (PONV) prophylaxis. When aprepitant is used for PONV prophylaxis, provide rationale as to why ondansetron is not a suitable alternative.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Request will also be reviewed for coverage under part B versus part D.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ARCALYST

Products Affected

• ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	1 YEAR AT A TIME
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

AUGTYRO

Products Affected

• AUGTYRO ORAL CAPSULE 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

AURYXIA

Products Affected

AURYXIA

PA Criteria	Criteria Details
Exclusion Criteria	Not approved for treatment of iron deficiency anemia in patients with ESRD on dialysis.
Required Medical Information	Diagnosis, CKD/ESRD Stage, prior therapy used and response to prior therapy. Required pre-treatment testing for hyperphosphatemia: serum calcium and phosphorus, serum creatinine and eGFR.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a nephrologist.
Coverage Duration	3 year
Other Criteria	For hyperphosphatemia, documentation of prior use of calcium acetate and one other drug: sevelamer or lanthanum.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

AYVAKIT

Products Affected

• AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to avapritinib.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by an oncologist or gastroenterologist - or - an allergist or immunologist, as appropriate to the diagnosis.
Coverage Duration	5 years
Other Criteria	For GIST: documentation of a PDGFRA exon 18 mutation or PDGFRA D842V mutation. For AdvSM, documentation of platelet count greater than 50 X 10-9th/L.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

BALVERSA

Products Affected

• BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior treatments, and outcome
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

BANZEL

Products Affected

• rufinamide

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by a specialist appropriate to the disease state such as a neurologist.
Coverage Duration	5 years
Other Criteria	For Lennox-Gastaut Syndrome: documenation of treatment with valproate and lamotrigine with outcomes (treatment failure or intolerance).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

BENLYSTA

Products Affected

• BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	Severe active lupus nephritis, active central nervous, use in combination with other biologics
Required Medical Information	Diagnosis, autoantibody testing, prior treatments including response.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a rheumatologist or nephrologist.
Coverage Duration	3 years
Other Criteria	Failed to demonstrate adequate response to TWO standard therapies at recommended doses: corticosteroids, antimalarials, NSAIDs, and/or immunosuppressants.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

BESREMI

Products Affected

• BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior treatments, and outcome
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Must have an intolerance, contraindication, or treatment failure with hydroxyurea and Peginterferon alpha-2a.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

BOSULIF

Products Affected

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to bosutinib. CBC and LFT lab test results are needed for continuation treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	Trial of OR intolerance/contraindication to imatinib.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

BRAFTOVI

Products Affected

• BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

BRIVIACT

Products Affected

• BRIVIACT ORAL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used and result of prior therapy. If continuation, response to brivaracetam.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by neurologist.
Coverage Duration	3 years
Other Criteria	Failure of treatment with levetiracetam and ONE additional Part D formulary anticonvulsant drug.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





BRONCHITOL

Products Affected

• BRONCHITOL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Patient has a diagnosis of cystic fibrosis (CF)
Age Restrictions	Individual is 18 years of age or older
Prescriber Restrictions	Must be prescribed by pulmonologist.
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





BRUKINSA

Products Affected

• BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies, and result of prior therapy.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by hematologist/oncologist.
Coverage Duration	5 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





CABOMETYX

Products Affected

CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





CALQUENCE

Products Affected

• CALQUENCE

• CALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
Exclusion Criteria	NA
Required Medical Information	Documentation of diagnosis, previous treatments, and response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by oncology
Coverage Duration	5 years
Other Criteria	Patient must have a trial of either Brukinsa or Imbruvica prior to approval of Calquence.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





CAPRELSA

Products Affected

• CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	HISTORY OF CONGENITAL LONG QT SYNDROME
Required Medical Information	DIAGNOSIS, PRIOR THERAPIES, TREATMENT RESPONSE
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST
Coverage Duration	5 years
Other Criteria	ECG, ELECTROLYTE(K,Mg,Ca), AND TSH MONITORING AT BASELINE, 2-4 WEEKS AND 8-12 WEEKS AFTER STARTING TREATMENT AND EVERY 3 MONTHS THEREAFTER
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





COMETRIQ

Products Affected

COMETRIQ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





COPIKTRA

Products Affected

COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or hematologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





CORLANOR

Products Affected

• CORLANOR ORAL SOLUTION

• ivabradine

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For heart failure in adult patients, only: ejection fraction less than or equal to 35% AND heart rate greater than 70 beats per minute AND in sinus rhythm AND on maximally tolerated beta-blocker OR has contraindication to beta-blocker (i.e., allergy, severe COPD limiting beta blocker usage).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a cardiologist.
Coverage Duration	12 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

COSENTYX

Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 150 MG/ML, 75 MG/0.5 ML
- COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other treatments tried and reasons for failure. Regular monitoring for TB required, both at baseline and during treatment
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribing limited to a rheumatologist or dermatologist.
Coverage Duration	3 years
Other Criteria	For arthritis conditions, must have a trial of or contraindication to one non-biologic DMARD (methotrexate, hydroxychloroquine, sulfasalazine, azathioprine). For diagnosis of plaque psoriasis, must have a trial of or contraindication to cyclosporine and methotrexate. For hidradenitis suppurativa: patient has tried ONE other therapy (e.g., intralesional or oral corticosteroids, systemic antibiotics, isotretinoin). The allowable quantity is dependent upon the induction dosing regimen for the applicable FDA-labeled indications as outlined in product labeling. Note: every two-week dosing requires demonstration of treatment failure of every four-week dosing.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





PA Criteria	Criteria Details
Part B Prerequisite	No





COTELLIC

Products Affected

• COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and/or failed, and treatment response.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by an oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

CROFELEMER

Products Affected

• MYTESI

PA Criteria	Criteria Details
Exclusion Criteria	Use when infectious diarrhea has not been ruled out
Required Medical Information	Diagnosis, use of antiretroviral therapy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	Infectious diarrhea needs to be ruled out prior to initiating treatment. Patient must have a history of using at least two prior treatments for diarrhea, including bismuth subsalicylate, kaolin, loperamide, or diphenoxylate/atropine.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

CYSTADROP

Products Affected

CYSTADROPS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to cysteamine.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Presribing limited to opthalmologist or corneal specialist.
Coverage Duration	1 year
Other Criteria	To start treatment, documentation of presence of corneal cystine crystal accumulation by slit lamp examination, baseline Corneal Cystine Cystal Score (CCCS) provided. For continuation: positive response to therapy (e.g., documentation showing improvement in vision with less pain and photophobia).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





DAURISMO

Products Affected

• DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies, and result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or hematologist
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

DAYBUE

Products Affected

• DAYBUE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of Rett disorder.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

DEFERIPRONE

Products Affected

• deferiprone

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, length of therapy, serum ferritin concentrations and dose/weight verification, & CBC
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Must have a contraindication to, or an inadequate response to, or have experienced clinically significant adverse effects to deferasirox.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





DEMSER

Products Affected

• metyrosine

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and whether the patient is a candidate for surgery
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





DIACOMIT

Products Affected

• DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	There is no clinical data to support the use of Diacomit alone to treat Dravet syndrome.
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to stiripentol.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a neurologist.
Coverage Duration	5 years
Other Criteria	Documentation must show co-administration of stiripentol with clobazam.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





DIFICID

Products Affected

• DIFICID ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior treatments, response to therapy
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by or in consultation with an infectious disease specialist
Coverage Duration	1 month
Other Criteria	History of failure, contraindication, or intolerance to oral Vancocin (vancomycin) capsules or vancomycin oral solution
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

DOPTELET

Products Affected

• DOPTELET (10 TAB PACK)

• DOPTELET (30 TAB PACK)

• DOPTELET (15 TAB PACK)

• DOFTELET (13 TAB PACK)	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and platelet count.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a specialist appropriate to the disease state such as a hematologist, oncologist, or gastroenterologist.
Coverage Duration	1 month for chronic liver disease, 6 months for chronic immune thrombocytopenia.
Other Criteria	For chronic liver disease-associated thrombocytpenia, the patient must be scheduled to undergo a pre-planned medical or dental procedure with treatment beginning 10 to 13 days prior to the scheduled procedure. Patients should undergo the procedure 5 to 8 days after the last dose.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





DRIZALMA

Products Affected

• DRIZALMA SPRINKLE ORAL CAPSULE, DELAYED REL SPRINKLE 20 MG, 30 MG, 40 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR TREATMENTS, RESPONSE TO THERAPY
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	Need reason why Duloxetine capsule cannot be used.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

DRONABINOL

Products Affected

• dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	Not covered for the treatment of pain.
Required Medical Information	Diagnosis, previous treatments, and the outcome.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For cancer related weight loss, must have a treatment failure or intolerance to megestrol.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

DUPIXENT

Products Affected

- DUPIXENT PEN SUBCUTANEOUS PEN INJECTOR 200 MG/1.14 ML, 300 MG/2 ML
- DUPIXENT SYRINGE SUBCUTANEOUS SYRINGE 100 MG/0.67 ML, 200 MG/1.14 ML, 300 MG/2 ML

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used and result of prior therapy. If continuation, prior response to dupilumab.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribing limited to an allergist, immunologist, pulmonologist, otolaryngologist, dermatologist or gastroenterologist.
Coverage Duration	3 years
Other Criteria	For atopic dermatitis: a trial on at least one topical corticosteroid (fluticasone, fluocinonide, desonide) - and - one at least one topical immunomodulator (tacrolimus, pimecrolimus). Immunomodulators will not be required for patients under 2 years of age. For chronic rhinosinusitis with nasal polyps (CRS with NP): documentation of inflammatory persistence for 12 weeks or longer and a trial of of a intranasal corticosteroId product (fluticasone, mometasone). For asthma: blood and/or tissue eosinophil testing with documentation of a trial of an interleukine drug (montelukast, zafirlukast). For COPD: Member has tried treatment with both of the following preferred combination therapies. LABA plus LAMA (e.g., Stiolto, tiotropium), AND ICS plus LABA (e.g., Wixela, budesonide/formoterol, Breo Ellipta). Must have a baseline eosinophil count of 300 cells/mcL or greater within previous 12 months.





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	Must have a documented postbronchodilator FEV1 of 30 to 70% of the predicted normal value.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





DUVYZAT

Products Affected

• DUVYZAT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. Medical records supporting the request including genetically confirmed diagnosis of Duchenne muscular dystrophy (DMD). Baseline function tests (e.g. time to wheelchair assistance, required respiratory assistance/pulmonary function tests, 4SC, 6MWT, time to walk/run 10 meters [10MWT], or NSAA) are documented.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	prescribed by or in consultation with a specialist (e.g. neurologist) with experience in the treatment of DMD.
Coverage Duration	1 year.
Other Criteria	Patient is ambulatory upon initiation of therapy. Patient is stable on baseline corticosteroids for 6 months.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ENBREL

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION
- ENBREL SUBCUTANEOUS SYRINGE
- ENBREL SURECLICK

SOLUTION	
PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with biologic therapy or targeted synthetic DMARD
Required Medical Information	Diagnosis, concurrent medications, previous therapies tried.
Age Restrictions	N/A
Prescriber Restrictions	RA/AS/JIA/JRA, prescribed by or in consult w/ rheumatologist. PsA, prescribed by or in consultation w/ rheumatologist or dermatologist. PP, prescribed by or in consult w/ dermatologist.GVHD, prescribed by or in consult w/ oncologist, hematologist, or physician affiliated w/ transplant center.Behcet's disease, prescribed by or in consult w/ rheumatologist, dermatologist, ophthalmologist, gastroenterologist, or neurologist. Uveitis, prescribed by or in consultation with an ophthalmologist.
Coverage Duration	3 year
Other Criteria	RA initial, patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to step back and try a conventional synthetic DMARD). JIA/JRA, approve if the pt has aggressive disease, as determined by the prescriber, or the pt has tried one other agent for this condition (eg, MTX, sulfasalazine, leflunomide, NSAID, biologic DMARD or the pt will be started on Enbrel concurrently with MTX, sulfasalazine, or leflunomide or the pt has an absolute contraindication to MTX (eg, pregnancy, breast feeding, alcoholic liver disease, immunodeficiency syndrome, blood dyscrasias), sulfasalazine, or leflunomide.Plaque psoriasis





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	(PP) initial approve if the patient meets one of the following conditions: 1) patient has tried at least one traditional systemic agent for at least 3 months for plaque psoriasis, unless intolerant (eg, MTX, cyclosporine, Soriatane, oral methoxsalen plus PUVA, (note: pts who have already tried a biologic for psoriasis are not required to step back and try a traditional agent first) OR 2) the patient has a contraindication to one oral agent for psoriasis such as MTX. GVHD. Tried or currently is receiving with etanercept 1 conventional GVHD tx (high-dose systemic corticosteroid, CSA, tacrolimus, MM, thalidomide, antithymocyte globulin, etc.). Behcet's. Has tried at least 1 conventional tx (eg, systemic corticosteroid, immunosuppressant, interferon alfa, MM, etc) or adalimumab or infliximab. Uveitis-tried one of the following: periocular, intraocular, or systemic coricosteroid, immunosuppressives, Humira or an infliximab product.RA/AS/JIA/PP/PsA Cont - must have a response to tx according to the prescriber. Behcet's, GVHD, Uveitis Cont-if the patient has had a response to tx according to the prescriber. Clinical criteria incorporated into the Enbrel 25 mg quantity limit edit, approve additional quantity (to allow for 50 mg twice weekly dosing) if one of the following is met: 1) Patient has plaque psoriasis, OR 2) Patient has RA/JIA/PsA/AS and is started and stabilized on 50 mg twice weekly dosing, OR 3) Patient has RA and the dose is being increased to 50 mg twice weekly for at least 2 months, unless MTX is contraindicated or intolerant, OR 4) Patient has JIA/PsA/AS and the dose is being increased to 50 mg twice weekly after taking 50 mg once weekly for at least 2 months.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ENDARI

Products Affected

• glutamine (sickle cell)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Patient has a diagnosis of sickle cell disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Patients must have an inadequate response, contraindication, or intolerance to hydroxyurea.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ENSPRYNG

Products Affected

ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to satralizumab-mwge.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by a neurologist or ophthalmologist.
Coverage Duration	3 years
Other Criteria	Confirmed diagnosis of neuromyelitis optica spectrum disorder (NMOSD) with a positive serologic test for anti-aquaporin-4 immunoglobulin G (AQP4-IgG)/NMO-IgG antibodies. Documented treatment failure with immunosuppressive therapy: corticosteroid therapy and mycophenolate.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

EPIDIOLEX

Products Affected

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by or in consultation with a neurologist.
Coverage Duration	5 years
Other Criteria	For Lennox-Gastaut Syndrome: documentation of use of valproate and lamotrigine and outcomes (treatment failure or intolerance). For Dravet Syndrome: documentation of use of valproate and topiramate and outcomes (treatment failure or intolerance). For refractory seizures: documentation of use of two different anti-convulsant drugs from different pharmacologic classes (valproate, topiramate, lamotrigine or similar) and outcomes (intolerance or treatment failure).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





EPRONTIA

Products Affected

• EPRONTIA

PA Criteria	Criteria Details
1 A CHICHA	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior treatments, and outcome
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	5 Years
Other Criteria	Must have a intolerance, contraindication, or medical reason the tablet or capsule are not acceptable.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ERGOTAMINE DERIVATIVES

Products Affected

• dihydroergotamine nasal

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior treatments and responses.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by a headache specialist, pain management specialist or neurologist.
Coverage Duration	1 year
Other Criteria	For treatment of migraine, unless contraindicated, a trial and failure of two different triptans (covered on the formulary): one oral tablet and one other formulation, either nasal spray or injection.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ERIVEDGE

Products Affected

• ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	PATIENTS WHO ARE CANDIDATES FOR SURGERY OR RADIATION
Required Medical Information	DIAGNOSIS, PRIOR TREATMENTS, RESPONSE TO THERAPY
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY A ONCOLOGIST OR DERMATOLOGIST
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ERLEADA

Products Affected

• ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and outcome, fall risk assessment, and seizure history (if any)
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by an oncologist or urologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ERYTHROPOIESIS STIMULATING AGENTS

Products Affected

- ARANESP (IN POLYSORBATE)
 INJECTION SOLUTION 100 MCG/ML,
 200 MCG/ML, 25 MCG/ML, 40
 MCG/ML, 60 MCG/ML
- ARANESP (IN POLYSORBATE) INJECTION SYRINGE
- EPOGEN INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/2 ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML

INJECTION SYRINGE	
PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension. Pure red cell aplasia that begins after ESA treatment.
Required Medical Information	Pre-treatment hemoglobin level less than 10 g/dL AND Patient has adequate iron stores prior to initiation of therapy defined as ferritin more than 100 mcg/L or serum transferrin saturation greater than 20% AND other causes of anemia such as iron deficiency, folate deficiency or B12 deficiency, hemolysis, gastrointestinal bleeding, other active or occult bleeding, or underlying hematologic disease (such as sickle cell anemia, thalassemia, and porphyria) have been ruled out.
Age Restrictions	N/A
Prescriber Restrictions	CKD - prescribed by a nephrologist or hematologist. Non-myeloid malignancies - prescribed by an oncologist/hematologist. Surgery - Prescribed by a surgeon. HIV - Prescribed by an infectious disease specialist.
Coverage Duration	Initial: 3 months. Renewal: CKD-12 months, Non-myeloid cancers, HIV-4 months. Surgery-3 months
Other Criteria	For renewal of CKD, for dialysis patients: Hb less than 11 g/dL or physician will decrease or interrupt dose and for non-dialysis patients: Hb less than 10 g/dL or physician will decrease or interrupt dose. For renewal of non-myeloid malignancies: Concurrent myelosuppressive chemotherapy and Hb is 12g/dL or less and there is measurable response after eight weeks. For renewal of zidovudine-treated HIV, Hb is 12g/dL or less AND





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	Zidovudine dose remains 4,200 mg/week or less and there is a measurable response after eight weeks (defined as an increase in Hb or a reduction in RBC transfusion requirements or documented dose escalation [up to max of 300 units/kg/dose])
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

EVRYSDI

Products Affected

• EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. If continuation, prior response to risdiplam.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by a pulmonologist, neurologist, orthopedist or gastroenterologist.
Coverage Duration	1 year
Other Criteria	Genetic testing to determine if the SMN1 gene is missing or damaged is required to initiate SMA treatment.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

FASENRA

Products Affected

• FASENRA

• FASENRA PEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Allergen test, baseline FEV1, FEV1 following bronchodilator, asthma medical history (including medications, emergency department visits, and hospitalizations), baseline eosinophil count
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by a Pulmonologist or Immunologist or Allergist
Coverage Duration	3 year
Other Criteria	Patient must be diagnosed with severe asthma, currently receiving inhaled and/or oral corticosteroid treatment, AND have a baseline eosinophil count of 150 cells/mcL or greater within previous 12 months.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

FILSUVEZ

Products Affected

• FILSUVEZ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of Epidermolysis Bullosa
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	n/a
Coverage Duration	1 year.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





FINTEPLA

Products Affected

• FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to fenfluramine.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a neurologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

FORTEO

Products Affected

- teriparatide subcutaneous pen injector 20 mcg/dose (600mcg/2.4ml)
 - TERIPARATIDE SUBCUTANEOUS PEN INJECTOR 20 MCG/DOSE (620MCG/2.48ML)

PA Criteria	Criteria Details
Exclusion Criteria	Not approved for a cumulative lifetime duration of abaloparatide and any other parathyroid hormone therapy (eg, teriparatide) of more than 2 years. Not approved for combination therapy of a PTH/PTHrP analog in combination with other osteoporosis agents.
Required Medical Information	Diagnosis, fracture history, prior therapy used and response to prior therapy. Required pretreatment testing: DXA, if not performed in the past two years: serum calcium, phosphorus, creatinine, alkaline phosphatase, albumin, 25-hydroxyvitamin D (25[OH]D), and, 24-hour urine calcium, creatinine (or fasting specimen for calcium/creatine ratio) to evaluate for baseline hypercalciuria.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an endocrinologist.
Coverage Duration	2 years
Other Criteria	Documentation of a trial on an oral bisphosphonate, or, if GI intolerant of oral bisphosphonates, use of a parenteral bisphosphonate - AND - a trial on denosumab.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
	All Medically-accepted Indications.





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Part B Prerequisite	Yes





Date Effective: 12/01/2024

FOTIVDA

Products Affected

• FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, response to tivozanib.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a specialist in hematology / oncology.
Coverage Duration	5 years
Other Criteria	Documentation of two prior lines of systemic drug therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





FRUZAQLA

Products Affected

• FRUZAQLA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of Colorectal cancer, Metastatic, previously treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, an anti-VEGF therapy, and an anti-EGFR therapy if RAS wild-type and medically appropriate
Age Restrictions	Member is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





FYCOMPA

Products Affected

• FYCOMPA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried, response to prior therapy
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be prescribed by neurologist
Coverage Duration	5 years
Other Criteria	Monitor at initiation and after dose increases for serious psychiatric and/or behavioral reactions.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





GALAFOLD

Products Affected

GALAFOLD

PA Criteria	Criteria Details
Exclusion Criteria	Not covered in combination with Fabrazyme
Required Medical Information	Confirmed diagnosis of Fabry disease and baseline renal function assessment
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a specialist in genetic disorders, or nephrologist
Coverage Duration	3 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

GATTEX

Products Affected

• GATTEX 30-VIAL

• GATTEX ONE-VIAL

PA Criteria	Criteria Details
Exclusion Criteria	Therapy should be discontinued in cases of intestinal malignancy.
Required Medical Information	Diagnosis, other therapies tried and treatment responses.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a gastroenterologist.
Coverage Duration	3 year
Other Criteria	For ADULT patients: A colonoscopy of the entire colon with removal of polyps must be done before initiating therapy, medical records documenting this procedure must be submitted. For PEDIATRIC patients: Perform fecal occult blood testing: if there is unexplained blood in the stool, perform colonoscopy / sigmoidoscopy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

GAUCHER'S DISEASE TREATMENT

Products Affected

CERDELGA

• miglustat

PA Criteria	Criteria Details
Exclusion Criteria	NOT APPROVED FOR TYPE II OR TYPE III GAUCHER'S DISEASE
Required Medical Information	Diagnosis. For MIGLUSTAT: rationale as to why ERT is not appropriate.
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY GENETICIST, HEMOTOLOGIST, OR METABOLIC SPECIALIST
Coverage Duration	3 YEAR
Other Criteria	USE OF MIGLUSTAT IS RESERVED FOR THOSE WHOM ENZYME REPLACEMENT THERAPY IS NOT AN OPTION
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





GAVRETO

Products Affected

GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to pralsetinib.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by an oncologist or pulmonologist.
Coverage Duration	5 years
Other Criteria	Diagnosis of non-small cell lung cancer (NSCLC) or Thyroid cancer that is verified by an FDA-approved diagnostic test to have rearranged during transfection (RET) fusion mutations.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





GILOTRIF

Products Affected

• GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies tried and/or failed
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





GLEOSTINE

Products Affected

• GLEOSTINE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous therapies tried and/or failed
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

GLP1

Products Affected

MOUNJARO

• OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2

MG/3 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)

- RYBELSUS
- TRULICITY

PA Criteria	Criteria Details
Exclusion Criteria	Not covered for weight loss
Required Medical Information	Documented diagnosis of type 2 diabetes and submitted A1c from past 6 months.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Member has tried Metformin or has a medical contraindication or intolerance to Metformin.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

GRALISE

Products Affected

• gabapentin oral tablet extended release 24 • GRALISE ORAL TABLET EXTENDED

RELEASE 24 HR 450 MG, 750 MG, 900 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and previous treatments, including dosage and outcome of previous treatments.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Must have a documented intolerance, contraindication to, or failure of generic regular relase gabapentin titrated to maximum tolerated dosage or rationale as to why generic regular release gabapentin cannot be used.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





GROWTH HORMONE

Products Affected

• NORDITROPIN FLEXPRO

• NUTROPIN AQ NUSPIN

PA Criteria	Criteria Details
Exclusion Criteria	PRESENCE OF CONTRAINDICATIONS TO THERAPY
Required Medical Information	DIAGNOSIS, HEIGHT AND WEIGHT, HISTORY OF GROWTH MEASUREMENT. REPLACEMENT THERAPY IN PATIENTS WITH GROWTH HORMONE DEFICIENCY WITH DIAGNOSIS CONFIRMED BY APPROPRIATE GROWTH HORMONE STIMULATION TESTING
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an an endocrinologist or nephrologist.
Coverage Duration	3 years
Other Criteria	Replacement therapy in patients with growth hormone deficiency with diagnosis confirmed by appropriate growth hormone stimulation testing.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

HADLIMA

Products Affected

- ADALIMUMAB-FKJP SUBCUTANEOUS PEN INJECTOR KIT • HADLIMA PUSHTOUCH
- ADALIMUMAB-FKJP SUBCUTANEOUS SYRINGE KIT
- HADLIMA
- HADLIMA(CF)
- HADLIMA(CF) PUSHTOUCH

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another biologic DMARD or targeted synthetic DMARD.
Required Medical Information	Diagnosis, concurrent medications, previous therapies tried.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	RA/JIA/JRA/Ankylosing spondylitis, prescribed by or in consultation with rheumatologist. Psoriatic arthritis (PsA), prescribed by or in consultation with a rheumatologist or dermatologist. Plaque psoriasis (PP), prescribed by or in consultation with a dermatologist. UC/CD, prescribed by or in consultation with a gastroenterologist. HS - dermatologist. UV-ophthalmologist
Coverage Duration	3 years
Other Criteria	RA: patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to 'step back' and try a conventional synthetic DMARD). JIA/JRA: patient has tried another a non-biologic DMARD (e.g., MTX, sulfasalazine, leflunomide) - or will be starting on adalimumab concurrently with MTX, sulfasalazine, or leflunomide. Approve without trying another agent if pt has absolute contraindication to MTX, sulfasalazine, or leflunomide or if pt has aggressive disease. PP: approve if the patient meets one of the following criteria: 1) pt has tried at least one





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	traditional systemic agent (e.g., MTX, cyclosporine, acitretin, PUVA) for at least 3 months, unless intolerant (note: patients who have already tried a biologic for psoriasis are not required to 'step back' and try a traditional agent first) OR 2) patient has a contraindication to MTX as determined by the prescribing physician. CD: patient has tried corticosteroids (CSs), or if CSs are contraindicated, or if pt currently on CSs, or patient has tried one other agent for CD (e.g., azathioprine, 6-mercaptopurine, MTX, OR patient had ilecolonic resection or enterocutaneous (perianal or abdominal) or rectovaginal fistulas. UC: patient has tried a systemic therapy (e.g., 6-mercaptopurine, azathioprine, mesalamine or a corticosteroid such as prednisone or methylprednisolone) for 2 months or was intolerant to one of these agents, or the patient has pouchitis and has tried therapy with an antibiotic, probiotic, corticosteroid enema, or mesalamine (Rowasa) enema. HS: patient has tried ONE other therapy (e.g., intralesional or oral corticosteroids, systemic antibiotics, isotretinoin). Clinical criteria incorporated into the Hadlima 40 mg quantity limit edit allow for approval of additional quantities to accommodate induction dosing. The allowable quantity is dependent upon the induction dosing regimen for the applicable FDA-labeled indications as outlined in product labeling. For all indications (excepting hidradenitis suppurativa), weekly dosing requires a demonstration of treatment failure of every other week dosing.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

HEMADY

Products Affected

• HEMADY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to dexamethasone.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Presribing limited to hematologist / oncologist.
Coverage Duration	5 years
Other Criteria	Clinical treatment plan to include combination therapy of dexamethasone with other anti-myeloma products. Dexamethasone dosing is to be in accordance with the Prescribing Information of the other anti-myeloma products used in the combination treatment.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





HEPATITIS C TREATMENT

Products Affected

• MAVYRET ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	PATIENT WEIGHT, GENOTYPE, HCV-RNA, LEVEL OF FIBROSIS, TREATMENT HISTORY
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY GASTROENTEROLOGIST, HEPATOLOGIST, OR INFECTIOUS DISEASE SPECIALIST
Coverage Duration	8 - 24 WEEKS. TREATMENT WILL BE APPROVED CONSISTENT WITH CURRENT AASLD-IDSA GUIDANCE.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





HEREDITARY ANGIOEDEMA

Products Affected

- HAEGARDA
- icatibant
- ORLADEYO

- sajazir
- TAKHZYRO

PA Criteria	Criteria Details
Exclusion Criteria	Dual prescribing of injectable and oral formulations for HAE prophylaxis are not covered to prevent risk of double-dosing.
Required Medical Information	Diagnosis and the results of immunologic laboratory testing that show low C4 and functional C1- inhibitor levels (less than the lower limits of laboratory reference ranges).
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by an allergist or immunologist.
Coverage Duration	1 year
Other Criteria	For icatibant initiation: Member is 18 years of age and older, with confirmed diagnosis of HAE. To continue icatibant in patients who have treated previous acute HAE attacks with icatibant: documentation demonstrating a favorable clinical response (e.g., decrease in the duration of HAE attacks, quick onset of symptom relief, complete resolution of symptoms, decrease in HAE acute attack frequency or severity). To initiate berotralstat (Orladeyo: Member is aged 12 or older. Confirmed diagnosis of HAE. For those 18 years or older, documented use and effectiveness of on-demand treatment with icatibant (Firazyr) and that adding prophylaxis treatment is appropriate to the care plan. To initiate C1 esterase inhibitor, (Haegrada): Member is aged 6 or older. Confirmed diagnosis of HAE. For those 18 years or older, documented use and effectiveness of on-demand treatment with icatibant (Firazyr) and that adding prophylaxis treatment is appropriate to the care plan. To initiate lanadelumab-flyo (Takhzyro):): Member is aged 2 or older. Confirmed diagnosis of HAE. For those 18





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	years or older, documented use and effectiveness of on-demand treatment with icatibant (Firazyr) and that adding prophylaxis treatment is appropriate to the care plan.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





HETLIOZ

Products Affected

• tasimelteon

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies and responses.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by a sleep specialist or neurologist.
Coverage Duration	1 year
Other Criteria	For a diagnosis of non-24-hour sleep-wake disorder, submit sleep log through a wrist activity monitor that supports diagnosis of non-24-hour sleep-wake disorder) AND sleep study has ruled out sleep apnea and periodic limb movement disorder. For continuation, positive clinical response demonstrated by: (1) increased total nighttime sleep and, (2) decreased daytime nap duration, as determined by treating physician. For nighttime sleep disturbances in Smith-Magenis Syndrome, documentation supporting the diagnosis.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA

Products Affected

• JUXTAPID

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, results of prior therapy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 year
Other Criteria	Patient must have a diagnosis of homozygous familial hypercholesterolemia. Liver function tests required at baseline and at least monthly during the first year of treatment.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

HUMIRA

Products Affected

• HUMIRA(CF) (ONLY NDCS STARTING WITH 00074)

SUBCUTANEOUS SYRINGE KIT 10 MG/0.1 ML

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with another biologic DMARD or targeted synthetic DMARD.
Required Medical Information	Diagnosis, concurrent medications, previous therapies tried.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	RA/JIA/JRA/Ankylosing spondylitis, prescribed by or in consultation with rheumatologist. Psoriatic arthritis (PsA), prescribed by or in consultation with a rheumatologist or dermatologist. Plaque psoriasis (PP), prescribed by or in consultation with a dermatologist. UC/CD, prescribed by or in consultation with a gastroenterologist. HS - dermatologist. UV-ophthalmologist
Coverage Duration	3 years
Other Criteria	RA: patient has tried one conventional synthetic DMARD for at least 3 months (note: patients who have already had a 3-month trial of a biologic for RA are not required to 'step back' and try a conventional synthetic DMARD). JIA/JRA: patient has tried another a non-biologic DMARD (e.g., MTX, sulfasalazine, leflunomide, NSAID) - or - biologic DMARD (e.g., etanercept, abatacept, infliximab, anakinra, tocilizumab) or will be starting on adalimumab concurrently with MTX, sulfasalazine, or leflunomide. Approve without trying another agent if pt has absolute contraindication to MTX, sulfasalazine, or leflunomide or if pt has aggressive disease. PP: approve if the patient meets one of the following criteria: 1) pt has tried at least one traditional systemic agent (e.g., MTX,





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	cyclosporine, acitretin, PUVA) for at least 3 months, unless intolerant (note: patients who have already tried a biologic for psoriasis are not required to 'step back' and try a traditional agent first) OR 2) patient has a contraindication to MTX as determined by the prescribing physician. CD: patient has tried corticosteroids (CSs), or if CSs are contraindicated, or if pt currently on CSs, or patient has tried one other agent for CD (e.g., azathioprine, 6-mercaptopurine, MTX, certolizumab, infliximab, ustekinumab, or vedolizumab) OR patient had ilecolonic resection or enterocutaneous (perianal or abdominal) or rectovaginal fistulas. UC: patient has tried a systemic therapy (e.g., 6-mercaptopurine, azathioprine, CSA, tacrolimus, infliximab, golimumab SC, or a corticosteroid such as prednisone or methylprednisolone) for 2 months or was intolerant to one of these agents, or the patient has pouchitis and has tried therapy with an antibiotic, probiotic, corticosteroid enema, or mesalamine (Rowasa) enema. HS: patient has tried ONE other therapy (e.g., intralesional or oral corticosteroids, systemic antibiotics, isotretinoin). Clinical criteria incorporated into the Humira 40 mg quantity limit edit allow for approval of additional quantities to accommodate induction dosing. The allowable quantity is dependent upon the induction dosing regimen for the applicable FDA-labeled indications as outlined in product labeling. For all indications (excepting hidradenitis suppurativa), weekly dosing requires demonstration of treatment failure of every other week dosing.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





IBRANCE (S)

Products Affected

• IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ICLUSIG

Products Affected

• ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or hematologist
Coverage Duration	5 years
Other Criteria	Liver function monitoring required at baseline and 3 months after initiation
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





IDHIFA

Products Affected

• IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous treatments, and outcome.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist/oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

IMBRUVICA

Products Affected

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION

• IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR TREATMENTS, RESPONSE TO THERAPY
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be prescribed by oncologist, hematologist, or transplant specialist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





INBRIJA

Products Affected

• INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to LEVODOPA ORAL INHALATION.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a neurologist.
Coverage Duration	1 year
Other Criteria	Required trial and failure of: 1) both carbidopa/levodopa IR and ER and, 2) at least one other Parkinson's Disease drug: entacapone, pramipexole, ropinirole, selegiline, rasagiline, or amantadine. Intention to continue use of carbidopa/levodopa.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

INCRELEX

Products Affected

• INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, HEIGHT AND WEIGHT MEASUREMENTS, GH LEVEL, IGF-1 LEVEL
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ENDOCRINOLOGIST
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





INHALED TOBRAMYCIN

Products Affected

• TOBI PODHALER

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, therapies tried, and outcome
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by Infectious disease specialist or pulmonologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





INLYTA

Products Affected

• INLYTA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR TREATMENTS, RESPONSE TO THERAPY
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY A ONCOLOGIST
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





INQOVI

Products Affected

• INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to decitabine / cedazuridine
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a hematologist / oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

INREBIC

Products Affected

• INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	Evaluate baseline thiamine levels prior to treatment initiation, do not initiate fedratinib in patients with thiamine deficiency. Replete thiamine prior to fedratinib initiation and during treatment if thiamine levels are low.
Required Medical Information	Diagnosis, Other therapies tried with treatment response, baseline thiamine level, baseline platelet level
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by Hematologist / Oncologist
Coverage Duration	5 years
Other Criteria	Documented baseline platelet count of at least 50,000 per cubic milimeter.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

INTERFERON ALPHA

Products Affected

• PEGASYS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	FOR HEPATITIS C: PATIENT WEIGHT, GENOTYPE, HCV-RNA QUANTITY AND DATE OF TEST, PRESENCE OF DIRRHOSIS (Y/N), TREATMENT HISTORY, HISTORY OF ANEMIA OR DEPRESSION. HEPATITIS B: HBEAG STATUS, HBV DNA QUANTITY, AND ALT LEVEL. OTHERS: DIAGNOSIS, OTHER THERAPIES TRIED AND FAILED, RESPONSE TO TREATMENT
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	2 YEARS FOR INDICATIONS OTHER THAN HEPATITIS C. HEPC APPROVALS FROM 12-48 WKS BASED ON DRUG REGIMEN
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

INVEGA IM

Products Affected

- INVEGA HAFYERA
- INVEGA SUSTENNA

• INVEGA TRINZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and other treatments tried and outcome.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a psychiatrist or mental health specialist.
Coverage Duration	5 years
Other Criteria	Patient must have a reason oral paliperidone cannot be used.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





IRESSA

Products Affected

• gefitinib

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, response to prior therapy
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST
Coverage Duration	5 years
Other Criteria	GEFITINIB IS COVERED AS MONOTHERAPY
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





IVERMECTIN

Products Affected

• ivermectin oral

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Must be prescribed by an infectious disease specialist or dermatologist.
Coverage Duration	1 month
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

IVIG

Products Affected

- BIVIGAM
- FLEBOGAMMA DIF INTRAVENOUS SOLUTION 10 %
- GAMMAGARD LIQUID
- GAMMAGARD S-D (IGA < 1 MCG/ML) PRIVIGEN
- GAMMAKED

- GAMMAPLEX
- GAMMAPLEX (WITH SORBITOL)
- GAMUNEX-C
- OCTAGAM

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and response to treatments.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Diagnosis and administration information will be reviewed to determine if coverage is available as a Medicare Part B or Part D benefit. Part B before Part D Step Therapy: SCIG will be reserved for members who cannot use IVIG due to poor access (on going access site issues unresolved by traditional means) - or - SCIG will be reserved for patients who continue to experience infusion reactions despite documented infusion rate adjustments and adequate pre-treatment. For Idiopathic thrombocytopenia purpura (ITP): trial and failure of oral corticosteroids at therapeutic dose (standard dosage of prednisone is 1 mg/kg/day) required. For Chronic Inflammatory Demyelinating Polyneuropathy (CIDP): trial and failure of oral corticosteroids at therapeutic dose (standard dosage of prednisone is 1-1.5 mg/kg/day) required.





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





IWILFIN

Products Affected

• IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





JAKAFI

Products Affected

JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, OTHER TREATMENTS TRIED AND FAILED, CBC AT BASELINE AND PERIODICALLY AFTER INITIATION, HISTORY OF RBC TRANSFUSIONS
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a hematologist, oncologist or transplant specialist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





JAYPIRCA

Products Affected

JAYPIRCA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis, previous treatments, and response to treatment.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by hematologist/oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





JOENJA

Products Affected

JOENJA

PA Criteria	Criteria Details
PA Criteria	Criteria Details
Exclusion Criteria	FOR TREATMENT OF ACTIVATED PHOSPHOINOSITIDE 3 KINASE DELTA SYNDROME (APDS): CANNOT BE USED IN COMBINATION WITH AN IMMUNOSUPPRESSIVE MEDICATION
Required Medical Information	COVERAGE FOR ACTIVATED PHOSPHOINOSITIDE 3 KINASE DELTA SYNDROME (APDS) REQUIRES ALL OF THE FOLLOWING: 1. A DIAGNOSIS OF APDS WITH AN ASSOCIATED PI3K DELTA MUTATION, 2. DOCUMENTED VARIANT IN EITHER PIK3CD OR PIK3R1, AND 3. DOCUMENTED SYMPTOMS ASSOCIATED WITH APDS SUCH AS NODAL AND/OR EXTRANODAL LYMPHOPROLIFERATION, HISTORY OF REPEATED OTO-SINO-PULMONARY INFECTIONS AND/OR ORGAN DYSFUNCTION (E.G. LUNG, LIVER).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





JYNARQUE

Products Affected

• JYNARQUE ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	History of signs or symptoms of significant liver impairment or injury
Required Medical Information	Documented diagnosis of polycystic kidney disease, ultrasound results
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by endocrinology or nephrology
Coverage Duration	6 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

KALYDECO

Products Affected

• KALYDECO ORAL GRANULES IN PACKET 13.4 MG, 25 MG, 5.8 MG, 50 MG, 75 MG

• KALYDECO ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and the presence of one or more specific gene mutations that the drug is FDA approved to treat.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





KERENDIA

Products Affected

KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	Not approved for serum potassium greater than 5.0 mEq/L.
Required Medical Information	Diagnosis, prior drug treatments and outcomes. Potassium level within 30 days.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by an endocrinologist, nephrologist or cardiologist.
Coverage Duration	1 year
Other Criteria	INITIATION: Documented diagnosis of chronic kidney disease (CKD) associated with diabetes mellitus, Type 2 (T2D). Documentation in the medical record that the patient is currently receiving the following standard of care background therapy with the requested agent: (a) a maximally tolerated dose of angiotensin converting enzyme (ACE) inhibitor, angiotensin receptor blocker (ARB), or a combination medication containing an ACE inhibitor or ARB therapy - AND - (b) an antidiabetic agent (e.g., metformin or an agent containing metformin, SGLT2 inhibitor, GLP-1) - OR - (c) according to the prescriber, the patient has contraindications to both ACE and ARB drug therapy. At baseline (prior to initiation of finerenone), (a) an estimated glomerular filtration rate greater than or equal to 25 mL/min/1.73m2 AND - (b) serum potassium level less than or equal to 5.0 mEq/L. CONTINUATION: Serum potassium within 30 days. Demonstrated response of GFR with finerenone therapy.
Indications	All Medically-accepted Indications.





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No





KINERET

Products Affected

KINERET

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with biologic therapy
Required Medical Information	Diagnosis, other therapies tried and/or failed
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by rheumatologist or pediatrician
Coverage Duration	3 year
Other Criteria	RA initial, Trial (3 month) and failure one formulary anti-TNF agents (adalimumab, etanercept, infliximab), or medically valid rationale as to why anti-TNF agents cannot be used (e.g. congestive heart failure (NYHA class III/IV) with an ejection fraction less than or equal to 50%). Trial (3 month) and failure to Rinvoq or medically valid rationale to avoid.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

KISQALI

Products Affected

KISQALI

• KISQALI FEMARA CO-PACK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies previously tried, and the outcome.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

KORLYM

Products Affected

• mifepristone oral tablet 300 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and outcome, HbA1c
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by endocrinologist
Coverage Duration	1 year
Other Criteria	Must have trial of ketoconazole therapy or have intolerance or contraindication to these medications. Must have failed surgery or not be a candidate for surgery. For continuation of therapy patient must show an improvement in HbA1c.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





KOSELUGO

Products Affected

KOSELUGO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by oncology
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





KRAZATI

Products Affected

• KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, diagnostic testing for muations, prior drug treatments and outcomes.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by an oncologist.
Coverage Duration	5 years
Other Criteria	Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC), as determined by an FDA-approved test, who have received at least one prior systemic therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





KUVAN

Products Affected

• sapropterin

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. BASELINE and FOLLOW-UP phenylalanine levels.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	2 MONTHS FOR INITIATION, 1 YEAR FOR CONTINUATION
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





LAZCLUZE

Products Affected

• LAZCLUZE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information
Prescriber Restrictions	Prescribed by or in consultation with an oncologist/hematologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





LENVIMA

Products Affected

• LENVIMA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





LEUKINE

Products Affected

• LEUKINE INJECTION RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

LIBTAYO

Products Affected

• LIBTAYO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and/or failed, and response to prior therapy
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by or in consultation with an oncologist or hematologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

LIDOCAINE TRANSDERMAL

Products Affected

- asperflex (lidocaine) topical cream
- lidocaine topical adhesive patch,medicated 5 %
- LIDOCAINE TOPICAL CREAM
- LMX 4 TOPICAL CREAM

- LMX 5
- rectasmoothe
- RECTICARE
- ULTRA LIDO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

LOKELMA

Products Affected

LOKELMA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of hyperkalemia, confirmed with laboratory test within the past month.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For initiation, must have a failure, contraindication, or intolerance to sodium polystyrene sulfonate (SPS). For continuation, must show response to therapy as demonstrated by normal potassium levels and patient remains at high risk for recurrence of hyperkalemia.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

LONSURF

Products Affected

• LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and/or failed, and treatment response
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

LORBRENA

Products Affected

LORBRENA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

LUMAKRAS

Products Affected

• LUMAKRAS ORAL TABLET 120 MG, 320 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, diagnostic testing for muations, prior drug treatments and outcomes.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by an oncologist.
Coverage Duration	5 years
Other Criteria	Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC), as determined by an FDA-approved test, who have received at least one prior systemic therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





LYBALVI

Products Affected

• LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	Not approved for dementia-related psychosis. Not approved for patients using opioids. Not approved for patients undergoing acute opioid withdrawal.
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Psychiatrist or in consulation with psychiatrist.
Coverage Duration	5 years
Other Criteria	Demonstrated positive clinical response but with unacceptable weight gain while on single-agent olanzapine AND trial/failure of one other formulary atypical anti-psychotic (e.g., risperidone, aripiprazole, quetiapine, ziprasidone) titrated to maximum tolerated dose. Rationale for combination therapy in medical record. Patient does not have a known opioid use disorder nor is dependent on opioids for a chronic health condition. Prior to initiating LYBALVI when prescription history shows opioid fills within the last 30 days, prescriber attestation required to initiate olanzapine/samidorphine (LYBALVI): 7-day opioid-free interval from the last use of short-acting opioids, and at least a 14-day opioid-free interval from the last use of long-acting opioids to avoid precipitation of opioid withdrawal.
Indications	All Medically-accepted Indications.





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No





LYNPARZA

Products Affected

LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR THERAPIES, GENETIC TESTING
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





LYTGOBI

Products Affected

 LYTGOBI ORAL TABLET 12 MG/DAY (4 MG X 3), 16 MG/DAY (4 MG X 4), 20 MG/DAY (4 MG X 5)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis, previous treatments, and response to treatment.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by hematologist/oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

MAVACAMTEN

Products Affected

• CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a cardiologist
Coverage Duration	3 years
Other Criteria	Must have an intolerance, contraindication, or treatment with at least one Non-vasodilating beta-blocker (e.g. metoprolol, propranolol or atenolol) AND one Non-dihydropyridine calcium channel blocker (e.g. verapamil, diltiazem)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

MEKINIST

Products Affected

• MEKINIST ORAL RECON SOLN

• MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	Wild-type BRAF melanoma
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. Documentation of BRAF mutation, as detected using an FDA-approved test.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

MEKTOVI

Products Affected

• MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	Must be used in combination with encorafenib.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





METHAMPHETAMINE

Products Affected

• methamphetamine

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies previously tried and failed, and response to treatment
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For the treatment of attention deficit disorder patient must have a trial of both methylphenidate and amphetamine/dextroamphetamine or rationale as to why these treatments are not suitable.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

MIRVASO

Products Affected

• brimonidine topical

D. C.	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and other treatments tried and outcome.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For the treatment of acne rosacea: doxycycline (oral) and topical metronidazole.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





MULPLETA

Products Affected

• MULPLETA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and platelet count
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a specialist appropriate to the disease state such as a hematologist, oncologist, or gastroenterologist.
Coverage Duration	1 month
Other Criteria	Patient must be scheduled to undergo a pre-planned medical or dental procedure with treatment beginning 8 to 14 days prior to the scheduled procedure. Patients should undergo the procedure 2 to 8 days after the last dose.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

MULTIPLE SCLEROSIS

Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT
- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)

- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)
- MAYZENT
- MAYZENT STARTER(FOR 1MG MAINT)
- MAYZENT STARTER(FOR 2MG MAINT)
- PLEGRIDY

• WAVENCEAD (/ TABLETTACK)	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For MS diagnosis: EDSS score, relapse history, physical or cognitive disability, TB test.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by a specialist appropriate to the disease state such as a neurologist.
Coverage Duration	3 year
Other Criteria	Trial of two of the following glatiramer/glatopa, Dimethyl Fumarate, Fingolimod, and Terifluomide.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

MU-OPIOID RECEPTOR ANTAGONIST.

Products Affected

• SYMPROIC

PA Criteria	Criteria Details
Exclusion Criteria	Will not be approved for cancer related pain
Required Medical Information	Documented diagnosis of opiate induced constipation (non-cancer pain)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Patient must have a trial of or contraindication to at least two different classes of laxative agents including bulk, osmotic, or stimulant laxatives.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





MYALEPT

Products Affected

• MYALEPT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





MYFEMBREE

Products Affected

• MYFEMBREE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Medically accepted indication
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	Trial of 2 oral contraceptives
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





NAYZILAM

Products Affected

NAYZILAM

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to midazolam.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a neurologist.
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





NERLYNX

Products Affected

NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, previous treatments, and outcome.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





NEXLETOL

Products Affected

NEXLETOL

NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to bempedoic acid (with or without ezetimibe).
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by or given in consultation with a cardiologist, endocrinologist, or physician who focuses on CV risk management and or lipid disorders.
Coverage Duration	3 years
Other Criteria	For Primary hyperlipidemia: LDL of 190 or higher (not associated with ASCVD, HeFH, or HoFH), approve if tried one high-intensity statin therapy (i.e., atorvastatin 40-80 mg daily, rosuvastatin 20-40 mg daily) and ezetimibe for at least 8 weeks and LDL remains 100 mg/dL or higher, unless statin intolerant. For Cardiovascular event risk, Statin intolerant with cardiovascular disease (CVD), or a high risk for a CVD event: History of atherosclerotic cardiovascular disease (ASCVD) or high risk of cardiovascular event. OR Established diagnosis of heterozygous familial hypercholesterolemia (HeFH) OR atherosclerotic cardiovascular disease (ASCVD) with history of ONE of the following (for ASCVD): Myocardial infarction (MI) OR Acute Coronary Syndrome (ACS) OR Stable or unstable angina OR Thromboembolic stroke OR Transient ischemic attack (TIA) OR Peripheral arterial disease (PAD) OR Coronary or other arterial revascularization. Patient has had a previous trial of or has a contraindication to a high-intensity statin (i.e., atorvastatin 40-80 mg daily, rosuvastatin 20-40 mg daily) and LDL-C remains greater than 100 mg/dL





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	with laboratory confirmation within the last 30 days. If patient cannot tolerate a high intensity statin, the patient is taking a maximally tolerated dose of any statin. If intolerate of all statins, patient has had a trial of ezetimbe, unless contraindicated, and LDL-C remains greater than 100 mg/dL with laboratory confirmation within the last 30 days. Patient will continue taking the maximally tolerated statin (unless contraindicated) in combination with Nexletol or Nexlizet.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





NINLARO

Products Affected

• NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and/or failed, and treatment response. Used in combination with Revlimid (lenalidomide) and dexamethasone.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or hematologist
Coverage Duration	5 years
Other Criteria	Must have an intolerance or contraindication to Velcade
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	Yes





Date Effective: 12/01/2024

NOXAFIL

Products Affected

• NOXAFIL ORAL SUSP, DELAYED RELEASE FOR RECON

• posaconazole oral

RELEASE FOR RECON	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, therapies tried, and outcome
Age Restrictions	N/A
Prescriber Restrictions	Prescribed Infectious disease specialist, transplant specialist, hematologist, or oncologist
Coverage Duration	6 months
Other Criteria	For the treatment of aspergillosis patient must have failure of, intolerance or contraindication to vorconizole OR rationale as to why vorconizole is not suitable.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





NUBEQA

Products Affected

• NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to darolutamide.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or urologist.
Coverage Duration	5 year
Other Criteria	If metastatic disease patient must have a failure, intolerance, or contraindication to abiraterone (Zytiga)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





NUCALA

Products Affected

• CINQAIR

NUCALA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For patients with asthma: allergen test, baseline FEV1, FEV1 following bronchodilator, asthma medical history (including medications, emergency department visits, and hospitalizations), baseline eosinophil count. For patients with eosinophilic granulomatosis with polyangiitis: documentation of diagnosis, prior therapies, and the outcome.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a: pulmonologist, immunologist, allergist, rheumatologist, hematologist or otolaryngologist.
Coverage Duration	3 years
Other Criteria	For patients with asthma: must be currently receiving inhaled and/or oral corticosteroid treatment, AND have a baseline eosinophil count of 300 cells/mcL or greater within previous 12 months or 150 cell/mcL within previous 6 weeks.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





NUEDEXTA

Products Affected

NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by or in consultation with a neurologist, psychiatrist or geriatrician.
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





NUPLAZID

Products Affected

NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Parkinson's disease psychosis: Diagnosis of Parkinson's disease. Patient has at least one of the following: hallucinations or delusions.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by or in consultation with a neurologist or psychiatrist
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

NURTEC

Products Affected

• NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	For acute migraine: Unless contraindicated per the FDA label, a trial of at least one-month a triptan. For chronic migraine: Patient must have an inadequate response, contraindication, or intolerance to two different chronic migraine prevention drugs. The two prerequisite drugs must be from different classes such as anticonvulsants (topiramate/valproate), beta blockers (propranolol, metoprolol), and antidepressants (nortriptyline/venlafaxine).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

OCALIVA

Products Affected

• OCALIVA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and failed, and response to therapy, baseline alkaline phosphatase (ALP) level for initiation, and ALP levels after first 3 months of therapy and then yearly for continuation of therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Use in combination with ursodiol in patients with an inadequate biochemical response to treatment (elevated ALP levels) with ursodiol dosed at 13-15 mg/kg/day for at least 1 year, may be used as monotherapy in patients unable to tolerate ursodiol. Must show improvement in ALP levels for continuation.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ODOMZO

Products Affected

ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Patient has recurring disease following surgery or radiation OR patient is not a candidate for surgery or radiation therapy
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or dermatologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





OGSIVEO

Products Affected

• OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





OJEMDA

Products Affected

 OJEMDA ORAL SUSPENSION FOR
 OJEMDA ORAL TABLET RECONSTITUTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, Documentation of BRAF mutation or BRAF fusion or rearrangement amd liver function tests (ALT, AST, bilirubin) at initiation.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information
Prescriber Restrictions	Prescription must be written by an oncologist or hematologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





OJJAARA

Products Affected

OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. Members must have Myelofibrosis, Intermediate or high risk, primary or secondary, with anemia.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	prescribed by an Oncologist or hematologist.
Coverage Duration	5 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ONUREG

Products Affected

ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	Do not substitute ONUREG for intravenous or subcutaneous azacitidine.
Required Medical Information	N/A
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by an oncologist or hematologist oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ORENCIA

Products Affected

• ORENCIA CLICKJECT

• ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other treatments tried and reasons for failure. Regular monitoring for TB required, both at baseline and during treatment.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY A RHEUMATOLOGIST
Coverage Duration	3 year
Other Criteria	For arthritic conditions, a 3 month trial at least one non-biologic DMARD (methotrexate, hydroxychloroquine, sulfasalazine, azathioprine) and documented reason for failure (or contraindication).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ORFADIN

Products Affected

• nitisinone

• ORFADIN ORAL SUSPENSION

	T
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS AND WEIGHT
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY AN ENDOCRINOLOGIST, GASTROENTEROLOGIST, HEMATOLOGIST, METABOLIC SPECIALIST, OR NEPHROLOGIST
Coverage Duration	6 months
Other Criteria	CLOSE MONITORING OF DISEASE MARKERS (ERYTHROCYTE PBG-SYNTHASE ACTIVITY, URINE 5-ALA, SUCCINYLACETONE) DURING THE FIRST 3 MONTHS OF TREATMENT TO ENSURE NORMALIZATION
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ORGOVYX

Products Affected

ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to relugolix.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Presribing limited to oncologist or urologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

ORKAMBI

Products Affected

 ORKAMBI ORAL GRANULES IN PACKET 100-125 MG, 150-188 MG, 75-94 MG • ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Patient has a diagnosis of cystic fibrosis (CF) AND Patient is homozygous for the F508del mutation in the CF transmembrane conductance regulator (CFTR) gene AND The presence of the mutation was documented by an FDA-cleared cystic fibrosis mutation test.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 years
Other Criteria	For renewal, Patient is benefiting from treatment (i.e., improvement in lung function [forced expiratory volume in one second [FEV1], decreased number of pulmonary exacerbations)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ORLISSA

Products Affected

• ORILISSA

DA Cuitoris	Cuitouio Dotoila
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies, and result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Must have a failure, contraindication, or intolerance to a continuous hormonal contraceptive AND progestin therapy (e.g. medroxyprogesterone, norethindrone).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ORSERDU

Products Affected

• ORSERDU

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis, previous treatments, and response to treatment.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by hematologist/oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

OTEZLA

Products Affected

• OTEZLA ORAL TABLET 20 MG, 30 MG

MG (51), 10 MG (4)-20 MG (4)-30 MG (47)

• OTEZLA STARTER ORAL TABLETS, DOSE PACK 10 MG (4)- 20

1ABLE15,DOSE PACK 10 MG (4)- 20	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by a Rheumatologist or Dermatologist
Coverage Duration	3 year
Other Criteria	For arthritis conditions, must have a trial of or contraindication to one non-biologic DMARD (including but not limited to methotrexate, hydroxychloroquine, sulfasalazine, azathioprine). For diagnosis of plaque psoriasis, must have a trial of or contraindication to cyclosporine or methotrexate
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





OXERVATE

Products Affected

OXERVATE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies, and result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an ophthalmologist
Coverage Duration	8 weeks
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

PALYNZIQ

Products Affected

PALYNZIQ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior treatments, and outcome. Baseline and follow up phenylalanine (Phe) concentrations.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Must have phenylalanine (Phe) concentrations greater than 600 micromol/L on existing management, and a failure, contraindication, or intolerance to Kuvan.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





PANRETIN

Products Affected

• PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





PDE-5 INHIBITORS FOR PAH

Products Affected

- alyq
- sildenafil (pulmonary arterial hypertension) oral suspension for reconstitution 10 mg/ml
- sildenafil (pulmonary arterial hypertension) oral tablet 20 mg
- tadalafil (pulm. hypertension)

reconstitution 10 mg/mi	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, right heart cath results.
Age Restrictions	N/A
Prescriber Restrictions	For PAH: prescribed by, or in consultation with, a cardiologist or a pulmonologist.
Coverage Duration	3 years
Other Criteria	Pulmonary arterial hypertension (PAH) WHO Group 1, are required to have had a right-heart catheterization to confirm diagnosis of PAH to ensure appropriate medical assessment. Sildenafil must be tried prior to the use of the other drugs included in these criteria, unless using ambrisentan (Letairis) plus tadalafil for treatment-naive, WHO functional class II or III PAH.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

PEMAZYRE

Products Affected

• PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to pemigatinib.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by an oncologist or gastroenterologist.
Coverage Duration	5 years
Other Criteria	Documentation of a susceptible fibroblast growth factor receptor 2 fusion or other genetic rearrangement (as detected by an FDA-approved test).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





PIQRAY

Products Affected

• PIQRAY

D. C.	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	Breast Cancer. Approve if the patient meets the following criteria (A, B, C, D, and E): A)The patient is a postmenopausal female or a male AND B)The patient has advanced or metastatic hormone receptor (HR)-positive disease AND C)The patient has human epidermal growth factor receptor 2 (HER2)-negative disease AND D)The patient has PIK3CA-mutated breast cancer as detected by an approved test AND E)The patient has progressed on or after at least one prior endocrine-based regimen (e.g., anastrozole, letrozole, exemestane, Faslodex, tamoxifen, toremifene).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





POMALYST

Products Affected

• POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or hematologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





PROLIA

Products Affected

• PROLIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed or in consultation with Endocrinologist, Hematologist/Oncologist, Obstetrician/Gynecologist, Rheumatologist or Urologist
Coverage Duration	3 years
Other Criteria	For treatment of postmenopausal osteoporosis / treatment of osteoporosis in men [a man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression], approve if the patient meets one of the following: (1) has had inadequate response after 6 months of therapy with an oral bisphosphonate or (2) had osteoporotic fracture or fragility fracture while receiving an oral bisphosphonate or (3) the patient cannot take an oral bisphosphonate because (s)he cannot swallow or has difficulty swallowing, cannot remain in an upright position, or has a pre-existing GI medical condition - AND - the patient has tried an IV bisphosphonate (e.g., ibandronate or zoledronic acid). Part B before Part D Step Therapy. Approve if the patient has severe chronic kidney disease (e.g., creatinine clearance less than 35 mL/min). Approve for treatment of bone loss in patient at high risk for fracture receiving ADT for nonmetastatic prostate cancer or receiving adjuvant AI therapy for breast cancer. For treatment of glucocorticoid





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	induced osteoporosis (GIO), approve if the patient has tried one oral bisphosphonate OR patient cannot take an oral bisphosphonate because the patient cannot swallow or has difficulty swallowing, or the patient cannot remain in an upright position post oral bisphosphonate administration, or has a pre-existing GI medical condition (e.g., esophageal lesions, esophageal ulcers, or abnormalities of the esophagus that delay esophageal emptying [stricture, achalasia]), OR patient has tried zoledronic acid (Reclast), OR patient has severe renal impairment (CrCL less than 35 mL/min) or has CKD or has had an osteoporotic fracture or fragility fracture.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	Yes





Date Effective: 12/01/2024

PROMACTA

Products Affected

PROMACTA

PA Criteria	Criteria Details
Exclusion Criteria	For ITP, eltrombopag should only be used if the degree of thrombocytopenia and clinical condition increase the risk for bleeding. For chronic hepatitis C, eltrombopag should only be used if the degree of thrombocytopenia prevents initiation of or limits the ability to maintain interferon-based therapy. Eltrombopag is not indicated for the treatment of myelodysplastic syndromes.
Required Medical Information	Diagnosis, other therapies tried and outcome. Platelet count.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

PULMONARY HYPERTENSION

Products Affected

- ambrisentan
- bosentan
- OPSUMIT
- ORENITRAM
- TRACLEER ORAL TABLET FOR SUSPENSION
- TYVASO
- TYVASO INSTITUTIONAL START KIT
- TYVASO REFILL KIT
- TYVASO STARTER KIT
- VENTAVIS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Confirmation of diagnosis of pulmonary arterial hypertension (WHO GROUP 1). Prior therapies used and responses to treatments.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a pulmonologist or cardiologist.
Coverage Duration	3 years
Other Criteria	PAH (WHO Group 1) diagnosis confirmed by a right heart catheterization to ensure appropriate medical assessment. For new starts in the Treatment Naive patient: must initiate treatment with dual therapy with tadalafil and ambrisentan, unless intolerant or contraindicated. For these drugs: bosentan, macitentan, treprostnil: must show documentation of prior drug treatment. Note: patients who are already established on any therapy and clinically responsive and stable, a step-back to dual therapy is not required.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





PA Criteria	Criteria Details
Part B Prerequisite	No





Date Effective: 12/01/2024

PURIXAN

Products Affected

• PURIXAN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR TREATMENTS, RESPONSE TO THERAPY
Age Restrictions	N/A
Prescriber Restrictions	prescriber must be a oncologist or hematologist
Coverage Duration	5 years
Other Criteria	Need reason why Mercaptopurine tablet cannot be used.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

PYRUKYND

Products Affected

PYRUKYND

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a hemoatologist or oncologist
Coverage Duration	3 months
Other Criteria	Continuation of therapy is dependant on response to therapy, determined by treating physician.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





QINLOCK

Products Affected

• QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to ripretinib.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a oncologist or gastroenterologist.
Coverage Duration	5 years
Other Criteria	For initiation, documentation of prior treatment with three or more kinase inhibitors, including imatinib.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

RADICAVA

Products Affected

• RADICAVA ORS

• RADICAVA ORS STARTER KIT SUSP

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to therapy
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by a neurologist or prescriber expertise with treating ALS
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





REGRANEX

Products Affected

REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prior use of collagenase (Santyl), unless contraindicated.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

REPATHA

Products Affected

REPATHA

• REPATHA SURECLICK

• REPATHA PUSHTRONEX

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use of Juxtapid or Praluent.
Required Medical Information	Current LDL-C (within 30 days of request), documentation of prior statin drug(s) and ezetimibe previously tried including dosage and response to therapy such as adverse event history (for example muscle pain) and/or inadequate reduction of LDL-C (provide lab value).
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by, or in consultation with, a cardiologist, endocrinologist, or a physician who focuses in the treatment of CV risk management and/or lipid disorders
Coverage Duration	Approve for 3 years
Other Criteria	Hyperlipidemia with HeFH - approve if: (1) diagnosis of HeFH - AND - (2) tried ezetimibe and ONE high intensity statin (e.g., atorvastatin greater than or equal to 40 mg daily or rosuvastatin greater than or equal to 20 mg daily) and LDL remains 70 mg/dL or higher, unless patient is statin intolerant defined by experiencing statin-related rhabdomyolysis or skeletal-related muscle symptoms while receiving separate trials of atorvastatin and rosuvastatin, and during both trials the symptoms resolved upon discontinuation. For hyperlipidemia with ASCVD, approve if: (1) has one of the following conditions: prior MI, h/o ACS, diagnosis of angina, h/o CVA or TIA, PAD, undergone a coronary or other arterial revascularization procedure - AND - (2) tried ezetimibe and ONE high intensity statin (defined above) and LDL remains 70 mg/dL or higher, unless patient is statin intolerant (defined above). For HoFH, approve if: (1) has one of the following: (a) genetic confirmation of two mutant alleles





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	at the LDLR, APOB, PCSK9, or LDLRAP1 gene locus, OR (b) untreated LDL greater than 500 mg/dL (prior to treatment), OR (c) treated LDL greater than or equal to 300 mg/dL (after treatment but prior to agents such as Repatha or Juxtapid), OR (d) has clinical manifestations of HoFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas or xanthelasma), AND - (2) tried ezetimibe and ONE high intensity statin (defined above) for 8 weeks or longer and LDL remains 70 mg/dL or higher unless statin intolerant (defined above). For primary hyperlipidemia with LDL of 190 or higher (not associated with ASCVD, HeFH, or HoFH), approve if tried one high-intensity statin therapy (defined above) and ezetimibe for at least 8 weeks and LDL remains 100 mg/dL or higher, unless statin intolerant (defined above). Maximally tolerated statin therapy may mean zero tolerance for those patients who cannot tolerate a statin.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

RETEVMO

Products Affected

• RETEVMO ORAL CAPSULE

• RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to selpercatinib.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a oncologist, pulmonologist or endocrinologist.
Coverage Duration	5 years
Other Criteria	For initiation, documentation of the presence of a RET gene fusion (with non-small cell lung cancer or thyroid cancer) or specific RET gene mutation (medullary thyroid cancer) in tumor specimens or plasma.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





REVCOVI

Products Affected

• REVCOVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documented diagnosis of adenosine deaminase (ADA) deficiency, IgA, IgM and IgG levels, CBC, and the prescence of mutations in the ADA gene at 20q13.11
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an immunologist
Coverage Duration	3 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





REVLIMID

Products Affected

• lenalidomide

	C. de de Detelle
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and previous therapies or drug regimens tried.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	MCL-approve if the patient meets one of the following 1) Pt has tried two prior therapies or therapeutic regimens OR 2) Pt has tried one prior therapy or therapeutic regimen and cannot take Velcade according to the prescribing physician. MDS-approve if the patient meets one of the following: 1) Pt has symptomatic anemia, OR 2) Pt has transfusion-dependent anemia, OR 3) Pt has anemia that is not controlled with an erythroid stimulating agent (eg, Epogen, Procrit [epoetin alfa injection], Aranesp [darbepoetin alfa injection]). Diffuse, Large B Cell Lymphoma (Non-Hodgkin's Lymphoma)-approve if the pt has tried one other medication treatment regimen. Myelofibrosis-approve if the pt has tried one other therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Part B Prerequisite	No





Date Effective: 12/01/2024

REYVOW

Products Affected

• REYVOW

PA Criteria	Criteria Details
Exclusion Criteria	Excluded for migraine prevention.
Required Medical Information	N/A
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by a headache specialist, pain management specialist or neurologist.
Coverage Duration	1 year
Other Criteria	Unless contraindicated per the FDA label, a trial of at least one-month of two different triptans: one oral tablet and one other formulation, either nasal spray or injection.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





REZLIDHIA

Products Affected

• REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis, previous treatments, and response to treatment.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by hematologist/oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

REZUROCK

Products Affected

REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR TREATMENTS, RESPONSE TO THERAPY
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be prescribed by oncologist, hematologist, or transplant specialist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Bate Effective: 12/01/202

RINVOQ

Products Affected

• RINVOQ

• RINVOQ LQ

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a rheumatologist, dermatologist, or gastroenterologist
Coverage Duration	For Ulcerative Colitis, 8 weeks for 45 mg daily. 3 years for all other indications
Other Criteria	Prior to receiving treatment with Rinvoq for arthritis related conditions, the patient must have trial and failure of at least one non-biologic DMARDs, including but not limited to methotrexate, hydroxychloroquine and sulfasalazine, azathioprine for at least three months or have a contraindication. For atopic dermatitis: a trial on at least one topical corticosteroid (fluticasone, fluocinonide, desonide) - and - one at least one topical immunomodulator (tacrolimus, pimecrolimus). For inflammatory bowel disease (CD and UC), the patient must have an inadequate response or intolerance to a tumor necrosis factor blockers. Dose adjustments and laboratory monitoring parameters should be adhered to as specified in the FDA approved drug labeling.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





PA Criteria	Criteria Details
Part B Prerequisite	No





Date Effective: 12/01/2024

ROZLYTREK

Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG
- ROZLYTREK ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, If continuation, prior response to ENTRECTINIB.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist.
Coverage Duration	5 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





RUBRACA

Products Affected

• RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	OVARIAN CANCER (epithelial ovarian, fallopian tube or primary peritoneal): Diagnosis of advanced ovarian cancer. Presence of deleterious BRCA mutation as detected by an FDA-approved diagnostic test. Patients are in a complete or partial response to platinum-based chemotherapy. Maintenance Therapy-Approve if the patient is in complete or partial response after at least two platinum-based chemotherapy regimens. PROSTATE CANCER: Diagnosis of advanced metastatic castration-resistant prostate cancer. Presence of deleterious BRCA mutation as detected by an FDA-approved diagnostic test. History of failure, contraindication, or intolerance to androgen receptor-directed therapy and a taxane-based chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





RYDAPT

Products Affected

• RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Newly diagnosed acute myeloid leukemia (AML), FMS-like tyrosine kinase 3 (FLT3) mutation-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test, used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation. Aggressive Systemic Mastocytosis (ASM), Systemic Mastocytosis with Associated Hematological Neoplasm (SM-AHN), Mast Cell Leukemia (MCL): Diagnosis of one of the following: aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL).
Age Restrictions	N/A
Prescriber Restrictions	All indications: Prescribed by or in consultation with a hematologist or oncologist.
Coverage Duration	5 years
Other Criteria	Approve for continuation of prior therapy
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

SAMSCA

Products Affected

• TOLVAPTAN ORAL TABLET 15 MG • tolvaptan oral tablet 30 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, patient has a serum sodium less than 125 mEq/L at baseline, OR member has less marked hyponatremia (serum sodium less than 135 mEq/L at baseline) AND is symptomatic (e.g. nausea, vomiting, headache, lethargy, confusion, and baseline LFTs).
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a specialist appropriate to the disease state sucha as an endocrinologist or nephrologist.
Coverage Duration	1 month
Other Criteria	Patient must have documented failure of two other therapies (e.g. fluid restriction, furosemide, demeclocycline).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





SCEMBLIX

Products Affected

• SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. Medical records supporting the request.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

SIGNIFOR

Products Affected

SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis and other treatments tried and failed. Documentation: of surgery with response (if performed), or when surgery is not a treatment option.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an endocrinologist
Coverage Duration	6 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

SIRTURO

Products Affected

• SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and/or failed. ECG and liver function tests are required at baseline and at intervals as specified in the FDA prescribing information to monitor for potentially severe adverse events.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a infectious disease specialist
Coverage Duration	24 weeks
Other Criteria	Drug therapy must be directly observed. Use of Sirturo is reserved for MDR-TB where other treatment options cannot be used for safety or efficacy reasons.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

SKYCLARYS

Products Affected

SKYCLARYS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis and Confirmation of diagnosis via genetic testing revealing two pathogenic mutations of the frataxin (FXN) gene. Obtain alanine aminotransferase (ALT), aspartate aminotransferase (AST), bilirubin, B-type natriuretic peptide (BNP), and lipid parameters.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by or in consultation with a Neurologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

SKYRIZI

Products Affected

- SKYRIZI SUBCUTANEOUS PEN INJECTOR
- SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML
- SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR 180 MG/1.2 ML (150 MG/ML), 360 MG/2.4 ML (150 MG/ML)

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by a gastroenterologist, dermatologist, or rheumatologist
Coverage Duration	3 years
Other Criteria	For diagnosis of plaque psoriasis, must have a trial of or contraindication to cyclosporine or methotrexate. For arthritis conditions, must have a trial of or contraindication to one non-biologic DMARD (methotrexate, hydroxychloroquine, sulfasalazine, azathioprine). For UC: patient has tried a systemic therapy (e.g., 6-mercaptopurine, azathioprine, CSA, or a corticosteroid such as prednisone or methylprednisolone) for 2 months or was intolerant to one of these agents, or the patient has pouchitis and has tried therapy with an antibiotic, probiotic, corticosteroid enema, or mesalamine (Rowasa) enema. For Crohns disease: patient has tried corticosteroids (CSs), or if CSs are contraindicated, or if pt currently on CSs, or patient has tried one other agent for CD (e.g., azathioprine, 6-mercaptopurine, MTX, OR patient had ilecolonic resection or enterocutaneous (perianal or abdominal) or rectovaginal fistulas.





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

SPRITAM

Products Affected

• SPRITAM

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, therapies tried, and outcome
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by Neurologist
Coverage Duration	5 years
Other Criteria	Rationale as to why generic levetiracetam is not suitable.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





SPRYCEL

Products Affected

• dasatinib

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST OR HEMATOLOGIST
Coverage Duration	5 years
Other Criteria	Must have a failure, intolerance or contraindication to imatinib.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

STELARA

Products Affected

- STELARA SUBCUTANEOUS SOLUTION
- STELARA SUBCUTANEOUS SYRINGE 45 MG/0.5 ML, 90 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies, and result of prior therapy. TB test results
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by gastroenterologist, dermatologist or rheumatologist.
Coverage Duration	3 year
Other Criteria	For arthritis conditions, must have a trial of or contraindication to one non-biologic DMARD (including but not limited to methotrexate, hydroxychloroquine, sulfasalazine, azathioprine). For diagnosis of plaque psoriasis, must have a trial of or contraindication to cyclosporine or methotrexate. For psoriasis and psoriatic arthritis, 90 mg dosing requires demonstration of treatment failure of 45 mg for 16 weeks and patient weight greater than 100kg. For inflammatory bowel disease (CD and UC), must have trial of or contraindication to at least one non-biologic conventional therapy, including but not limited to sulfasalazine, mesalamine, azathioprine, or methotrexate.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





PA Criteria	Criteria Details
Part B Prerequisite	No





STIVARGA

Products Affected

• STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to regorafenib. LFT lab test results are needed for continuation treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

SUNOSI

Products Affected

• SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior drug treatments and outcomes.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a pulmonologist, sleep specialist or neurologist.
Coverage Duration	1 year
Other Criteria	INITIATION: 1) Demonstration that co-existing hypertension has been controlled before initiating treatment with solriamfetol. 2) A one-month trial of modafinil and armodafinil, both titrated to maximum tolerated dose.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





SUTENT

Products Affected

• sunitinib malate

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR THERAPIES, AND TREATMENT RESPONSE. DOCUMENATION OF FAILURE OF IMATINIB FOR PATIENTS WITH GIST. DOCUMENTATION OF FAILURE OF VOTRIENT FOR PATIENTS WITH RENAL CELL CARCINOMA. IF NO PREVIOUS FAILURE, RATIONALE AS TO WHY PREFERRED AGENT CANNOT BE USED
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST
Coverage Duration	5 years
Other Criteria	GIST PATIENTS REQUIRE A FOLLOW-UP CT SCAN BETWEEN 8 AND 12 WEEKS.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

SYMDEKO

Products Affected

SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of CF AND homozygous for the F508del mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor based on in vitro data and/or clinical evidence. If the patients genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of a CFTR mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use. FEV1 at baseline and continuation, baseline LFT and continuation, review for drug interactions CYP3A inducers
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by a pulmonologist or doctor specializing in cystic fibrosis
Coverage Duration	5 year
Other Criteria	For renewal, Patient is benefiting from treatment (i.e., improvement in lung function [forced expiratory volume in one second (FEV1)], decreased number of pulmonary exacerbations)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





SYMLIN

Products Affected

• SYMLINPEN 120

• SYMLINPEN 60

PA Criteria	Criteria Details
Exclusion Criteria	GASTROPARESIS OR USE OF DRUGS TO STIMULATE GASTROINTESTINAL MOTILITY
Required Medical Information	HBA1C AND CURRENT DIABETES MEDICATIONS
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 YEAR
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective. 12/01/2024

SYMPAZAN

Products Affected

• SYMPAZAN ORAL FILM 10 MG, 20 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	Must have failure, intolerance, or contraindication to generic clobazam
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TABRECTA

Products Affected

• TABRECTA

DA C '	
PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to capmatinib.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by an oncologist or pulmonologist.
Coverage Duration	5 years
Other Criteria	Documentation of a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an FDA-approved test.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TAFAMIDIS

Products Affected

VYNDAMAX

VYNDAQEL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Confirmation of diagnosis with appropriate testing.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a cardiologist
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TAFINLAR

Products Affected

• TAFINLAR ORAL CAPSULE

• TAFINLAR ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	Wild-type BRAF melanoma
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. Documentation of BRAF mutation, as detected using an FDA-approved test.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TAGRISSO

Products Affected

• TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis supported with an approved test for the detection of mutations named in FDA label found in tumor or plasma specimens. Other therapies tried and responses to treatments.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by an oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TALZENNA

Products Affected

TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TARCEVA

Products Affected

• erlotinib

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR THERAPIES, AND TREATMENT RESPONSE
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TARGRETIN

Products Affected

• bexarotene topical

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, response to bexarotene.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribing limited to a specialist in dermatology, hematolgy, or oncology.
Coverage Duration	5 months
Other Criteria	Documentation that the patient has refractory or persistent cutaneous T-cell lymphoma (stage IA and IB) or who has not tolerated other therapies
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TASIGNA

Products Affected

• TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, OTHER THERAPIES TRIED AND FAILED, RESPONSE TO TREATMENT, POTENTIAL DRUG INTERACTIONS
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY AN ONCOLOGIST
Coverage Duration	5 years
Other Criteria	Must have a failure, intolerance or contraindication to imatinib and Sprycel.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TAVALISSE

Products Affected

TAVALISSE

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried, outcome, and platelet count less than 50,000/microL for at least 3 months
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by a hematologist or oncologist
Coverage Duration	1 year
Other Criteria	Patient must have a failure, contraindication, or intolerance to at least two of the following therapies: corticosteroids, IVIG, Rituxan, or Promacta. For continuation of therapy the platelet counts must be to a level sufficient to avoid clinically important bleeding.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TAVNEOS

Products Affected

TAVNEOS

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior treatments, response to therapy
Age Restrictions	N/A
Prescriber Restrictions	Prescription must written hematologist, rheumatologist, neurologist, nephrologist, and immunologist
Coverage Duration	3 years
Other Criteria	Patient must have tried and failed two of the following: azathioprine, methotrexate, mycophenolate, rituximab, or cyclophosphamide
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TAZORAC

Products Affected

• tazarotene topical cream

• tazarotene topical gel

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 YEAR
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TAZVERIK

Products Affected

TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to tazemetostat.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribing limited to a hematologist/oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TEGSEDI

Products Affected

• TEGSEDI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of polyneuropathy of hereditary transthyretin-mediated amyloidosis (hATTR). For continuation of therapy, demonstrated positive response to therapy (improved neurologic impairment, motor function, or slowing of disease progression)
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by a geneticist or neurologist
Coverage Duration	6 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TEPMETKO

Products Affected

• TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to TEPOTINIB.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by an oncologist or pulmonologist.
Coverage Duration	5 years
Other Criteria	For NSCLC: confirmation of genertic alteration with biomarker testing for mesenchymal-epithelial transition (MET) exon 14 skipping alterations.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Prior Authorization Criteria

HAP CareSourceTM MI Health Link (Medicare-Medicaid Plan) 2024

Date Effective: 12/01/2024

TESTOSTERONE

Products Affected

- testosterone transdermal gel
- testosterone transdermal gel in metereddose pump 12.5 mg/1.25 gram (1 %), 20.25 mg/1.25 gram (1.62 %)

• testosterone transdermal gel in packet

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of primary hypogonadism (congenital or acquired) in males. Diagnosis of secondary (hypogonadotropic) hypogonadism (congenital or acquired) in males. Hypogonadism (primary or secondary) in males, serum testosterone level. [Man is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.]
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Hypogonadism (primary or secondary) in males - initial therapy, approve if all of the following criteria are met: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) [eg, depressed mood, decreased energy, progressive decrease in muscle mass, osteoporosis, loss of libido, AND 2) patient has had two pre-treatment serum testosterone (total or available) measurements, each taken in the morning on two separate days, AND 3) the two serum testosterone levels are both low, as defined by the normal laboratory reference values. hypogonadism has been confirmed by a low for age serum testosterone (total or free) level defined by the normal laboratory reference values. Hypogonadism (primary or secondary) in males - continuing therapy, approve if the patient meets all





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	of the following criteria: 1) patient has persistent signs and symptoms of androgen deficiency (pre-treatment) AND 2) patient had at least one pre-treatment serum testosterone level that was low. [Note: male is defined as an individual with the biological traits of a man, regardless of the individual's gender identity or gender expression.]
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TIBSOVO

Products Affected

• TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TRANSMUCOSAL IMMEDIATE RELEASE FENTANYL

Products Affected

• fentanyl citrate buccal lozenge on a handle 1,200 mcg, 200 mcg, 600 mcg

PA Criteria	Criteria Details
Exclusion Criteria	NOT approved for patients who are NOT tolerant to opioid drug treatment - AND - are NOT receiving long-acting opioids
Required Medical Information	Diagnosis and prior drug treatments.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Documentation that shows prior use of two formulary short-acting opioid analgesics such as oxycodone, morphine, or hydromorphone as immediate-release (IR) tablet formulations or oral solution. Explanation of treatment failure or product intolerance must also explain why this unique method of administration is medically necessary and why a traditional short-acting oral opiate medication cannot be continued.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TRELSTAR DEPOT, TRELSTAR LA

Products Affected

• TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY AN ONCOLOGIST.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TRETINOIN

Products Affected

• tretinoin topical

PA Criteria	Criteria Details
Exclusion Criteria	Coverage is not provided for cosmetic use.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TRIENTINE

Products Affected

• trientine oral capsule 250 mg

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and the outcome
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist or hepatatologist.
Coverage Duration	6 months
Other Criteria	Must have a failure, contraindication, or intolerance to penicillamine.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TRUQAP

Products Affected

• TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of Breast cancer, Locally advanced or metastatic, hormone receptor-positive, HER2-negative, with one or more PIK3CA/AKT1/PTEN-alteration
Age Restrictions	Member is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TUKYSA

Products Affected

• TUKYSA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to tucatinib.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a oncologist.
Coverage Duration	5 years
Other Criteria	For initiation, documentation of human epidermal growth factor receptor 2 (HER2) testing. Documentation of one or more prior anti-HER2-based regimens in the metastatic setting.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

TURALIO

Products Affected

• TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis: symptomatic tenosynovial giant cell tumor associated with severe morbidity or functional limitations and not amenable to improvement with surgery in adults. If continuation, prior response to pexidartinib.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TYKERB

Products Affected

• lapatinib

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR AND CURRENT THERAPIES, TREATMENT RESPONSE
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





TYMLOS

Products Affected

• TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	Not approved for a cumulative lifetime duration of abaloparatide and any other parathyroid hormone therapy (eg, teriparatide) of more than 2 years. Not approved for combination therapy of a PTH/PTHrP analog in combination with other osteoporosis agents.
Required Medical Information	Diagnosis, fracture history, prior therapy used and response to prior therapy. Required pretreatment testing: DXA, if not performed in the past two years: serum calcium, phosphorus, creatinine, alkaline phosphatase, albumin, 25-hydroxyvitamin D (25[OH]D), and, 24-hour urine calcium, creatinine (or fasting specimen for calcium/creatine ratio) to evaluate for baseline hypercalciuria.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an endocrinologist.
Coverage Duration	2 years
Other Criteria	Documentation of a trial on an oral bisphosphonate, or, if GI intolerant of oral bisphosphonates, use of a parenteral bisphosphonate - AND - a trial on denosumab.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	Yes





Date Effective: 12/01/2024

UBRELVY

Products Affected

ubrelvy

PA Criteria	Criteria Details
Exclusion Criteria	Excluded for migraine prevention.
Required Medical Information	N/A
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Unless contraindicated per the FDA label, a trial of at least one-month of a triptan.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

UCERIS

Products Affected

• budesonide oral tablet,delayed and ext.release

PA Criteria	Criteria Details
Exclusion Criteria	Not approved for maintenance of remission or in patients with severe disease (UCDAI score = 10)
Required Medical Information	Diagnosis, other therapies tried and/or failed, including anti-inflammatory and immunosuppressant drugs
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a gastroenterology specialist
Coverage Duration	3 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





UPTRAVI

Products Affected

• UPTRAVI ORAL

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Confirmation of diagnosis of PAH, other therapies tried, and documentation of response to therapy
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by a cardiologist or pulmonologist
Coverage Duration	3 years
Other Criteria	Must have PAH WHO group 1. Prior to receiving treatment with Uptravi, patient must have a contraindication, intolerance to, or history of taking a PDE5 inhibitor (sildenafil or Adcirca) AND an ERA (bosentan or ambrisentan)
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

UREA CYCLE DISORDER

Products Affected

• RAVICTI

• sodium phenylbutyrate oral powder

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VALCHLOR

Products Affected

VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VALTOCO

Products Affected

• LIBERVANT

VALTOCO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Documentation of diagnosis
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribing limited to a neurologist.
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

VANFLYTA

Products Affected

VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Acute Myeloid Leukemia (AML): Newly diagnosed acute myeloid leukemia (AML), FLT3 internal tandem duplication (ITD)-positive) mutation-positive as detected by a U.S. Food and Drug Administration (FDA)-approved test, used in combination with standard cytarabine and anthracycline induction and cytarabine consolidation and as maintenance monotherapy following consolidation chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist.
Coverage Duration	5 years
Other Criteria	Approve for continuation of prior therapy
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VELTASSA

Products Affected

• VELTASSA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Evidence of episodes of moderate to severe hyperkalemia (serum potassium level = 5.1 mEq/L) requiring discontinuation or dose reduction of angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, and/or aldosterone antagonists AND receives medication regimen that allows for practical administration of Veltassa 3 hours before or 3 hours after other oral medications
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Prior use of sodium polystryrene (SPS) and sodium zirconium cyclosilicate (Lokelma).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

VENCLEXTA

Products Affected

VENCLEXTA

• VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. Other treatments tried and response to therapies.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VEOZAH

Products Affected

VEOZAH

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried, and outcome. Baseline hepatic function (including ALT, AST, and serum bilirubin [total and direct] before initiating therapy)
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	Patients must have an inadequate response, contraindication, or intolerance to two different medications such as conjugated estrogens, Venlafaxine, gabapentin, or clonidine.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VERZENIO

Products Affected

VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	NA
Required Medical Information	Documentation of diagnosis, previous treatments, response to treatment, and LFTs.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by oncology
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VITRAKVI

Products Affected

VITRAKVI

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies, and result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VIZIMPRO

Products Affected

VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VONJO

Products Affected

VONJO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to treatment
Age Restrictions	N/A
Prescriber Restrictions	Must be prescribed by oncology and hematology
Coverage Duration	3 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VORANIGO

Products Affected

VORANIGO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. Medical records supporting the request, including evidence of IDH1 or IDH2 mutation.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information
Prescriber Restrictions	Prescribed by or in consultation with an oncologist/hematologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VORICONAZOLE

Products Affected

• voriconazole intravenous

PA Criteria	Criteria Details
Exclusion Criteria	Not approved for topical use such as Foot Bath, Nasal Rinse, Mouthwash, etc. applications.
Required Medical Information	Medically accepted indication.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	N/A
Coverage Duration	1 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VOTRIENT

Products Affected

• pazopanib

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY AN ONCOLOGIST
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VOXZOGO

Products Affected

VOXZOGO

PA Criteria	Criteria Details
Exclusion Criteria	limb surgery
Required Medical Information	Documentation of achondroplasia confirmed by genetic testing for variants in the fibroblast growth factor receptor 3 (FGFR3) gene, members baseline annualized growth velocity, open epiphyses AND prescriber attests that there are no plans for the member to have limb-lengthening surgery and the member has not had limb-lengthening surgery.
Age Restrictions	5 years or older
Prescriber Restrictions	Prescribed by or in consultation with a board-certified geneticist, endocrinologist, neurologist, orthopedic surgeon, or specialist with experience in treating achondroplasia.
Coverage Duration	5 year
Other Criteria	Documentation of members positive clinical response as demonstrated by improvement in annualized growth velocity, open epiphyses AND prescriber attests that there are no plans for the member to have limblengthening surgery.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





VUITY

Products Affected

• VUITY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis
Age Restrictions	N/A
Prescriber Restrictions	Prescription must by written by an optometrist or ophthalmologist
Coverage Duration	3 years for all medically accepted indications
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

WAINUA

Products Affected

• WAINUA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of Polyneuropathy due to amyloidosis, Hereditary transthyretin-mediated.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	prescribed by a neurologist.
Coverage Duration	3 years.
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





WAKIX

Products Affected

WAKIX

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to pitolisant.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescriber must be a neurologist or sleep specialist.
Coverage Duration	1 year
Other Criteria	For narocolepsy wiithout cateplexy: an adequate trial (at least 2 months) on one analeptic drug (i.e., modafinil or armodafinil) AND one CNS stimulant drug (i.e., amphetamine, methylphenidate or amphetamine/dextroamphetamine), unless intolerant or contraindicated.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

WEIGHT LOSS

Products Affected

- CONTRAVE
- QSYMIA
- SAXENDA

- WEGOVY
- XENICAL

• SAXENDA	
Criteria Details	
N/A	
Patient must have a BMI greater than or equal to 30 kg/m2 OR Patient must have a BMI greater than or equal to than 27 kg/m2 but less than 30 kg/m2 and at least one of the following risk factors: hypertension, coronary artery disease, diabetes, dyslipidemia, or sleep apnea OR For Wegovy, pediatric patients must have an initial BMI at the 95th percentile or greater for age and sex (obesity). Prescriber attests to patient has absence of any contraindications to use of the requested product AND Prescriber attests that the patient is not pregnant or lactating AND Prescriber attests that at least one previously documented weight reduction attempt in the past year AND Prescriber attests medication therapy is part of a total treatment plan including a calorie and fat restricted diet and exercise regimen.	
Patient age greater than or equal to 12 years (Wegovy, Xenical, Saxenda) OR Patient age greater than or equal to 18 years (Contrave, Qsymia)	
N/A	
6 months	
N/A	
All Medically-accepted Indications.	
N/A	





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Part B Prerequisite	No





Date Effective. 12/01/202-

WEILREG

Products Affected

WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

WINREVAIR

Products Affected

WINREVAIR

PA Criteria	Criteria Details
Exclusion Criteria	n/a
Required Medical Information	Diagnosis. Medical records supporting the request. Confirmation of diagnosis of pulmonary arterial hypertension (WHO GROUP 1). Prior therapies used and responses to treatments.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by, or in consultation with, a cardiologist or a pulmonologist.
Coverage Duration	1 year
Other Criteria	Pulmonary arterial hypertension (PAH) WHO Group 1, are required to have had a right-heart catheterization to confirm diagnosis of PAH to ensure appropriate medical assessment. Patient is currently receiving at least two other PAH therapies from the following different pharmacologic categories: phosphodiesterase type 5 inhibitors (PDE5i), endothelin receptor antagonists (ERAs), soluble guanylate cyclase stimulator (sGCs), and prostacyclins OR Patient is currently receiving at least one other PAH therapy and is intolerant to combination therapy with a phosphodiesterase type 5 inhibitors (PDE5i), endothelin receptor antagonists (ERAs), soluble guanylate cyclase stimulator (sGCs), or prostacyclin.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Part B Prerequisite	No





Date Effective: 12/01/2024

XATMEP

Products Affected

• XATMEP

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior treatments, response to therapy
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or rheumatologist
Coverage Duration	3 years
Other Criteria	Medical justification as to why member cannot use methotrexate tablets or injectable solution
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

XCOPRI

Products Affected

XCOPRI

XCOPRI TITRATION PACK

• XCOPRI MAINTENANCE PACK

PA Criteria	Criteria Details
Exclusion Criteria	Not approved for patients with familial short QT syndrome.
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to cenobamate.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescription must be written by a neurologist.
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





XDEMVY

Products Affected

• XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For treatment of symptomatic Demodex blepharitis (DB). Symptoms defined as redness, inflammation, missing or misdirected eyelashes, itching along the eyelid base, and the presence of collarettes.
Age Restrictions	Member is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Diagnosed by an ophthalmologist.
Coverage Duration	6 weeks
Other Criteria	Diagnosis must include: eyelash epilation for examination by light microscopy for identification and confirmation of Demodex infestation OR collarettes that are visible on slit lamp examination.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

XELJANZ

Products Affected

- XELJANZ ORAL SOLUTION
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, response to prior therapy. Documentation of negative TB test.
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.
Prescriber Restrictions	Prescribed by a specialist appropriate to the disease state, such as a rheumatologist or gastroenterologist.
Coverage Duration	3 year , Xelzanz XR 22 mg - 16 weeks
Other Criteria	Prior to receiving treatment with Xeljanz for ARTHRITIS-RELATED conditions, the patient must have trial of, or contraindication to, at least one non-biologic DMARD (including but not limited to methotrexate, hydroxychloroquine, sulfasalazine, azathioprine). Dose adjustments and laboratory monitoring parameters should be adhered to as specified in the FDA approved drug labeling. Prior to receiving treatment with Xeljanz XR 22 mg for an ULCERATIVE COLITIS condition, the patient must have an inadequate response or intolerance to tumor necrosis factor blockers (includng but not limited to adalimumab). Dose adjustments and laboratory monitoring parameters should be adhered to as specified in the FDA approved drug labeling.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Part B Prerequisite	No





XERMELO

Products Affected

XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of carcinoid syndrome diarrhea AND diarrhea is inadequately controlled by a stable dose of somatostatin analog (SSA) therapy (e.g., octreotide, Somatuline Depot) for at least 3 months AND used in combination with SSA therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist
Coverage Duration	3 years
Other Criteria	For continuation of therapy-Documentation of a positive clinical response to Xermelo therapy AND Xermelo will continue to be used in combination with SSA therapy.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





XGEVA

Products Affected

• XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and/or failed
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by oncologist
Coverage Duration	5 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





XIFAXAN

Products Affected

• XIFAXAN ORAL TABLET 200 MG, 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	Not covered for PROPHYLAXIS of traveler's diarrhea. For TREATMENT of traveler's diarrhea, not covered for diarrhea due to pathogens other than E. coli, and, not covered for diarrhea complicated by fever or blood in the stool.
Required Medical Information	Small bowel bacterial overgrowth syndrome (SIBO): The patients diagnosis has been confirmed by ONE of the following: A) quantitative culture of upper gut aspirate, B) breath testing (e.g., lactulose hydrogen or glucose hydrogen breath test). OR The patient is experiencing a recurrence of small intestinal bacterial overgrowth (SIBO) after completion of a successful course of the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months for all medically accepted indications, excepting 2 years for hepatic encephalopathy.
Other Criteria	For hepatic encephalopathy: treatment failure, intolerance, or contraindication to lactulose. For travelers diarrhea: treatment failure, intolerance, or contraindication to a fluoroquinolone (such as ciprofloxacin) and azithromycin. For IBS-D: treatment failure, intolerance, or contraindication to at least two of these drugs: loperamide, dicyclomine or diphenoxylate/atropine. For recurrent C. difficile: treatment failure, intolerance, or contraindication to vancomycin. For SIBO symptoms that do not respond to a 7- to 10-day therapeutic trial of ONE of the following, unless the treatment is medically inadvisable or the patient has a history of intolerance to the treatment: metronidazole.





Date Effective: 12/01/2024

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





XOLAIR

Products Affected

• XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	For asthma: weight, IgE level at baseline, baseline FEV1, FEV1 following bronchodilator, medication history, ER visits, and hospitalizations. Evidence of a positive skin test or in-vitro reactivity to a perennial aeroallergen. For uticaria: documentation of persistence of hives associated with itching and prior treatments with outcome.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by a pulmonologist, immunologist, allergist, dermatologist or otolaryngologist.
Coverage Duration	3 year
Other Criteria	Moderate to severe persistent asthma approve if the patient meets criteria 1 and 2: 1) patient has received combination therapy with an inhaled corticosteroid and at least one the following: long-acting beta-agonist (LABA), long-acting muscarinic antagonist (LAMA), leukotriene receptor antagonist, or theophylline, and 2) patient's asthma is uncontrolled or was uncontrolled prior to receiving any Xolair or anti-IL-4/13 therapy (Dupixent) therapy as defined by ONE of the following (a, b, c, d, or e): a) The patient experienced two or more asthma exacerbations requiring treatment with systemic corticosteroids in the previous year OR b) The patient experienced one or more asthma exacerbation requiring hospitalization or an Emergency Department (ED) visit in the previous year OR c) Patient has a forced expiratory volume in 1 second (FEV1) less than 80% predicted OR d) Patient has an FEV1/forced vital capacity (FVC) less than 0.80 OR e) The patient's asthma worsens upon tapering of oral





Date Effective: 12/01/2024

PA Criteria	Criteria Details
	corticosteroid therapy. NOTE: An exception to the requirement for a trial of one additional asthma controller/maintenance medication can be made if the patient has already received anti-IL-4/13 therapy (Dupixent) used concomitantly with an ICS for at least 3 consecutive months. For chronic idiopathic urticaria (CIU), approve if the patient has documented CIU for at least 6 weeks AND failure, intolerance, or contraindication to cyclosporine and montelukast. For continuation of asthma treatment - The patient has responded to therapy as determined by the prescribing physician and continues to receive therapy with one inhaled corticosteroid or inhaled corticosteroid containing combination product. For continuation of CIU treatment - The patient must have responded to therapy as determined by the prescribing physician.
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





XOSPATA

Products Affected

XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapies, and result of prior therapy.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or hematologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





XPOVIO

Products Affected

• XPOVIO

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, other therapies tried and treatment responses.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or hematologist.
Coverage Duration	5 year
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

XTANDI

Products Affected

• XTANDI ORAL CAPSULE

• XTANDI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis, prior therapy used, result of prior therapy. If continuation, prior response to enzalutamide.
Age Restrictions	N/A
Prescriber Restrictions	Prescription must be written by an oncologist or urologist
Coverage Duration	5 years
Other Criteria	Patient must undergo evaluation of seizure risk. For metastatic castration resistant prostate cancer (CRPC) OR metastatic castration-sensitive prostate cancer (CSPC), patient must have a failure, intolerance, or contraindication to abiraterone (Zytiga) prior to initiation of therapy with enzalutamide (Xtandi).
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





XURIDEN

Products Affected

• XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	The starting dose will be approved for 60 mg/kg once daily. Higher doses will be approved (up to 120 mg/kg once daily) in the following situations: Levels of orotic acid in urine remain above normal or increase above the usual or expected range for the patient OR Laboratory values (e.g., red blood cell or white blood cell indices) affected by hereditary orotic aciduria show evidence of worsening OR Worsening of other signs or symptoms of the disease
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Date Effective: 12/01/2024

XYREM

Products Affected

• sodium oxybate

• XYREM

PA Criteria	Criteria Details	
Exclusion Criteria	Not to be used in patients concurrently using alcohol or sedative-hypnotic agents	
Required Medical Information	Diagnosis, other therapies tried and failed	
Age Restrictions	N/A	
Prescriber Restrictions	Prescription must be written by a sleep medicine specialist or neurologist	
Coverage Duration	6 months	
Other Criteria	Dosing approved up to 9 grams per day	
Indications	All Medically-accepted Indications.	
Off-Label Uses	N/A	
Part B Prerequisite	No	





Date Effective: 12/01/2024

ZEJULA

Products Affected

• ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	Diagnosis. Medical records supporting the request.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





ZELBORAF

Products Affected

ZELBORAF

PA Criteria	Criteria Details	
Exclusion Criteria	PATIENTS WITH WILD-TYPE BRAF MELANOMA, PREGNANCY	
Required Medical Information	FOR METASTATIC MELANOMA: DOCUMENTATION OF DIAGONSIS AND BRAF V600E MUTATION AS DETECTED USING AN FDA-APPROVED TEST. FOR ERDHEIM-CHESTER DISEASE: DOCUMENTATION OF DIAGNOSIS AND BRAF V600 MUTATION.	
Age Restrictions	N/A	
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY AN ONCOLOGIST.	
Coverage Duration	5 years	
Other Criteria	PERFORM DERMATOLOGICAL EVALUATIONS PRIOR TO INITIATION OF THERAPY AND EVERY 2 MONTHS WHILE ON THERAPY TO MONITOR FOR NEW PRIMARY MELANOMAS. MONITOR LIVER FUNCTION TESTS PRIOR TO INITIATION OF THERAPY AND AS CLINICALLY INDICATED THEREAFTER. MONITOR ECGs PRIOR TO INITIATION OF THERAPY, AT DAY 15, THEN MONTHLY, AND EVERY 3 MONTHS THEREAFTER.MONITOR PATIENTS FOR OPHTHALMOLOGIC REACTIONS AS CLINICALLY INDICATED.	
Indications	All Medically-accepted Indications.	
Off-Label Uses	N/A	





PA Criteria	Criteria Details
Part B Prerequisite	No





ZILBRYSQ

Products Affected

ZILBRYSQ

PA Criteria	Criteria Details	
Exclusion Criteria	N/A	
Required Medical Information	Diagnosis, Members must have confirmed anti-acetylcholine receptor antibody-positive generalized myasthenia gravis.	
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.	
Prescriber Restrictions	prescribed by a neurologist.	
Coverage Duration	1 year.	
Other Criteria	Patient must have an inadequate response, contraindication, or intolerance to two different immunosuppressant therapies: Azathioprine, mycophenolate, methotrexate, cyclosporine, or tacrolimus.	
Indications	All Medically-accepted Indications.	
Off-Label Uses	N/A	
Part B Prerequisite	No	





ZOLINZA

Products Affected

• ZOLINZA

PA Criteria	Criteria Details	
Exclusion Criteria	N/A	
Required Medical Information	Documentation of diagnosis, previous treatments, and response to treatment.	
Age Restrictions	N/A	
Prescriber Restrictions	Must be prescribed by an oncologist/hematologist.	
Coverage Duration	5 years	
Other Criteria	N/A	
Indications	All Medically-accepted Indications.	
Off-Label Uses	N/A	
Part B Prerequisite	No	





Date Effective: 12/01/2024

ZOLPIDEM

Products Affected

• zolpidem oral tablet

• zolpidem oral tablet,ext release multiphase

PA Criteria	Criteria Details	
Exclusion Criteria	N/A	
Required Medical Information	Diagnosis and prior drug treatments.	
Age Restrictions	N/A	
Prescriber Restrictions	N/A	
Coverage Duration	1 year	
Other Criteria	Patient must have a trial of Trazodone.	
Indications	All Medically-accepted Indications.	
Off-Label Uses	N/A	
Part B Prerequisite	No	





Date Effective: 12/01/2024

ZONISADE

Products Affected

• ZONISADE

PA Criteria	Criteria Details	
Exclusion Criteria	N/A	
Required Medical Information	Diagnosis, previous treatments, and response therapy	
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information	
Prescriber Restrictions	N/A	
Coverage Duration	5 years	
Other Criteria	Must have a intolerance, contraindication, or medical reason the capsule are not acceptable.	
Indications	All Medically-accepted Indications.	
Off-Label Uses	N/A	
Part B Prerequisite	No	





ZTALMY

Products Affected

• ZTALMY

PA Criteria	Criteria Details	
Exclusion Criteria	N/A	
Required Medical Information	Documentation of diagnosis	
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.	
Prescriber Restrictions	Prescribed by Neurologist	
Coverage Duration	5 years	
Other Criteria	N/A	
Indications	All Medically-accepted Indications.	
Off-Label Uses	N/A	
Part B Prerequisite	No	





Date Effective: 12/01/2024

ZURZUVAE

Products Affected

• ZURZUVAE

PA Criteria	Criteria Details	
Exclusion Criteria	N/A	
Required Medical Information	Diagnosis of Postpartum Depression	
Age Restrictions	Beneficiary is of appropriate age for use per FDA prescribing information.	
Prescriber Restrictions	N/A	
Coverage Duration	3 months	
Other Criteria	N/A	
Indications	All Medically-accepted Indications.	
Off-Label Uses	N/A	
Part B Prerequisite	No	





ZYDELIG

Products Affected

• ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	N/A
Required Medical Information	DIAGNOSIS, PRIOR TREATMENTS, RESPONSE TO THERAPY. FOR PATIENTS WITH CLL, DOCUMENTATION OF PRIOR TREATMENT WITH IMBRUVICA. IF NO PREVIOUS FAILURE, RATIONALE AS TO WHY PREFERRED AGENT CANNOT BE USED
Age Restrictions	N/A
Prescriber Restrictions	PRESCRIPTION MUST BE WRITTEN BY ONCOLOGIST OR HEMATOLOGIST
Coverage Duration	5 years
Other Criteria	N/A
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Part B Prerequisite	No





Prior Authorization Criteria

HAP CareSourceTM MI Health Link (Medicare-Medicaid Plan) 2024

Date Effective: 12/01/2024

PART B VERSUS PART D

Products Affected

- ABELCET INTRAVENOUS SUSPENSION 5 MG/ML
- acetylcysteine solution 100 mg/ml (10 %), 200 mg/ml (20 %)
- acyclovir sodium intravenous solution 50 mg/ml
- albuterol sulfate inhalation solution for nebulization 0.63 mg/3 ml, 1.25 mg/3 ml, 2.5 mg/3 ml (0.083 %), 2.5 mg/0.5 ml, 5 mg/ml
- AMBISOME INTRAVENOUS SUSPENSION FOR RECONSTITUTION 50 MG
- amphotericin b injection recon soln 50 mg
- arformoterol inhalation solution for nebulization 15 mcg/2 ml
- AZASAN ORAL TABLET 100 MG, 75 MG
- azathioprine oral tablet 100 mg, 50 mg, 75 mg
- bleomycin injection recon soln 15 unit, 30
 unit
- budesonide inhalation suspension for nebulization 0.25 mg/2 ml, 0.5 mg/2 ml, 1 mg/2 ml
- CLINIMIX 5%/D15W SULFITE FREE INTRAVENOUS PARENTERAL SOLUTION 5 %
- CLINIMIX 4.25%/D10W SULF FREE INTRAVENOUS PARENTERAL SOLUTION 4.25 %
- CLINIMIX 4.25%/D5W SULFIT FREE INTRAVENOUS PARENTERAL SOLUTION 4.25 %

- CLINIMIX 5%-D20W(SULFITE-FREE) INTRAVENOUS PARENTERAL SOLUTION 5 %
- CLINIMIX E 2.75%/D5W SULF FREE INTRAVENOUS PARENTERAL SOLUTION 2.75 %
- CLINIMIX E 4.25%/D10W SUL FREE INTRAVENOUS PARENTERAL SOLUTION 4.25 %
- CLINIMIX E 4.25%/D5W SULF FREE INTRAVENOUS PARENTERAL SOLUTION 4.25 %
- CLINIMIX E 5%/D15W SULFIT FREE INTRAVENOUS PARENTERAL SOLUTION 5 %
- CLINIMIX E 5%/D20W SULFIT FREE INTRAVENOUS PARENTERAL SOLUTION 5 %
- CLINISOL SF 15 % INTRAVENOUS PARENTERAL SOLUTION 15 %
- cromolyn inhalation solution for nebulization 20 mg/2 ml
- cyclophosphamide oral capsule 25 mg, 50 mg
- CYCLOPHOSPHAMIDE ORAL TABLET 25 MG, 50 MG
- cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg
- cyclosporine modified oral solution 100 mg/ml
- cyclosporine oral capsule 100 mg, 25 mg
- ENGERIX-B (PF) INTRAMUSCULAR SUSPENSION 20 MCG/ML
- ENGERIX-B (PF) INTRAMUSCULAR SYRINGE 20 MCG/ML





- ENGERIX-B PEDIATRIC (PF)
 INTRAMUSCULAR SYRINGE 10
 MCG/0.5 ML
- ENVARSUS XR ORAL TABLET EXTENDED RELEASE 24 HR 0.75 MG, 1 MG, 4 MG
- everolimus (immunosuppressive) oral tablet 0.25 mg, 0.5 mg, 0.75 mg, 1 mg
- formoterol fumarate inhalation solution for nebulization 20 mcg/2 ml
- gengraf oral capsule 100 mg, 25 mg
- gengraf oral solution 100 mg/ml
- granisetron hcl oral tablet 1 mg
- HEPLISAV-B (PF) INTRAMUSCULAR SYRINGE 20 MCG/0.5 ML
- IMOVAX RABIES VACCINE (PF) INTRAMUSCULAR RECON SOLN 2.5 UNIT
- intralipid intravenous emulsion 20 %
- INTRALIPID INTRAVENOUS EMULSION 30 %
- ipratropium bromide inhalation solution 0.02 %
- ipratropium-albuterol inhalation solution for nebulization 0.5 mg-3 mg(2.5 mg base)/3 ml
- levalbuterol hcl inhalation solution for nebulization 0.31 mg/3 ml, 0.63 mg/3 ml, 1.25 mg/0.5 ml, 1.25 mg/3 ml
- methylprednisolone oral tablet 16 mg, 32 mg, 4 mg, 8 mg
- mycophenolate mofetil oral capsule 250 mg
- mycophenolate mofetil oral suspension for reconstitution 200 mg/ml
- mycophenolate mofetil oral tablet 500 mg
- mycophenolate sodium oral tablet,delayed release (dr/ec) 180 mg, 360 mg
- ondansetron hcl oral solution 4 mg/5 ml

- ondansetron hcl oral tablet 4 mg, 8 mg
- ondansetron oral tablet, disintegrating 4 mg, 8 mg
- pentamidine inhalation recon soln 300 mg
- PLENAMINE INTRAVENOUS PARENTERAL SOLUTION 15 %
- PREHEVBRIO (PF) INTRAMUSCULAR SUSPENSION 10 MCG/ML
- premasol 10 % intravenous parenteral solution 10 %
- PROGRAF ORAL GRANULES IN PACKET 0.2 MG, 1 MG
- PROSOL 20 % INTRAVENOUS PARENTERAL SOLUTION
- PULMOZYME INHALATION SOLUTION 1 MG/ML
- RABAVERT (PF) INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 2.5 UNIT
- RECOMBIVAX HB (PF)
 INTRAMUSCULAR SUSPENSION 10
 MCG/ML, 40 MCG/ML, 5 MCG/0.5 ML
- RECOMBIVAX HB (PF) INTRAMUSCULAR SYRINGE 10 MCG/ML, 5 MCG/0.5 ML
- sirolimus oral solution 1 mg/ml
- sirolimus oral tablet 0.5 mg, 1 mg, 2 mg
- tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg
- tobramycin in 0.225 % nacl inhalation solution for nebulization 300 mg/5 ml
- tobramycin inhalation solution for nebulization 300 mg/4 ml
- travasol 10 % intravenous parenteral solution 10 %
- TROPHAMINE 10 % INTRAVENOUS PARENTERAL SOLUTION 10 %
- YUPELRI INHALATION SOLUTION FOR NEBULIZATION 175 MCG/3 ML





Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.





Index

A	amphotericin b injection recon soln 50 mg
ABELCET INTRAVENOUS	300
SUSPENSION 5 MG/ML300	aprepitant21
ABILIFY MAINTENA2	ARALAST NP 1
acetylcysteine solution 100 mg/ml (10 %),	ARANESP (IN POLYSORBATE)
200 mg/ml (20 %)300	INJECTION SOLUTION 100 MCG/ML,
ACTIMMUNE3	200 MCG/ML, 25 MCG/ML, 40
acyclovir sodium intravenous solution 50	MCG/ML, 60 MCG/ML 67, 68
mg/ml300	ARANESP (IN POLYSORBATE)
ADALIMUMAB-FKJP SUBCUTANEOUS	INJECTION SYRINGE 67, 68
PEN INJECTOR KIT 87, 88	ARCALYST 22
ADALIMUMAB-FKJP SUBCUTANEOUS	arformoterol inhalation solution for
SYRINGE KIT 87, 88	nebulization 15 mcg/2 ml 300
adapalene topical cream5	armodafinil16
adapalene topical gel 0.3 %5	asenapine maleate
adapalene topical gel with pump5	asperflex (lidocaine) topical cream 131
adapalene topical solution5	AUGTYRO ORAL CAPSULE 40 MG 23
adapalene topical swab5	AURYXIA24
ADBRY SUBCUTANEOUS SYRINGE 6	AUVELITY 17, 18
ADEMPAS 7	AVONEX INTRAMUSCULAR PEN
ADLARITY 8	INJECTOR KIT 146
AIMOVIG AUTOINJECTOR10	AVONEX INTRAMUSCULAR SYRINGE
AJOVY AUTOINJECTOR11	KIT146
AJOVY SYRINGE11	AYVAKIT
AKEEGA 12	AZASAN ORAL TABLET 100 MG, 75
albuterol sulfate inhalation solution for	MG300
nebulization 0.63 mg/3 ml, 1.25 mg/3 ml,	azathioprine oral tablet 100 mg, 50 mg, 75
2.5 mg/3 ml (0.083 %), 2.5 mg/0.5 ml, 5	mg 300
mg/ml300	В
ALECENSA	BALVERSA 26
ALUNBRIG14	BENLYSTA SUBCUTANEOUS 28
alyq177	BESREMI 29
AMBISOME INTRAVENOUS	bexarotene topical228
SUSPENSION FOR	BIVIGAM111, 112
RECONSTITUTION 50 MG 300	bleomycin injection recon soln 15 unit, 30
ambrisentan	unit300
	bosentan





BOSULIF ORAL CAPSULE 100 MG, 50	CLINIMIX E 4.25%/D10W SUL FREE
MG30	INTRAVENOUS PARENTERAL
BOSULIF ORAL TABLET 100 MG, 400	SOLUTION 4.25 % 300
MG, 500 MG 30	CLINIMIX E 4.25%/D5W SULF FREE
BRAFTOVI31	INTRAVENOUS PARENTERAL
brimonidine topical144	SOLUTION 4.25 % 300
BRIVIACT ORAL32	CLINIMIX E 5%/D15W SULFIT FREE
BRONCHITOL33	INTRAVENOUS PARENTERAL
BRUKINSA 34	SOLUTION 5 % 300
budesonide inhalation suspension for	CLINIMIX E 5%/D20W SULFIT FREE
nebulization 0.25 mg/2 ml, 0.5 mg/2 ml, 1	INTRAVENOUS PARENTERAL
mg/2 ml 300	SOLUTION 5 % 300
budesonide oral tablet, delayed and	CLINISOL SF 15 % INTRAVENOUS
ext.release249	PARENTERAL SOLUTION 15 % 300
C	COMETRIQ
CABOMETYX 35	CONTRAVE269, 270
CALQUENCE	COPIKTRA39
CALQUENCE (ACALABRUTINIB MAL)	CORLANOR ORAL SOLUTION 40
36	COSENTYX (2 SYRINGES) 41, 42
CAMZYOS140	COSENTYX PEN 41, 42
CAPLYTA 19, 20	COSENTYX PEN (2 PENS) 41, 42
CAPRELSA ORAL TABLET 100 MG, 300	COSENTYX SUBCUTANEOUS
MG37	SYRINGE 150 MG/ML, 75 MG/0.5 ML
CERDELGA 80	41, 42
CINQAIR	COSENTYX UNOREADY PEN 41, 42
CLINIMIX 5%/D15W SULFITE FREE	COTELLIC
INTRAVENOUS PARENTERAL	cromolyn inhalation solution for
SOLUTION 5 % 300	nebulization 20 mg/2 ml 300
CLINIMIX 4.25%/D10W SULF FREE	cyclophosphamide oral capsule 25 mg, 50
INTRAVENOUS PARENTERAL	mg 300
SOLUTION 4.25 % 300	CYCLOPHOSPHAMIDE ORAL TABLET
CLINIMIX 4.25%/D5W SULFIT FREE	25 MG, 50 MG300
INTRAVENOUS PARENTERAL	cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg
SOLUTION 4.25 % 300	25 mg, 50 mg 300
CLINIMIX 5%-D20W(SULFITE-FREE)	cyclosporine modified oral solution 100
INTRAVENOUS PARENTERAL	mg/ml300
SOLUTION 5 %	cyclosporine oral capsule 100 mg, 25 mg300
CLINIMIX E 2.75%/D5W SULF FREE	CYSTADROPS45
INTRAVENOUS PARENTERAL	D
SOLUTION 2.75 % 300	dasatinib213





DAURISMO ORAL TABLET 100 MG, 25	EPIDIOLEX62		
MG46	EPOGEN INJECTION SOLUTION 10,000		
DAYBUE47	UNIT/ML, 2,000 UNIT/ML, 20,000		
deferiprone	UNIT/2 ML, 20,000 UNIT/ML, 3,000		
DIACOMIT50	UNIT/ML, 4,000 UNIT/ML 67, 68		
diclofenac sodium topical gel 3 %4	EPRONTIA 63		
DIFICID ORAL TABLET51	ERIVEDGE 65		
dihydroergotamine nasal64	ERLEADA ORAL TABLET 240 MG, 60		
DOPTELET (10 TAB PACK) 52	MG6		
DOPTELET (15 TAB PACK) 52	erlotinib227		
DOPTELET (30 TAB PACK) 52	everolimus (antineoplastic)		
DRIZALMA SPRINKLE ORAL	everolimus (immunosuppressive) oral tablet		
CAPSULE, DELAYED REL SPRINKLE	0.25 mg, 0.5 mg, 0.75 mg, 1 mg 301		
20 MG, 30 MG, 40 MG, 60 MG 53	EVRYSDI69		
dronabinol54	F		
DUPIXENT PEN SUBCUTANEOUS PEN	FANAPT19, 20		
INJECTOR 200 MG/1.14 ML, 300 MG/2	FASENRA 70		
ML55, 56	FASENRA PEN70		
DUPIXENT SYRINGE SUBCUTANEOUS	fentanyl citrate buccal lozenge on a handle		
SYRINGE 100 MG/0.67 ML, 200	1,200 mcg, 200 mcg, 600 mcg		
MG/1.14 ML, 300 MG/2 ML 55, 56	FETZIMA ORAL CAPSULE,EXT REL		
DUVYZAT 57	24HR DOSE PACK 20 MG (2)- 40 MG		
E	(26)17, 18		
ENBREL MINI 58, 59	FETZIMA ORAL CAPSULE,EXTENDED		
ENBREL SUBCUTANEOUS SOLUTION	RELEASE 24 HR 17, 18		
	FILSUVEZ71		
ENBREL SUBCUTANEOUS SYRINGE58,	FINTEPLA72		
59	FIRDAPSE15		
ENBREL SURECLICK 58, 59	FLEBOGAMMA DIF INTRAVENOUS		
ENGERIX-B (PF) INTRAMUSCULAR	SOLUTION 10 % 111, 112		
SUSPENSION 20 MCG/ML 300	formoterol fumarate inhalation solution for		
ENGERIX-B (PF) INTRAMUSCULAR	nebulization 20 mcg/2 ml		
SYRINGE 20 MCG/ML 300	FOTIVDA75		
ENGERIX-B PEDIATRIC (PF)	FRUZAQLA76		
INTRAMUSCULAR SYRINGE 10	FYCOMPA		
MCG/0.5 ML301	G		
ENSPRYNG61	gabapentin oral tablet extended release 24 hr		
ENVARSUS XR ORAL TABLET	85		
EXTENDED RELEASE 24 HR 0.75 MG,	GALAFOLD78		
1 MG, 4 MG 301	GAMMAGARD LIQUID 111, 112		





GAMMAGARD S-D (IGA < 1 MCG/ML)	IMBRUVICA ORAL SUSPENSION 100	
	IMBRUVICA ORAL TABLET 140 MG,	
GAMMAKED111, 112	280 MG, 420 MG 100	
GAMMAPLEX111, 112	IMOVAX RABIES VACCINE (PF)	
GAMMAPLEX (WITH SORBITOL) 111,	INTRAMUSCULAR RECON SOLN 2.5	
112	UNIT301	
GAMUNEX-C 111, 112	INBRIJA INHALATION CAPSULE,	
GATTEX 30-VIAL79	W/INHALATION DEVICE101	
GATTEX ONE-VIAL 79	INCRELEX 102	
GAVRETO 81	INLYTA ORAL TABLET 1 MG, 5 MG 104	
gefitinib109	INQOVI 105	
gengraf oral capsule 100 mg, 25 mg 301	INREBIC106	
gengraf oral solution 100 mg/ml301	intralipid intravenous emulsion 20 % 301	
GILOTRIF 82	INTRALIPID INTRAVENOUS	
GLASSIA 1	EMULSION 30 %	
GLEOSTINE 83	INVEGA HAFYERA 108	
glutamine (sickle cell)60	INVEGA SUSTENNA 108	
GRALISE ORAL TABLET EXTENDED	INVEGA TRINZA108	
RELEASE 24 HR 450 MG, 750 MG, 900	ipratropium bromide inhalation solution	
MG85	0.02 % 301	
granisetron hel oral tablet 1 mg 301	ipratropium-albuterol inhalation solution for	
H	nebulization $0.5 \text{ mg}-3 \text{ mg}(2.5 \text{ mg base})/3$	
HADLIMA 87, 88	ml 301	
HADLIMA PUSHTOUCH 87, 88	ivabradine40	
HADLIMA(CF)87, 88	ivermectin oral	
HADLIMA(CF) PUSHTOUCH 87, 88	IWILFIN 113	
HAEGARDA 91, 92	J	
HEMADY89	JAKAFI114	
HEPLISAV-B (PF) INTRAMUSCULAR	JAYPIRCA 115	
SYRINGE 20 MCG/0.5 ML301	JOENJA116	
HUMIRA(CF) (ONLY NDCS STARTING	JUXTAPID94	
WITH 00074) SUBCUTANEOUS	JYNARQUE ORAL TABLETS,	
SYRINGE KIT 10 MG/0.1 ML 95, 96	SEQUENTIAL117	
I	K	
IBRANCE	KALYDECO ORAL GRANULES IN	
icatibant	PACKET 13.4 MG, 25 MG, 5.8 MG, 50	
ICLUSIG	MG, 75 MG 118	
IDHIFA99	KALYDECO ORAL TABLET 118	
IMBRUVICA ORAL CAPSULE 140 MG,	KERENDIA 119, 120	
70 MG 100	KINERET 121	





KISQALI 122	MAYZENT STARTER(FOR 1MG	
KISQALI FEMARA CO-PACK 122	MAINT)146	
KOSELUGO 124	MAYZENT STARTER(FOR 2MG	
KRAZATI125	MAINT)146	
L	MEKINIST ORAL RECON SOLN 141	
lapatinib246	MEKINIST ORAL TABLET 0.5 MG, 2	
LAZCLUZE 127	MG141	
lenalidomide	MEKTOVI142	
LENVIMA 128	methamphetamine 143	
LEUKINE INJECTION RECON SOLN 129	methylprednisolone oral tablet 16 mg, 32	
levalbuterol hel inhalation solution for	mg, 4 mg, 8 mg301	
nebulization 0.31 mg/3 ml, 0.63 mg/3 ml,	metyrosine49	
1.25 mg/0.5 ml, 1.25 mg/3 ml 301	mifepristone oral tablet 300 mg 123	
LIBERVANT253	miglustat80	
LIBTAYO 130	MOUNJARO 84	
lidocaine topical adhesive patch,medicated 5	MULPLETA 145	
%131	MYALEPT148	
LIDOCAINE TOPICAL CREAM 131	mycophenolate mofetil oral capsule 250 mg	
LMX 4 TOPICAL CREAM 131		
LMX 5131	mycophenolate mofetil oral suspension for	
LOKELMA 132	reconstitution 200 mg/ml301	
LONSURF 133	mycophenolate mofetil oral tablet 500 mg	
LORBRENA		
LUMAKRAS ORAL TABLET 120 MG,	mycophenolate sodium oral tablet,delayed	
320 MG 135	release (dr/ec) 180 mg, 360 mg 301	
LYBALVI136, 137	MYFEMBREE149	
LYNPARZA 138	MYTESI44	
LYTGOBI ORAL TABLET 12 MG/DAY	N	
(4 MG X 3), 16 MG/DAY (4 MG X 4),	NAYZILAM 150	
20 MG/DAY (4 MG X 5) 139	NERLYNX 151	
M	NEXLETOL 152, 153	
MAVENCLAD (10 TABLET PACK) 146	NEXLIZET 152, 153	
MAVENCLAD (4 TABLET PACK) 146	NINLARO 154	
MAVENCLAD (5 TABLET PACK) 146	nitisinone	
MAVENCLAD (6 TABLET PACK) 146	NORDITROPIN FLEXPRO 86	
MAVENCLAD (7 TABLET PACK) 146	NOXAFIL ORAL SUSP, DELAYED	
MAVENCLAD (8 TABLET PACK) 146	RELEASE FOR RECON155	
MAVENCLAD (9 TABLET PACK) 146	NUBEQA156	
MAVYRET ORAL TABLET90	NUCALA	
MAYZENT 146	NUEDEXTA158	





NUPLAZID159	OZEMPIC SUBCUTANEOUS PEN	
NURTEC ODT 160	INJECTOR 0.25 MG OR 0.5 MG (2	
NUTROPIN AQ NUSPIN 86	MG/3 ML), 1 MG/DOSE (4 MG/3 ML),	
0	2 MG/DOSE (8 MG/3 ML) 84	
OCALIVA161	P	
OCTAGAM 111, 112	PALYNZIQ175	
ODOMZO162	PANRETIN176	
OGSIVEO ORAL TABLET 100 MG, 150	pazopanib	
MG, 50 MG 163	PEGASYS 107	
OJEMDA ORAL SUSPENSION FOR	PEMAZYRE178	
RECONSTITUTION 164	pentamidine inhalation recon soln 300 mg	
OJEMDA ORAL TABLET 164		
OJJAARA 165	PIQRAY179	
ondansetron hcl oral solution 4 mg/5 ml. 301	PLEGRIDY146	
ondansetron hcl oral tablet 4 mg, 8 mg 301	PLENAMINE INTRAVENOUS	
ondansetron oral tablet, disintegrating 4 mg,	PARENTERAL SOLUTION 15 % 301	
8 mg 301	POMALYST180	
ONUREG 166	posaconazole oral155	
OPSUMIT184, 185	PREHEVBRIO (PF) INTRAMUSCULAR	
ORENCIA CLICKJECT167	SUSPENSION 10 MCG/ML 301	
ORENCIA SUBCUTANEOUS SYRINGE	premasol 10 % intravenous parenteral	
125 MG/ML, 50 MG/0.4 ML, 87.5	solution 10 % 301	
MG/0.7 ML 167	PRIVIGEN111, 112	
ORENITRAM184, 185	PROGRAF ORAL GRANULES IN	
ORFADIN ORAL SUSPENSION 168	PACKET 0.2 MG, 1 MG 301	
ORGOVYX169	PROLASTIN-C INTRAVENOUS	
ORILISSA171	SOLUTION 1	
ORKAMBI ORAL GRANULES IN	PROLIA 181, 182	
PACKET 100-125 MG, 150-188 MG, 75-	PROMACTA 183	
94 MG170	PROSOL 20 % INTRAVENOUS	
ORKAMBI ORAL TABLET 170	PARENTERAL SOLUTION 301	
ORLADEYO91, 92	PULMOZYME INHALATION	
ORSERDU 172	SOLUTION 1 MG/ML 301	
OTEZLA ORAL TABLET 20 MG, 30 MG	PURIXAN186	
	PYRUKYND 187	
OTEZLA STARTER ORAL	Q	
TABLETS, DOSE PACK 10 MG (4)- 20	QINLOCK 188	
MG (51), 10 MG (4)-20 MG (4)-30 MG	QSYMIA269, 270	
(47)173		
$\overrightarrow{OXERVATE}$ 174		





R	\mathbf{S}		
RABAVERT (PF) INTRAMUSCULAR	sajazir91, 92		
SUSPENSION FOR	sapropterin		
RECONSTITUTION 2.5 UNIT 301	SAXENDA 269, 270		
RADICAVA ORS 189	SCEMBLIX ORAL TABLET 100 MG, 20		
RADICAVA ORS STARTER KIT SUSP	MG, 40 MG 206		
189	SECUADO19, 20		
RAVICTI 251	SIGNIFOR		
RECOMBIVAX HB (PF)	sildenafil (pulmonary arterial hypertension)		
INTRAMUSCULAR SUSPENSION 10	oral suspension for reconstitution 10		
MCG/ML, 40 MCG/ML, 5 MCG/0.5 ML	mg/ml 177		
301	sildenafil (pulmonary arterial hypertension)		
RECOMBIVAX HB (PF)	oral tablet 20 mg 177		
INTRAMUSCULAR SYRINGE 10	sirolimus oral solution 1 mg/ml 301		
MCG/ML, 5 MCG/0.5 ML 301	sirolimus oral tablet 0.5 mg, 1 mg, 2 mg 301		
rectasmoothe 131	SIRTURO 208		
RECTICARE 131	SKYCLARYS209		
REGRANEX190	SKYRIZI SUBCUTANEOUS PEN		
REPATHA 191, 192	INJECTOR210, 2		
REPATHA PUSHTRONEX 191, 192	SKYRIZI SUBCUTANEOUS SYRINGE		
REPATHA SURECLICK 191, 192	150 MG/ML210, 211		
RETEVMO ORAL CAPSULE 193	SKYRIZI SUBCUTANEOUS WEARABLE		
RETEVMO ORAL TABLET 120 MG, 160	INJECTOR 180 MG/1.2 ML (150		
MG, 40 MG, 80 MG 193	MG/ML), 360 MG/2.4 ML (150 MG/ML)		
REVCOVI	210, 211		
REXULTI ORAL TABLET 19, 20	sodium oxybate		
REYVOW197	sodium phenylbutyrate oral powder 251		
REZLIDHIA198	SPRITAM 212		
REZUROCK 199	STELARA SUBCUTANEOUS		
RINVOQ200, 201	SOLUTION214, 215		
RINVOQ LQ200, 201	STELARA SUBCUTANEOUS SYRINGE		
ROZLYTREK ORAL CAPSULE 100 MG,	45 MG/0.5 ML, 90 MG/ML 214, 215		
200 MG202	STIVARGA216		
ROZLYTREK ORAL PELLETS IN	sunitinib malate		
PACKET202	SUNOSI217		
RUBRACA	SYMDEKO219		
rufinamide27	SYMLINPEN 120220		
RYBELSUS 84	SYMLINPEN 60		
RYDAPT204	SYMPAZAN ORAL FILM 10 MG, 20 MG,		
	5 MG221		





SYMPROIC 147	tolvaptan oral tablet 30 mg	
T	TRACLEER ORAL TABLET FOR	
TABRECTA 222	SUSPENSION 184, 185	
tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg	travasol 10 % intravenous parenteral	
	solution 10 % 301	
tadalafil (pulm. hypertension) 177	TRELSTAR INTRAMUSCULAR	
TAFINLAR ORAL CAPSULE 224	SUSPENSION FOR	
TAFINLAR ORAL TABLET FOR	RECONSTITUTION240	
SUSPENSION 224	tretinoin topical	
TAGRISSO	trientine oral capsule 250 mg 242	
TAKHZYRO91, 92	TRINTELLIX 17, 18	
TALZENNA	TROPHAMINE 10 % INTRAVENOUS	
TASIGNA ORAL CAPSULE 150 MG, 200	PARENTERAL SOLUTION 10 % 301	
MG, 50 MG229	TRULICITY 84	
tasimelteon93	TRUQAP243	
TAVALISSE230	TUKYSA244	
TAVNEOS231	TURALIO ORAL CAPSULE 125 MG 245	
tazarotene topical cream232	TYMLOS247	
tazarotene topical gel232	TYVASO 184, 185	
TAZVERIK233	TYVASO INSTITUTIONAL START KIT	
TEGSEDI234		
TEPMETKO	TYVASO REFILL KIT 184, 185	
teriparatide subcutaneous pen injector 20	TYVASO STARTER KIT 184, 185	
mcg/dose (600mcg/2.4ml) 73, 74	U	
TERIPARATIDE SUBCUTANEOUS PEN	ubrelvy	
INJECTOR 20 MCG/DOSE	ULTRA LIDO131	
(620MCG/2.48ML)73, 74	UPTRAVI ORAL250	
testosterone transdermal gel 236, 237	\mathbf{V}	
testosterone transdermal gel in metered-dose	VALCHLOR	
pump 12.5 mg/ 1.25 gram (1 %), 20.25	VALTOCO	
mg/1.25 gram (1.62 %)236, 237	VANFLYTA	
testosterone transdermal gel in packet 236,	VELTASSA255	
237	VENCLEXTA	
TIBSOVO238	VENCLEXTA STARTING PACK 256	
TOBI PODHALER103	VENTAVIS 184, 185	
tobramycin in 0.225 % nacl inhalation	VEOZAH257	
solution for nebulization 300 mg/5 ml 301	VERSACLOZ 19, 20	
tobramycin inhalation solution for	VERZENIO	
nebulization 300 mg/4 ml 301	vilazodone	
TOLVAPTAN ORAL TABLET 15 MG 205	VITRAKVI	





Date Effective: 12/01/2024

VIZIMPRO 260	XGEVA	280
VONJO261	XIFAXAN ORAL TABLET 200 MG, 550	
VORANIGO 262	MG281, 282	
voriconazole intravenous	XOLAIR	283, 284
VOXZOGO265	XOSPATA	•
VRAYLAR ORAL CAPSULE 19, 20	XPOVIO	286
VUITY	XTANDI ORAL CAPSULE	287
VYNDAMAX	XTANDI ORAL TABLET	287
VYNDAQEL	XURIDEN	288
W	XYREM	289
WAINUA267	Y	
WAKIX268	YUPELRI INHALATION SOLUTION	
WEGOVY269, 270	FOR NEBULIZATION 175 MCG/3 ML	
WELIREG		301
WINREVAIR272, 273	Z	
X	ZEJULA ORAL TABLET	290
XALKORI ORAL CAPSULE13	ZELBORAF	291, 292
XALKORI ORAL PELLET 150 MG, 20	ZEMAIRA	1
MG, 50 MG13	ZILBRYSQ	293
XATMEP274	ZOLINZA	294
XCOPRI275	zolpidem oral tablet	295
XCOPRI MAINTENANCE PACK 275	zolpidem oral tablet,ext release multiphase	
XCOPRI TITRATION PACK275	-	295
XDEMVY276	ZONISADE	296
XELJANZ ORAL SOLUTION 277, 278	ZTALMY	297
XELJANZ ORAL TABLET 277, 278	ZURZUVAE	298
XELJANZ XR277, 278	ZYDELIG	299
XENICAL269, 270	ZYKADIA	13
XERMELO 279		

HAP CareSource™ MI Health Link (Medicare-Medicaid Plan) is a health plan that contracts with both Medicare and Michigan Medicaid to provide benefits of both programs to enrollees.