

ADULTHEDIS® CODING GUIDE 2022-2023



This guide provides HEDIS coding information only, not necessarily payment guidance. Refer to your state's guidance for payment details and telehealth regulations.

MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS
Prevention and S	creening		
Breast Cancer Screening (BCS)* Females 50-74 years	Women 50-74 years of age who had a mammogram to screen for breast cancer once every 27 months.	Biopsies, breast ultrasounds or MRIs do not count towards this measure.	CPT®: 77061-3, 77065-7 Potential exclusion for Bilateral Mastectomy in patient history ICD-10: Z90.13
Care of Older Adults (COA) 66 years and older	Adults 66 years and older who had each of the following during the measurement year: • Medication Review • Functional Status Assessment • Pain Assessment Services rendered during a telephone visit, e-visit or virtual check-in meet criteria for the Advance Care Planning, Functional Assessments and Pain Assessment indicators.	Medication Review: A complete medication list, signed and dated during the measurement year by the appropriate practitioner type; member not required to be present. Functional Status Assessment: Documentation must include evidence of a complete functional status assessment and the date it was performed. Must include one of the following: • ADLs • IADLs • Standardized Functional Assessment Tool Pain Assessment: Evidence of assessment and date performed. Must include one of the following: • Documentation that patient was assessed for pain • Use of standardized assessment tool and result	Medication Review CPT: 90863, 99483, 99605-6 CPT II: 1160F - AND - Medication List HCPCS: G8427 CPT II: 1159F - OR - Transitional Care Management CPT: 99495-6 Functional Status Assessment CPT: 99483 HCPCS: G0438-9 CPT II: 1170F Pain Assessment CPT II: 1125F, 1126F Note: CPT II codes are for quality reporting purposes only, not for payment.



MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS		
Prevention and Screening					
Cervical Cancer Screening (CCS)* Females 21-64 years	Women 21-64 years of age who were screened for cervical cancer using one of the following	Cervical cytology during the measurement year or the two years prior.	High Risk HPV CPT: 87624-5 HCPCS: G0476		
Temales 21 04 years	methods: • Women 21-64 years of age who had cervical cytology performed within the last three years	Documentation must include both : • A note indicating the date when the cervical cytology was performed • The result or findings	Cervical Cytology CPT: 88141-3, 88147-8, 88150, 88152-3, 88164-7, 88174-5 HCPCS: G0123-4, G0141, G0143-5, G0147-8, P3000-1, Q0091		
	 Women 30-64 years of age who had cervical high-risk human papillomavirus (hrHPV) testing performed during the measurement year or the four years prior, and who were 30 years or older as of the date of testing Women 30-64 years of age who had cervical cytology/hrHPV cotesting during the measurement year or the four years prior 	 Documentation must include both: A note indicating the date when the cervical cytology and/or the HPV test were performed; the cervical cytology and HPV test must be from the same data source The results or findings 	Potential exclusion from measure for Hysterectomy in patient history ICD-10: Q51.5, Z90.710, Z90.712 CPT: 51925, 56308, 57530-1, 57540, 57545, 57550, 57555-6, 58150, 58152, 58200, 58210, 58240, 58260, 58262-3, 58267, 58270, 58275, 58280, 58285, 58290-4, 58548, 58550, 58552-4, 58570-3, 58575, 58951, 58953-4, 58956, 59135		
Chlamydia Screening in Women (CHL) Females 16-24 years	Women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.	Women are considered sexually active if there is evidence of the following: • Contraceptives are prescribed • Via medical coding	CPT: 87110, 87270, 87320, 87490-2, 87810		
Colorectal Cancer Screening (COL)* 45-75 years	Adults 45-75 years of age who had appropriate screening for colorectal cancer. One or more screenings for colorectal cancer. One or more screenings for colorectal cancer. Any of the following meet criteria: • Fecal occult blood test Yearly • FIT sDNA test Every 3 Years • CT Colonography Every 5 Years • Flexible sigmoidoscopy Every 5 years • Colonoscopy Every 10 Years	Documentation in the medical record must include a note indicating the date the colorectal cancer screening was performed. A result is not required if the documentation is clearly part of the "medical history" section of the record; if this is not clear, the result or finding must also be present (this ensures that the screening was performed and not merely ordered).	FOBT CPT: 82270, 82274 HCPCS: G0328 FIT sDNA CPT: 81528 CT Colonography CPT: 74261-3 Flexible Sigmoidoscopy CPT: 45330-5, 45337-8, 45340-2, 45346-7, 45349-50 HCPCS: G0104 Colonoscopy CPT: 44388-94, 44397, 44401-8, 45355, 45378-93, 45398 HCPCS: G0105, G0121		

MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS
Prevention and S Colorectal Cancer Screening (COL)* 45-75 years	creening		Potential exclusion from measure Colorectal Cancer ICD-10: Z85.038, Z85.048, C18.0-9, C19, C20, C21.2, C21.8, C78.5 Total Colectomy CPT: 44150-3, 44152-3, 44155-8, 44210-12
Respiratory Cond	litions		
Asthma Medication Ratio (AMR) 5-64 years	The percentage of those 5-64 years with persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.	 Medications given as oral, inhaler, or as an injection are counted. Controller medication(s) should account for ≥ 0.50 of total asthma medications dispensed. 	Compliance occurs only if patient fills prescription. Encourage patient to fill prescriptions on time and take medications as prescribed.
Appropriate Testing for Pharyngitis (CWP) 3 years and older	Members 3 years and older with a diagnosis of pharyngitis, dispensed an antibiotic, and received a group A streptococcus (strep) test for the episode.	Documentation in the medical record must include all of the following: • Diagnosis of pharyngitis • Antibiotic dispensed on or up to three days after date of service • And received group A strep test	Need evidence of all three components: Strep Test CPT: 87070-1, 87081, 87430, 87650-2, 87880 - WITH - Pharyngitis Diagnostic ICD-10: J02.0, J02.8-9, J03.00-1, J03.80-1, J03.90-1 - AND - Prescribed antibiotic is filled by a pharmacy
Cardiovascular C	onditions		
Controlling High Blood Pressure (CBP)* 18-85 years	Adults 18-85 years with a diagnosis of essential hypertension and whose BP was adequately controlled during the measurement year. Telephone visits, e-visits and virtual check-ins are appropriate settings for BP readings. BPs can be taken by any digital device.	Criteria for control BP < 140/90 on or after the date of the second diagnosis of hypertension. Exclusions: Patients with evident ESRD Diagnosis of pregnancy during the current year Patients who had an admission to a non-acute inpatient setting in the current year	Record Review Notation of the most recent BP in the medical record. Blood Pressure CPT II: 3074F, 3075F, 3077F, 3078F, 3079F, 3080F - OR - Taken during Outpatient, without Revenue Code: CPT: 99201-5, 99211-5, 99241-5, 99341-5, 99347-50, 99381-7, 99391-7, 99401-4, 99411-2, 99429, 99455-6, 99483 HCPCS: G0402, G0438, G0439, G0463, T1015 - OR - Telephone Visit CPT: 98966-8, 99441-3 - OR - Online Assessment CPT: 98969-72, 99421-3, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061, G2062, G2063 Note: CPT II codes are for quality reporting purposes only, not for payment.

MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS
Cardiovascular C	onditions		
Statin Therapy for Patients With Cardiovascular Disease (SPC)* Males 21-75 years Females 40-75 years	Adults who were identified as having clinical ASCVD and met the following criteria: • Received statin therapy • Were adherent to therapy at least 80% of treatment period	Patients should be dispensed at least one high or moderate-intensity statin and stay on medication for at least 80% of treatment period. Include patients with a discharge diagnosis of MI. Patients with a diagnosis of CABG, PCI or any other revascularization	Compliance occurs only if patient fills prescription. Encourage patient to fill prescriptions on time and take medications as prescribed. Telehealth can be used to prescribe to eligible patients, if appropriate for the patient. Exclusions: Frailty and advanced illness (must meet both), palliative care, ESRD,
		process are automatically included in measure.	cirrhosis, pregnancy or IVF (current or prior year), and muscular pain or disease*
Diabetes Care			
Statin Therapy for Patients With Diabetes (SPD)* 40-75 years	Adults who were identified as having diabetes and DO NOT HAVE clinical ASCVD and met the following criteria:	Patients who were identified as having diabetes with diagnosis of MI, CABG, PCI or any other revascularization process are automatically excluded in measure.	Compliance occurs only if patient fills prescription. Encourage patient to fill prescriptions on time and take medications as prescribed.
.0 10 900.0	 Received statin therapy Were adherent to therapy at least 80% of treatment period 	Patients should be dispensed at least one high or moderate-intensity statin and stay on medication for at least 80% of treatment period.	Telehealth can be used to prescribe to eligible patients, if appropriate for the patient.
Hemoglobin A1c Control for Patients With Diabetes (HBD)*	levels during the measurement year:	Notation of the most recent HbA1c screening noting date performed and result performed in current year.	HbA1c CPT: 83036-7 CPT II: 3044F, 3046F, 3051F, 3052F
18-75 years with type 1 or type 2 diabetes		Ranges and thresholds do not meet criteria for this indicator. A distinct numeric result is required for numerator compliance when documented in the medical record.	Note: CPT II codes are for quality reporting purposes only, not for payment.
Eye Exam for Patients With Diabetes (EED)* 18-75 years with type 1 or type 2 diabetes	Adults who had a screening or monitoring for diabetic retinal disease in the measurement year.	A retinal or dilated eye exam by an optometrist or ophthalmologist in current year, a negative retinal or dilated exam (negative for retinopathy) done by an optometrist or ophthalmologist in previous year.	Eye Exam by Eye Care Professional CPT: 67028, 67030-1, 67036, 67039-43, 67101, 67105, 67107-8, 67110, 67113, 67121, 67141, 67145, 67208, 67210, 67218, 67220-1, 67227-8, 92002, 92004, 92012, 92014, 92018-9, 92134, 92201-2, 92225-8, 92230, 92235, 92240, 92250, 92260, 99203-5, 99213-5, 99242-5 HCPCS: S0620, S0621, S3000
			Eye Exam by any Professional CPT: 92229 (automated eye exam) CPT II: 2022F, 2023F, 2024F, 2025F, 2026F, 2033F, 3072F
			Note: CPT II codes are for quality reporting

purposes only, not for payment.

MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS
Diabetes Care			
Blood Pressure Control for Patients With Diabetes (BPD)*	Adults with diabetes whose blood pressure was adequately controlled (< 140/90 mm Hg) during	Note the most recent BP in the medical record.	Blood Pressure CPT II: 3074F, 3075F, 3077F, 3078F, 3079F, 3080F
18-75 years with type 1 or type 2 diabetes	the measurement year.		Taken During Outpatient CPT: 99201-5, 99211-5, 99241-5, 99341-5, 99347-50, 99381-7, 99391-7, 99401-4, 99411-2, 99429, 99455-6, 99483 HCPCS: G0402, G0438, G0439, G0463, T1015 Blood Pressure CPT II: 3074F, 3075F, 3077F, 3078F, 3079F, 3080F Online Assessment CPT: 98969-72, 99421-3, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061 Note: CPT II codes are for quality reporting purposes only, not for payment.
Kidney Health Evaluation for Patients With Diabetes (KED)* 18-85 years with type 1 or type 2 diabetes	Percentage of adults with diabetes (type 1 and type 2) who received a kidney health evaluation during the measurement year.	Defined by an estimated glomerular filtration rate (eGFR) AND a urine albumin-creatinine ratio (uACR); both quantitative urine albumin test and urine creatinine test with service dates four or less days apart. Exclusion: ESRD or dialysis at any time during patients history.	eGFR CPT: 80047-8, 80050, 80053, 80069, 82565 With Urine Albumin Creatinine Ratio Lab Test (uACR) - OR - Quantitative Urine Albumin CPT: 82043 With Urine Creatinine CPT: 82570
Behavioral Health	Conditions		
Antidepressant Medication Management (AMM) 18 years and over	The percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment. • Acute Phase: The percentage of members who remained on an antidepressant medication for at least 84 days (12 weeks).	Telehealth, virtual check-in or e-visit can be used to prescribe to eligible patients, if appropriate for the patient. Exclusion: Patients in hospice at any time during the measurement year.	Compliance occurs only if patient fills prescription. Encourage patient to fill prescriptions on time and take medications as prescribed.



MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS
Behavioral Healt	h Conditions		
Antidepressant Medication Management (AMM) 18 years and over	Continuation Phase: The percentage of members who remained on an antidepressant medication for at least 180 days (6 months).		
Medication Mana	agement and Care Co	ordination	
Transitions of Care (TRC) 18 years and over	The percentage of adult discharges who had each of the following:	Notification of inpatient admission requires documentation in medical record of receipt of notification of	Any of following meet patient engagement: Outpatient Visit

18 years and over of the following: record of receipt of notification of **Outpatient Visit CPT:** 99201-5, 99211-5, 99241-5, Medicare only Notification of inpatient inpatient admission on the day of admission admission through two days after 99341-5, 99347-50, 99381-7, Receipt of discharge the admission (three total days). 99391-7, 99401-4, 99411-2, 99429, information 99455-6, 99483 Patient engagement Receipt of discharge information **HCPCS:** G0402, G0438-9, G0463, T1015 after inpatient discharge documented in medical record on Medication the day of discharge through two **Telephone Visit** reconciliation postdays after the discharge (three total **CPT:** 98966-8, 99441-3 discharge days). **Transitional Care Management (TCM)** Patient engagement provided within **Services CPT:** 99495-6 30 days after discharge **Online Assessment CPT:** 98969-72, Medication reconciliation by a 99421-3, 99444, 99457 prescribing practitioner, clinical **HCPCS:** G0071, G2010, G2012, G2061, pharmacist, physician assistant G2062, G2063 or registered nurse on the date of discharge through 30 days after **Medication Reconciliation CPT:** 99483. discharge. 99495-6 **CPT II:** 1111F Note: CPT II codes are for quality reporting purposes only, not for payment.

DESCRIPTION COMPLIANCE CODES & MEASURE DOCUMENTATION TIPS OF MEASURE **MEASURE TIPS** Medication Management and Care Coordination Follow-Up After The percentage of adult Follow-up within seven days after **Outpatient Visit** emergency department the ED visit (eight total days). Visits **CPT:** 99201-05, 99211-5, 99241-5. **Emergency Department Visit for** visits who have multiple that occur on the date of the ED visit 99341-5, 99347-50, 99381-7, 99391-7, **People with Multiple** high-risk chronic do count. 99401-4, 99411-2, 99429, 99455-6, **High-Risk Chronic** conditions who had a 99483. 99492-4 **Conditions** follow-up service within This measure addresses the need **HCPCS:** G0402, G0438, G0439, G0463, seven days of the ED visit. for medication management and (FMC) T1015 18 years and over care coordination for this vulnerable **UBREV:** 0510-23, 0526-9, 0982-3 Medicare only population. -OR-

medication management and te coordination for this vulnerable bulation.

To Coordination for this vulnerable bulation.

To Coordination for this vulnerable bulation.

To Coordination for this vulnerable bulation.

99401-4, 99411-2, 99429, 99455-6, 99483, 99492-4 HCPCS: G0402, G0438, G0439, G0463, T1015 UBREV: 0510-23, 0526-9, 0982-3 - OR Telephone Visit CPT: 98966-8, 99441-3 - OR TMC Services CPT: 99495-6 - OR Case Management CPT: 99366 HCPCS: T1016-7, T2022-3 - OR Complex Care Management CPT: 99439, 99487, 99489-91 HCPCS: G0506 - OR Outpatient/Telehealth Behavioral Visit,

Setting Unspecified CPT: 90791-2, 90832-4, 90836-40, 90845, 90847, 90849, 90853, 90875-6, 99221-3, 99231-3, 99238-9, 99251-5 **With POS:** 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71, 72

– OR –

BH Outpatient

CPT: 98960-2, 99078, 99201-5, 99211-5, 99241-5, 99341-5, 99347-50, 99381-7, 99391-7, 99401-4, 99411-2, 99483, 99510

HCPCS: G0155, G0176-7, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036-7, H0039, H0040, H2000, H2010-1, H2013-20, T1015 **Revenue:** 0510, 0513, 0515-7,

0519-23, 0526-9, 0900, 0902-4, 0911, 0914-7, 0919, 0982-3

- OR -

CMHC Outpatient Visit Setting, Unspecified CPT With CMHC POS: 53

- OR -

Telehealth Visit Outpatient Setting, Unspecified CPT With Telehealth POS:02, 10

MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS
Medication Mana	gement and Care Cod	ordination	
Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions (FMC) 18 years and over Medicare only			− <i>OR</i> − Observation Visit CPT: 99217-20 − <i>OR</i> − Online Assessment CPT: 98969-72, 99421-3, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061-3 − <i>OR</i> − SUD Services CPT: 99408-9 HCPCS: G0396-7, G0443, H0001, H0005, H0007, H0015-6, H0022, H0047, H0050, H2035-6, T1006, T1012 UBREV: 0906, 0944-5
Access/Availabili	ty of Care		
Adults' Access to Preventive/ Ambulatory Health Services (AAP) 20 years and over	Adults who had an ambulatory or preventive care visit.	This measure looks at whether adult patients receive preventive and ambulatory services. To qualify, the patient must receive an evaluation and management care during an ambulatory visit with a medical professional. Care received in an emergency department or inpatient setting does not qualify. Telehealth is an option for this measure.	CPT: 99201-5, 99211-5, 99241-5, 99341-2, 99341-2, 99343-5, 99347-50, 99381-7, 99391-7, 99401-4, 99411-2, 99429, 99483, 92002, 92004, 92012, 92014, 99304-10, 99315-6, 99318, 99324-8, 99334-7 HCPCS: G0402, G0438-9, G0463, T1015, S0620-1 ICD10: Z00.00-01, Z00.121, Z00.129, Z00.3, Z00.5, Z00.8, Z02.0-6, Z02.71, Z02.79, Z02.81-3, Z02.89, Z02.9, Z76.1, Z76.2 Revenue Code: 0510-7, 0519-23, 0526-9, 0982-3, 0524-5 Telephone Visit CPT: 98966-8, 99441-3 Online Assessment CPT: 98969-72, 99421-44, 99457 HCPCS: G0071, G2010, G2012, G2061-3
Prenatal and Postpartum Care (PPC) All Ages	The measure assesses the following facets of prenatal and postpartum care: • Timeliness of Prenatal Care The percentage of deliveries that received a prenatal care visit in the first trimester, on or before the enrollment start date or within 42 days of enrollment in the organization.	A qualified prenatal care visit with an OB/GYN or other prenatal care practitioner or PCP. Documentation must include the date the visit occurred and include at least one of the following: • Auscultation for fetal heart tones • Pelvic exam with OB observations (a pap test alone does not count) • Measurement of fundal height	Stand-Alone Prenatal Visit CPT: 99500 CPT II: 0500F, 0501F, 0502F HCPCS: H1000-4 - OR - Prenatal Bundled Services CPT: 59400, 59425-6, 59510, 59610, 59618 HCPCS: H1005

MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS
Access/Availability	y of Care		
Prenatal and Postpartum Care (PPC) All Ages	• Postpartum Care The percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery. Services provided via telephone, e-visit or virtual check-in are eligible for both measures.	 Basic OB visit that includes one of the following prenatal procedures: Complete OB lab panel TORCH antibody panel Rubella antibody with Rh incompatibility blood typing Ultrasound of pregnant uterus Documentation indicating pregnancy which includes: Standardized prenatal flow sheet LMP or EDD or gestational age Prenatal risk assessment and counseling/education A complete obstetrical history Gravidity and parity Positive pregnancy test result Visits with a PCP or other family practitioner must follow the same guidelines but also include a documented diagnosis of pregnancy. A qualified postpartum visit must include a note indicating the date the visit occurred and include at least one of the following: Notation of postpartum care Pelvic exam Evaluation of weight, blood pressure, breasts and abdomen (must have all four components) Perineal or cesarean incision/wound check Screening for depression, anxiety, tobacco use, substance use disorder or preexisting mental health disorders Glucose screening for women with gestational diabetes Documentation of infant care or breastfeeding, resumption of intercourse, birth spacing or family planning, sleep/fatigue, resumption of physical activity, attainment of healthy weight 	- OR - Any of the following WITH an appropriate pregnancy diagnosis Prenatal Visit CPT: 99201-5, 99211-5, 99241-5, 99483 HCPCS: G0463, T1015 Telephone Visit CPT: 98966-8, 99441-3 Online Assessment CPT: 98969-72, 99421-4, 99457 HCPCS: G0071, G2010, G2012, G2061-3 Postpartum Visit CPT: 57170, 58300, 59430, 99501 HCPCS: G0101 CPT II: 0503F ICD-10: Z01.411, Z01.419, Z01.42, Z30.430, Z39.1, Z39.2 - OR - Postpartum Bundled CPT: 59400, 59410, 59510, 59515, 59610, 59614, 59618, 59622 Cervical Cytology CPT: 88141-3, 88147-8, 88150, 88152-4, 88164-7, 88174-5 HCPCS: G0123-4, G0141, G0143-5, G0147-8, P3000, P3001, Q0091 Note: CPT II codes are for quality reporting purposes only, not for payment.

MEASURE	DESCRIPTION OF MEASURE	DOCUMENTATION TIPS	COMPLIANCE CODES & MEASURE TIPS
Overuse/Appropri	iateness		
Use of Opioids at High Dosage (HDO)* 18 years and over	The proportion of members 18 years and older receiving prescription opioids at a high dosage (average morphine milligram equivalent dose [MME] ≥ 90mg) for ≥ 15 days during the measurement year.	Reduce the number of adults prescribed high dose opioids for ≥ 15 days. A lower rate indicates better performance. Increasing total MME dose of opioids is related to increased risk of overdose and adverse events. Necessity of use of high doses should be clear. Patients with cancer, sickle cell disease or members receiving palliative care are excluded from this measure.	Patients are considered out of compliance if their prescription average MME was ≥ 90mg during the treatment period. This measure does not include the following opioid medications: Injectables Opioid cough and cold products Ionsys® (fentanyl transdermal patch) Methadone for the treatment of opioid use disorder
Use of Opioids from Multiple Providers (UOP) 18 years and over	The proportion of members 18 years and older, receiving prescription opioids for ≥ 15 days during the measurement year who received opioids from multiple providers. Three rates are reported: Multiple Prescribers Patients receiving prescriptions for opioids from four or more different prescribers during the calendar year. Multiple Pharmacies Patients receiving prescriptions for opioids from four or more different pharmacies during the current calendar year. Multiple Prescribers and Multiple Pharmacies Patients receiving prescriptions for opioids from four or more different pharmacies Patients receiving prescriptions for opioids from four or more different prescribers and four or more different prescribers and four or more different pharmacies during the calendar year.	Reduce the number of adults prescribed opioids for ≥ 15 days by multiple providers. A lower rate indicates better performance for all three rates. Patient use of increasing number of prescribers or pharmacies may signal risk for uncoordinated care. Clinical correlation is encouraged so that providers can evaluate for risk of diversion, misuse or a substance use disorder. Providers are encouraged to communicate with each other for ideal management of member.	Patients are considered out of compliance if they received prescription opioids from four or more different prescribers. Multiple Pharmacies Patients are considered out of compliance if they received prescription opioids from four or more different pharmacies. Multiple Prescribers and Multiple Pharmacies Patients are considered out of compliance if they received prescription opioids from four or more different prescribers and four or more different prescribers and four or more different pharmacies. The following opioid medications are excluded from this measure: Injectables Opioid cough and cold products Single-agent and combination buprenorphine products used as part of medication-assisted treatment of opioid use disorder Ionsys® (fentanyl transdermal patch) Methadone for the treatment of opioid use disorder

calendar year.



*Palliative Care is a required exclusion for this measure
HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).

Please Note: The codes in this document are derived from the NCQA HEDIS Volume 2 Technical Specifications for Health Plans. These codes are examples of codes typically billed for this type of service and are subject to change. Billing these codes does not guarantee payment. CPT II codes are for quality reporting purposes only. Submitting claims using these codes helps improve reporting of quality measure performance.