

**INDIANA HEALTH COVERAGE PROGRAMS (IHCP)
PCSK9 INHIBITORS AND SELECT LIPO TROPICS PRIOR AUTHORIZATION (PA) REQUEST FORM**



CareSource Pharmacy Prior Authorization Form
P.O. Box 8738
Dayton, OH 45401-8738
Fax: (866) 930-0019

Today's Date

/ /

Non-Urgent ☐

Urgent ☐

Note: This form must be completed by the prescribing provider.

All sections must be completed or the request will be returned*

Patient's CareSource # <input type="text"/>	Date of Birth <input type="text"/> / <input type="text"/> / <input type="text"/>
Patient's Name	Prescriber's Name
Prescriber's Indiana License # <input type="text"/>	Specialty
Prescriber's NPI # <input type="text"/>	Prescriber's Signature
Prescriber Fax <input type="text"/> - <input type="text"/> - <input type="text"/>	Prescriber Phone <input type="text"/> - <input type="text"/> - <input type="text"/>
Prescriber Address:	Date(s) of Service: Start Date:
Diagnosis:	Diagnosis Code:

I attest that the information on this form is accurate:

Physician Signature: _____ **Date:** _____

Requested Medication	Strength	Quantity	Dosage Regimen

PA Requirements for Evkeeza (evinacumab-dgnb):

- Member has a diagnosis of homozygous familial hypercholesterolemia (HoFH)? ☐ Yes ☐ No
- Medication prescribed by, or in consultation with, a cardiologist or endocrinologist? ☐ Yes ☐ No
- Select one of the following:
 - ☐ Member is 5 years of age or older and less than 7 years of age.
 - ☐ Member is 7 years of age or older and less than 10 years of age and one of the following:

PA Requirements for Evkeeza (evinacumab-dgnb) continued:

- i. Member has trial and failure history of at least 90 days of therapy with rosuvastatin 20 mg?
☐ Yes ☐ No
 - ii. Provider has submitted documentation of intolerance/contraindication to rosuvastatin?
☐ Yes ☐ No
 - ☐ Member is 10 years of age or older and less than 18 years of age and one of the following:
 - i. Member has trial and failure history with Repatha (evolocumab)? ☐ Yes ☐ No
Drug/dose/date(s): _____
 - ii. Member has trial and failure history of at least 90 days of high dose rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy concurrently with ezetimibe (or documented intolerance/contraindication to statins/ezetimibe) AND provider has submitted medical justification for use of Evkeeza (evinacumab-dgnb) over Repatha (evolocumab)? ☐ Yes ☐ No
Drug/dose/date(s): _____
 - ☐ Member is 18 years of age or older and one of the following:
 - i. Member has trial and failure history with Praluent (alirocumab) OR Repatha (evolocumab)?
☐ Yes ☐ No
Drug/dose/date(s): _____
 - ii. Member has trial and failure history of at least 90 days of high dose rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy concurrently with ezetimibe (or documented intolerance/contraindication to statins/ezetimibe) AND provider has submitted medical justification for use of Evkeeza (evinacumab-dgnb) over Praluent (alirocumab) and Repatha (evolocumab)? ☐ Yes ☐ No
Drug/dose/date(s): _____
4. Select one of the following:
- ☐ Member will utilize maximally tolerated statin therapy with or without ezetimibe concurrently with Evkeeza (for those 7 years of age and older).
 - ☐ Provider has submitted documented intolerance to statin and/or ezetimibe therapy or medical rationale against use of statin or ezetimibe therapy.
5. Requested dose is 15 mg/kg every 4 weeks or less? ☐ Yes ☐ No Member weight: _____ LB / KG (circle one)

PA Requirements for Juxtapid (lomitapide mesylate):

- 1. Member is enrolled in the Juxtapid/lomitapide REMS program and prescriber is monitoring in accordance with REMS requirements? ☐ Yes ☐ No
- 2. Member is 18 years of age or older? ☐ Yes ☐ No
- 3. Medication prescribed by, or in consultation with, a cardiologist or endocrinologist? ☐ Yes ☐ No
- 4. Select one of the following:
 - ☐ Member has trial and failure history of Praluent (alirocumab) or Repatha (evolocumab).
Drug/dose/date(s): _____

PA Requirements for Juxtapid (lomitapide mesylate) continued:

- ☐ Member has trial and failure history of at least 90 days of high dose rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy concurrently with ezetimibe (or documented intolerance/contraindication to statins/ezetimibe) AND provider has submitted medical justification for use of Juxtapid (lomitapide mesylate) over Praluent (alirocumab) and Repatha (evolocumab).

Drug/dose/date(s): _____

5. For those of childbearing potential, documentation of a negative pregnancy test obtained in the past 30 days is attached and prescriber has counseled member on risks associated with conceiving while utilizing Juxtapid and appropriate methods of contraception? ☐ Yes ☐ No

Prescriber Name and Signature: _____

6. Select one of the following:

- ☐ Member will utilize maximally tolerated statin therapy with or without ezetimibe concurrently with Juxtapid.
- ☐ Provider has submitted documented intolerance to statin and/or ezetimibe therapy or medical rationale against use of statin or ezetimibe therapy.

7. Requested dose is 60 mg/day or less? ☐ Yes ☐ No

PA Requirements for Leqvio (inclisiran):

1. Select one of the following:

- ☐ Member has a diagnosis of primary hyperlipidemia with clinical atherosclerotic cardiovascular disease (ASCVD) or is at increased risk for ASCVD with a baseline LDL-C level of ≥ 55 mg/dL (documentation required).
- ☐ Member has diagnosis of heterozygous familial hypercholesterolemia (HeFH) with a baseline LDL-C level of ≥ 70 mg/dL (documentation required).

2. Member is 18 years of age or older? ☐ Yes ☐ No

3. Prescribed by, or in consultation with, a cardiologist or endocrinologist? ☐ Yes ☐ No

4. Select one of the following:

- ☐ Member has trial and failure history of Praluent (alirocumab) or Repatha (evolocumab).

Drug/dose/date(s): _____

- ☐ Member has trial and failure history of at least 90 days of high dose rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy concurrently with ezetimibe (or documented intolerance/contraindication to statins/ezetimibe) AND provider has submitted medical justification for use of Leqvio (inclisiran) over Praluent (alirocumab) and Repatha (evolocumab).

Drug/dose/date(s): _____

5. Select one of the following:

- ☐ Member will utilize maximally tolerated statin therapy with or without ezetimibe concurrently with Leqvio.
- ☐ Provider has submitted documented intolerance to statin and/or ezetimibe therapy or medical rationale against use of statin or ezetimibe therapy.

6. Select one of the following:

- ☐ Member is initiating therapy and requested dose does not exceed 284 mg every three months.
- ☐ Member is established on therapy and requested dose does not exceed 284 mg every six months.

PA Requirements for Niacin ER

1. Diagnosis of severe hypertriglyceridemia (baseline triglycerides ≥ 500 mg/dL)? ☐ Yes ☐ No

If **Yes**, then select one of the following:

- ☐ Member is on concurrent therapy with all of the following for at least 90 days: omega-3 fatty acid (omega-3-acid ethyl esters or icosapent ethyl), fibric acid derivative and statin therapy.
Drug/dose/date(s): _____
- ☐ Member has a documented intolerance of omega-3 fatty acid, fibric acid derivative **AND** statin therapy **OR** medical justification for use of Niacin ER over omega-3 fatty acid, fibric acid derivative **AND** statin therapy.
Please explain: _____

2. Member is 17 years of age or older? ☐ Yes ☐ No

PA Requirements for Praluent (alirocumab):

1. Select one of the following:

- ☐ Member has a diagnosis of clinical ASCVD, is at Very High Risk requiring therapy for secondary prevention, AND has persistently elevated LDL-C (≥ 55 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy or has documented intolerance of both rosuvastatin and atorvastatin OR medical rationale against the use of statin therapy.*
- ☐ Member has a diagnosis of clinical ASCVD, is NOT at Very High Risk requiring therapy for secondary prevention, AND has persistently elevated LDL-C (≥ 70 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy WITH ezetimibe or has documented intolerance of rosuvastatin and atorvastatin and/or ezetimibe OR medical rationale against the use of statin therapy and/or ezetimibe therapy.
- ☐ Member has a diagnosis of clinical ASCVD, with a baseline LDL-C ≥ 190 mg/dL, not due to secondary causes, without clinical or genetic diagnosis of familial hypercholesterolemia, requiring therapy for secondary prevention AND has persistently elevated LDL-C (≥ 70 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy or has documented intolerance of both rosuvastatin and atorvastatin OR medical rationale against the use of statin therapy.*
- ☐ Member has a diagnosis of clinical ASCVD, is at Very High Risk with a baseline LDL-C ≥ 190 mg/dL not due to secondary causes, a diagnosis of familial hypercholesterolemia, requiring therapy for secondary prevention, AND has persistently elevated LDL-C (≥ 55 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy or has documented intolerance of both rosuvastatin and atorvastatin OR medical rationale against the use of statin therapy.*

PA Requirements for Praluent (alirocumab) continued:

☐ Member has a diagnosis of primary hyperlipidemia, without clinical ASCVD, with a baseline LDL-C ≥ 190 mg/dL not due to secondary causes, with or without concomitant ASCVD risk factors, requiring therapy for primary prevention AND persistently elevated LDL-C (≥ 100 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy or has documented intolerance of both rosuvastatin and atorvastatin OR medical rationale against the use of statin therapy.*

☐ Member has a diagnosis of homozygous familial hypercholesterolemia (HoFH) or heterozygous familial hypercholesterolemia (HeFH) AND persistently elevated LDL-C (≥ 70 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy WITH ezetimibe or has documented intolerance of rosuvastatin and atorvastatin and/or ezetimibe OR medical rationale against the use of statin therapy and/or ezetimibe therapy.

*** For members requiring >25% additional lowering of LDL-C ONLY ($\leq 25\%$ LDL-C lowering must utilize high intensity statin therapy WITH ezetimibe as first line).**

For above diagnoses that require medical justification for use of Praluent over statin and/or ezetimibe therapy, please provide justification here:

2. Member is 18 years of age or older? ☐ Yes ☐ No

3. Select one of the following:

- ☐ Member will utilize maximally tolerated statin therapy with or without ezetimibe concurrently with Praluent.
- ☐ Provider has submitted documented intolerance to statin and/or ezetimibe therapy or medical rationale against use of statin or ezetimibe therapy.

4. Select one of the following:

- ☐ Requested dose is 75 mg every two weeks.
- ☐ Requested dose is 300 mg every four weeks.
- ☐ Requested dose is 150 mg every two weeks **AND** the member has one of the following:
 - ☐ Diagnosis of homozygous familial hypercholesterolemia.
 - ☐ Diagnosis of heterozygous familial hypercholesterolemia and member is undergoing LDL apheresis.

Member has not achieved clinically meaningful response after at least 4 weeks of dosing at 75 mg every 2 weeks or 300 mg every 4 weeks.

PA Requirements for Repatha (evolocumab):

1. Select one of the following:

- ☐ Member has a diagnosis of clinical ASCVD, is at Very High Risk requiring therapy for secondary prevention, AND has persistently elevated LDL-C (≥ 55 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy or has documented intolerance of both rosuvastatin and atorvastatin OR medical rationale against the use of statin therapy.*
- ☐ Member has a diagnosis of clinical ASCVD, is NOT at Very High Risk requiring therapy for secondary prevention, AND has persistently elevated LDL-C (≥ 70 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy WITH ezetimibe or has documented intolerance of rosuvastatin and atorvastatin and/or ezetimibe OR medical rationale against the use of statin therapy and/or ezetimibe.
- ☐ Member has a diagnosis of clinical ASCVD, with a baseline LDL-C ≥ 190 mg/dL, not due to secondary causes, without clinical or genetic diagnosis of familial hypercholesterolemia, requiring therapy for secondary prevention AND has persistently elevated LDL-C (≥ 70 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy or has documented intolerance of both rosuvastatin and atorvastatin OR medical rationale against the use of statin therapy.*
- ☐ Member has a diagnosis of clinical ASCVD, is at Very High Risk with a baseline LDL-C ≥ 190 mg/dL not due to secondary causes, a diagnosis of familial hypercholesterolemia, requiring therapy for secondary prevention, AND has persistently elevated LDL-C (≥ 55 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy or has documented intolerance of both rosuvastatin and atorvastatin OR medical rationale against the use of statin therapy.*
- ☐ Member has a diagnosis of primary hyperlipidemia, without clinical ASCVD, with a baseline LDL-C ≥ 190 mg/dL not due to secondary causes, with or without concomitant ASCVD risk factors, requiring therapy for primary prevention AND persistently elevated LDL-C (≥ 100 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy or has documented intolerance of both rosuvastatin and atorvastatin OR medical rationale against the use of statin therapy.*
- ☐ Member has a diagnosis of homozygous familial hypercholesterolemia (HoFH) or heterozygous familial hypercholesterolemia (HeFH) AND persistently elevated LDL-C (≥ 70 mg/dL) despite treatment with 90 days of therapy with high intensity rosuvastatin (20 mg/40 mg) or atorvastatin (40 mg/80 mg, if rosuvastatin intolerant) therapy WITH ezetimibe or has documented intolerance of rosuvastatin and atorvastatin and/or ezetimibe OR medical rationale against the use of statin therapy and/or ezetimibe.

2. **For members requiring >25% additional lowering of LDL-C ONLY ($\leq 25\%$ LDL-C lowering must utilize high intensity statin therapy WITH ezetimibe as first line).**

Note: Documentation of any and all intolerances to statins and/or ezetimibe must be provided.

For any of the above diagnoses that have medical rationale against the use of statin and/or ezetimibe therapy please provide here:

PA Requirements for Repatha (evolocumab) continued:

2. Select one of the following:

- ☐ Member is 18 years of age or older.
- ☐ Member is 10 years of age or older and has a diagnosis of either HoFH or HeFH.

3. Select one of the following:

- ☐ Member will utilize maximally tolerated statin therapy with or without ezetimibe concurrently with Repatha.
- ☐ Provider has submitted documented intolerance to statin and/or ezetimibe therapy or medical rationale against use of statin or ezetimibe therapy.

4. Select one of the following:

- ☐ Requested dose is 140 mg every two weeks.
- ☐ Requested dose is 420 mg once monthly.
- ☐ Requested dose is 420 mg every two weeks **AND** the member has one of the following:
 - ☐ Diagnosis of HoFH and has not achieved clinically meaningful response after at least 12 weeks at 420 mg once monthly dosing.
 - ☐ Member is receiving lipid apheresis.

CONFIDENTIAL INFORMATION

This facsimile and any attached document are confidential and are intended for the use of individual or entity to which it is addressed. If you have received this in error, please notify us by telephone immediately at **1-844-607-2831**.