

PHARMACY POLICY STATEMENT

Marketplace

DRUG NAME	Visudyne (verteporfin)
BENEFIT TYPE	Medical
STATUS	Prior Authorization Required

Visudyne is a light activated drug used in photodynamic therapy (PDT) to treat certain cases of choroidal neovascularization (CNV). A course of therapy is a 2-step process. First, Visudyne is administered. Second, Visudyne is activated with light from a nonthermal diode laser. Photoactivation of Visudyne is controlled by the light dose delivered. Patients must avoid exposure of skin or eyes to direct sunlight or bright indoor light for 5 days following the procedure.

CNV is the creation of new blood vessels in the choroid layer of the eye and can lead to vision loss. Age-related macular degeneration (AMD) is the most common cause of CNV.

Visudyne (verteporfin) will be considered for coverage when the following criteria are met:

Choroidal Neovascularization (CNV)

For **initial** authorization:

1. Member is at least 18 years of age; AND
2. Medication must be prescribed by or in consultation with an ophthalmologist; AND
3. Member has a documented diagnosis of predominantly classic subfoveal choroidal neovascularization (CNV) due to one of the following:
 - a) "Wet" age-related macular degeneration (AMD)
 - b) Pathologic myopia
 - c) Presumed ocular histoplasmosis; AND
4. Member has tried and failed bevacizumab; AND
5. Member does NOT have predominantly occult subfoveal CNV.
6. **Dosage allowed/Quantity limit:** 6 mg/m² body surface area IV

If all the above requirements are met, the medication will be approved for 3 months (1 dose per eye).

For **reauthorization**:

1. Chart notes must document positive clinical response (e.g., slowed progression of vision loss) following photodynamic treatment; AND
2. Choroidal neovascular leakage has recurred as detected on fluorescein angiography (FA) or optical coherence tomography (OCT).

If all the above requirements are met, the medication will be approved for an additional 3 months (1 dose per eye).

CareSource considers Visudyne (verteporfin) not medically necessary for the treatment of conditions that are not listed in this document. For any other indication, please refer to the Off-Label policy.

DATE	ACTION/DESCRIPTION
10/19/2021	New policy created for Visudyne.
10/04/2023	Revised/added references.

References:

1. Visudyne [prescribing information]. Bausch & Lomb; 2023.
2. Soubrane G, Bressler NM. Treatment of subfoveal choroidal neovascularisation in age related macular degeneration: focus on clinical application of verteporfin photodynamic therapy. *Br J Ophthalmol*. 2001;85(4):483-495. doi:10.1136/bjo.85.4.483
3. Wormald R, Evans J, Smeeth L, Henshaw K. Photodynamic therapy for neovascular age-related macular degeneration. *Cochrane Database Syst Rev*. 2007;(3):CD002030. Published 2007 Jul 18. doi:10.1002/14651858.CD002030.pub3
4. Flaxel CJ, Adelman RA, Bailey ST, et al. Age-Related Macular Degeneration Preferred Practice Pattern® [published correction appears in *Ophthalmology*. 2020 Sep;127(9):1279]. *Ophthalmology*. 2020;127(1):P1-P65. doi:10.1016/j.optha.2019.09.024
5. Zhu Y, Zhang T, Xu G, Peng L. Anti-vascular endothelial growth factor for choroidal neovascularisation in people with pathological myopia. *Cochrane Database Syst Rev*. 2016;12(12):CD011160. Published 2016 Dec 15. doi:10.1002/14651858.CD011160.pub2
6. Chen Y, Han X, Gordon I, et al. A systematic review of clinical practice guidelines for myopic macular degeneration. *J Glob Health*. 2022;12:04026. Published 2022 Mar 26. doi:10.7189/jogh.12.04026
7. Thuruthumaly C, Yee DC, Rao PK. Presumed ocular histoplasmosis. *Curr Opin Ophthalmol*. 2014;25(6):508-512. doi:10.1097/ICU.0000000000000100

Effective date: 04/01/2024

Revised date: 10/04/2023