

PHARMACY POLICY STATEMENT

North Carolina Marketplace

DRUG NAME	Enzyme Replacement Therapy (ERT) for Fabry Disease: Fabrazyme (agalsidase beta) and Elfabrio (pegunigalsidase alfa-iwxj)
BENEFIT TYPE	Medical
STATUS	Prior Authorization Required

Fabrazyme, approved by the FDA in 2003, is an enzyme replacement therapy (ERT) indicated for the treatment of confirmed Fabry disease, to replace the enzyme alpha-galactosidase A (alpha-Gal A). Fabry disease, a lysosomal storage disorder, is a rare genetic disease caused by certain mutations of the *GLA* gene resulting in deficient alpha-Gal A. Normally this enzyme breaks down certain lipids in lysosomes, such as globotriaosylceramide (GL-3). Without it, GL-3 accumulates in blood vessels, the kidneys, heart, nerves, and other organs. The continuous build-up of GL-3 results in progressive cell damage and subsequent symptoms and manifestations in the affected organ systems. Elfabrio is a “biobetter” of Fabrazyme and was designed to have an increased half-life and reduced immunogenicity.

ERT for Fabry Disease will be considered for coverage when the following criteria are met:

Fabry Disease

For **initial** authorization:

1. For Fabrazyme: Member is at least 2 years of age OR for Elfabrio: Member is at least 18 years of age; AND
2. Medication must be prescribed by or in consultation with a medical geneticist, nephrologist, cardiologist, neurologist, or metabolic specialist; AND
3. Member has a diagnosis of Fabry disease confirmed by genetic testing which identifies a pathogenic mutation of the *GLA* gene; AND
4. Member displays symptoms of Fabry disease (i.e., neuropathic pain, renal disease, cardiac disease, abdominal pain, impaired sweating)
NOTE: Exception-- Males with "classic" gene variants do not need to be symptomatic to qualify for treatment. Males with "non-classic" gene variants and asymptomatic females may be treated if there is evidence of injury to the heart, kidney, or central nervous system (CNS); AND
5. ERT will NOT be used in combination with Galafold.
6. **Dosage allowed/Quantity limit:** 1 mg/kg every 2 weeks as an IV infusion.

If all the above requirements are met, the medication will be approved for 6 months.

For **reauthorization**:

1. Chart notes must show positive clinical response such as stabilized kidney function (e.g., GFR, proteinuria), reduced plasma or tissue GL-3 levels, or other improved Fabry symptoms (such as neuropathic pain).

If all the above requirements are met, the medication will be approved for an additional 12 months.

CareSource considers ERT for Fabry Disease not medically necessary for the treatment of conditions that are not listed in this document. For any other indication, please refer to the Off-Label policy.

DATE	ACTION/DESCRIPTION
06/17/2021	New policy for Fabrazyme created.
11/22/2022	Annual review; added reference.
06/28/2023	Changed name of policy and added Elfabrio. Clarified note in #4.
04/30/2025	Updated references.

References:

1. Fabrazyme (agalsidase beta) [package insert]. Cambridge, MA; Genzyme Corporation; 2024.
2. Banikazemi M, Bultas J, Waldek S, et al. Agalsidase-beta therapy for advanced Fabry disease: a randomized trial. *Ann Intern Med*. 2007;146(2):77-86. doi:10.7326/0003-4819-146-2-200701160-00148
3. Eng CM, Guffon N, Wilcox WR, et al. Safety and efficacy of recombinant human alpha-galactosidase A replacement therapy in Fabry's disease. *N Engl J Med*. 2001;345(1):9-16. doi:10.1056/NEJM200107053450102
4. Wallace EL, Goker-Alpan O, Wilcox WR, et al. Head-to-head trial of pegunigalsidase alfa versus agalsidase beta in patients with Fabry disease and deteriorating renal function: results from the 2-year randomised phase III BALANCE study. *J Med Genet*. 2024;61(6):520-530. Published 2024 May 21. doi:10.1136/jmg-2023-109445
5. Laney DA, Bennett RL, Clarke V, et al. Fabry disease practice guidelines: recommendations of the National Society of Genetic Counselors. *J Genet Couns*. 2013;22(5):555-564. doi:10.1007/s10897-013-9613-3
6. Hopkin RJ, Jefferies JL, Laney DA, et al. The management and treatment of children with Fabry disease: A United States-based perspective. *Mol Genet Metab*. 2016;117(2):104-113. doi:10.1016/j.ymgme.2015.10.007
7. Ortiz A, Germain DP, Desnick RJ, et al. Fabry disease revisited: Management and treatment recommendations for adult patients. *Mol Genet Metab*. 2018;123(4):416-427. doi:10.1016/j.ymgme.2018.02.014
8. Germain DP, Fouilhoux A, Decramer S, et al. Consensus recommendations for diagnosis, management and treatment of Fabry disease in paediatric patients. *Clin Genet*. 2019;96(2):107-117. doi:10.1111/cge.13546
9. Elfabrio. [prescribing information]. Chiesi USA, Inc.; 2024.
10. Schiffmann R, Goker-Alpan O, Holida M, et al. Pegunigalsidase alfa, a novel PEGylated enzyme replacement therapy for Fabry disease, provides sustained plasma concentrations and favorable pharmacodynamics: A 1-year Phase 1/2 clinical trial. *J Inherit Metab Dis*. 2019;42(3):534-544. doi:10.1002/jimd.12080
11. Germain DP, Altarescu G, Barriaes-Villa R, et al. An expert consensus on practical clinical recommendations and guidance for patients with classic Fabry disease. *Mol Genet Metab*. 2022;137(1-2):49-61. doi:10.1016/j.ymgme.2022.07.010
12. Biegstraaten M, Arngrímsson R, Barbey F, et al. Recommendations for initiation and cessation of enzyme replacement therapy in patients with Fabry disease: the European Fabry Working Group consensus document. *Orphanet J Rare Dis*. 2015;10:36. Published 2015 Mar 27. doi:10.1186/s13023-015-0253-6

Effective date: 10/01/2025

Revised date: 04/30/2025