

MEDICAL POLICY STATEMENT Wisconsin Marketplace

wisconsin warketpiace				
Policy Name & Number	Date Effective			
Mechanical Stretching Devices-WI MP-MM-1685	06/01/2025-09/30/2025			
Policy Type				
MEDICAL				

Medical Policy Statements are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased, or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage or Certificate of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other plan policies and procedures.

Medical Policy Statements do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage or Certificate of Coverage) for the service(s) referenced in the Medical Policy Statement. Except as otherwise required by law, if there is a conflict between the Medical Policy Statement and the plan contract, then the plan contract will be the controlling document used to make the determination.

According to the rules of Mental Health Parity Addiction Equity Act (MHPAEA), coverage for the diagnosis and treatment of a behavioral health disorder will not be subject to any limitations that are less favorable than the limitations that apply to medical conditions as covered under this policy.

Table of Contents

Α.	Subject	2
	Background	
	Definitions	
	Policy	
	Conditions of Coverage	
	Related Policies/Rules	
	Review/Revision History	
ш	Peferences	1



A. Subject

Mechanical Stretching Devices

B. Background

Mechanical stretching devices are intended to restore range of motion (ROM) for joint stiffness or contracture by stretching joints. These devices provide passive stretching to an adjustable degree for a selected duration for multiple sessions. A variety of mechanical stretching devices are available for extension or flexion of the shoulder, elbow, wrist, fingers, knee, ankle, and toes. These devices can provide stretching for longer periods than a physical therapist and are generally used as adjunct treatment to physical therapy and/or exercise.

Mechanical Stretching Devices (also known as dynamic splinting systems) include:

- low-load prolonged duration stretch devices (LLPS)
- static progressive stretch (SPS) splint devices
- patient actuated serial stretch (PASS) devices

C. Definitions

- Low-load Prolonged Duration Stretch Devices (LLPS) These devices permit
 resisted active and passive motion (elastic traction) within a limited range. LLPS
 devices maintain a set level of tension by means of incorporated rubber bands or
 springs.
- Patient Actuated Serial Stretch (PASS) Devices These devices hold the joint in a set position but allow for manual modification of the joint angle and may allow for active motion without resistance (inelastic traction). This type of device itself does not exert a stress on the tissue unless the joint angle is set at the maximum ROM.
- Static Progressive Stretch Devices (SPS) These devices hold the joint in a set position but allow for manual modification of the joint angle and may allow for active motion without resistance (inelastic traction).

D. Policy

- I. Common Ground Healthcare Cooperative (CGHC) considers dynamic splinting devices medically necessary durable medical equipment (DME) as an adjunct treatment to physical therapy, massage and/or exercise for an existing joint contracture when the following clinical criteria is met:
 - A. medically necessary only for the following joints: knee, elbow, wrist, finger, ankle and toe
 - B. after three weeks of exercise and skilled therapy in the initial subacute injury or post-operative period in members with:
 - 1. signs and symptoms of persistent joint stiffness or contracture
 - 2. limited range of motion that poses a meaningful functional limitation as judged by a physician
 - C. may be used for an initial period of 4 weeks, a subsequent 4-week period with reevaluation, and then up to 4 months based on continued improvement



- II. In the acute post-operative period for members who have undergone additional surgery to improve the range of motion of a previously affected joint, CGHC considers use of an LLPS device medically necessary for:
 - A. an initial four-week period
 - B. an additional four-week period, if improvement was noted after the initial four weeks, for up to 4 months

III. Non-Covered Services

- A. CGHC considers the use of dynamic splinting experimental and investigational for the following indications, including but not limited to:
 - 1. adhesive capsulitis
 - 2. carpal tunnel syndrome
 - 3. cerebral palsy
 - 4. foot drop associated with neuromuscular diseases,
 - 5. hallux valgus
 - 6. head and spinal cord injuries
 - 7. Improvement of outcomes following botulinum toxin injection for treatment of limb spasticity
 - 8. injuries of the ankle and shoulder
 - 9. multiple sclerosis
 - 10. muscular dystrophy
 - 11. plantar fasciitis
 - 12. rheumatoid arthritis
 - 13. stroke
 - 14. trismus
- B. CGHC considers the following devices experimental and investigational due to insufficient scientific evidence of efficacy:
 - 1. patient actuated serial stretch (PASS) devices (for example, ERMI Knee Extensionater® and ERMI Shoulder Extensionater®)
 - 2. static progressive stretch devices (SPS) (for example, Joint Active Systems (JAS) splints (for example, JAS Elbow, JAS Shoulder, JAS Ankle, JAS Knee, JAS Wrist, and JAS Pronation-Supination)

E. Conditions of Coverage

All claims for LLPS are subject to post-payment review by CGHC.

F. Related Policies/Rules

NA



G. Review/Revision History

	DATE	ACTION
Date Issued	08/14/2024	New Policy. Approved at Committee.
Date Revised	03/12/2025	Updated references. Approved at Committee
Date Effective	06/01/2025	
Date Archived		This Policy is no longer active and has been archived. Please note that there could be other Policies that may have some of the same rules incorporated and CareSource reserves the right to follow CMS/State/NCCI guidelines without a formal documented Policy.

H. References

- 1. Dynamic Joint Extension and Flexion Devices (ACG: A-0882 (AC)). MCG. 28th ed. Accessed January 29, 2025. www.careweb.careguidelines.com
- Furia JP, Willis FB, Shanmugam R, Curran SA. Systematic review of contracture reduction in the lower extremity with dynamic splinting. *Adv Ther*. 2013;30(8):763-770. doi:10.1007/s12325-013-0052-1
- 3. Glasgow C, Tooth LR, Fleming J, et al. Dynamic splinting for the stiff hand after trauma: predictors of contracture resolution predictors of contracture resolution. *J Hand Ther*. 2011;24(3):195-205. doi:10.1016/j.jht.2011.03.001
- Harvey LA, Katalinic OM, Herbert RD, et al. Stretch for the treatment and prevention of contractures. *Cochrane Database Syst Rev.* 2017;1:CD007455. doi:10.1002/14651858.CD007455.pub3
- Medical Technology Directory. Mechanical Stretching Devices for the Treatment of Joint Contractures of the Extremities. Hayes; 2022. Accessed January 29, 2025. www.evidence.hayesinc.com
- Jongs RA, Harvey LA, Gwinn T, Lucas BR. Dynamic splints do not reduce contracture following distal radial fracture: a randomised controlled trial. J Physiotherapy. 2012;58(3):173-180. doi:10.1016/S1836-9553(12)70108-X
- Zatarain LA, Smith DK, Deng J, et al. A randomized feasibility trial to evaluate use of the jaw dynasplint to prevent trismus in patients with head and neck cancer receiving primary or adjuvant radiation-based therapy. *Integr Cancer Ther*. 2018;17(3):960-967. doi:10.1177/1534735418784363

Independent medical review -12/21