

MEDICAL POLICY STATEMENT WEST VIRGINIA MARKETPLACE PLANS

Policy Name		Policy Number	Date Effective	
Breast Reduction Surgery		MM-0248	08/01/2020	
Policy Type				
MEDICAL	Administrative	Pharmacy	Reimbursement	

Medical Policy Statement prepared by CSMG Co. and its affiliates (including CareSource) are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other policies and procedures.

Medical Policy Statements prepared by CSMG Co. and its affiliates (including CareSource) do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage) for the service(s) referenced in the Medical Policy Statement. If there is a conflict between the Medical Policy Statement and the plan contract (i.e., Evidence of Coverage), then the plan contract (i.e., Evidence of Coverage) will be the controlling document used to make the determination.

Table of Contents

A.	Subject	2
	Background	
	Definitions	
	Policy	
	Conditions of Coverage	
	Related Polices/Rules	
	Review/Revision History	
	Peferances	



MM-0248 Effective Date: 08/01/2020



Breast Reduction Surgery

B. Background

Women who suffer from macromastia (or excessively large breasts) seeking breast reduction typically present with complaints of a feeling of heaviness, chronic pain and tension in the neck, shoulders and upper back. Macromastia commonly causes permanent grooving and ulceration of the shoulder following years of wearing support bras to try to minimize symptoms. As much as two to five pounds of excess breast tissue is routinely removed during a reduction mammoplasty.

Reduction mammoplasty is a surgical procedure reducing the weight and volume of the breast. Indications for surgery include: chronic pain and skin conditions, neuropathy, breast discomfort, physical impairment and psychological symptoms that can be associated with poor self-esteem and loss of desire to engage in activities.

C. Definitions

- Macromastia (Breast Hypertrophy) An increase in the volume and weight of breast tissue relative to the general body habitus.
- Functional/Physical or Physiological Impairment Physical/functional or
 physiological impairment causes deviation from the normal function of a tissue or
 organ. This results in a significantly limited, impaired or delayed capacity to move
 and coordinate actions, or perform physical activities and is exhibited by difficulties in
 one or more of the following areas: physical and motor tasks; independent
 movement; performing basic life functions.
- Symptomatic Breast Hypertrophy A syndrome of persistent neck and shoulder pain, shoulder grooving from brassiere straps, chronic intertriginous rash of the inframammary fold and/or frequent episodes of headache, backache, and upper extremity neuropathies caused by an increase in the volume and weight of breast tissue beyond normal proportions.
- Cellulitis An acute spreading bacterial infection in the deeper layers of skin associated with an abrasion or cut and characterized by redness, warmth and swelling.
- Intertriginous Rash Dermatitis occurring between juxtaposed folds of skin, caused by retention of moisture and warmth and providing an environment favoring overgrowth of normal skin micro-organisms.
- **Kyphosis** Over-curvature of the thoracic vertebrae (upper back) associated with: degenerative diseases such as arthritis, developmental problems, or with osteoporotic compression fractures of vertebral bodies.
- The Schnur Sliding Scale Has been promoted for use in calculating the amount of breast tissue to be removed in reduction mammoplasty (Appendix A).
- Cosmetic Procedures Procedures that correct an anatomical congenital anomaly without improving or restoring physiologic function are considered cosmetic procedures.



MM-0248

Effective Date: 08/01/2020



- Breast reduction surgery for non-cosmetic indications requires a prior authorization and is considered medically necessary when ALL of the following clinical criteria are met:
 - A. Member must be eighteen (18) years or older, or growth is complete.
 - B. Breast size interferes with activities of daily living, as indicated by 1 or more of the following:
 - 1. Arm numbness consistent with brachial plexus compression syndrome
 - 2. Cervical pain;
 - 3. Chronic breast pain;
 - 4. Headaches:
 - 5. Nipple position greater than 21 cm below suprasternal notch;
 - 6. Persistent redness and erythema (intertrigo) below breasts;
 - 7. Restriction of physical activity;
 - 8. Severe bra strap grooving or ulceration of shoulder;
 - 9. Shoulder pain;
 - 10. Thoracic kyphosis; or
 - 11. Upper or lower back pain
 - C. Failure to relieve symptoms with nonsurgical treatment that includes 1 or more of the following:
 - 1. Medically supervised weight loss program for overweight or obese patient;
 - 2. Topical and oral antifungal agents for intertrigo;
 - 3. Trial of nonsteroidal anti-inflammatory drugs to treat pain in neck, shoulder, upper or lower back, or breast; or
 - 4. Wound care for skin ulceration.
 - D. Preoperative evaluation by surgeon concludes that amount of breast tissue to be removed (by mass or volume) will provide a reasonable expectation of symptomatic relief.
 - E. No evidence of breast cancer
 - 1. As evidenced by results of a physical exam completed by a physician within the last year if under 40.
 - 2. Women 40 to 54 years of age or older must have documentation of a mammogram negative for cancer performed within the year prior to the date of the planned breast reduction surgery.
 - 3. Women 55 and older may switch to mammograms every 2 years.
- II. Breast reduction surgery following mastectomy to achieve symmetry is covered as part of the *Women's Health and Cancer Rights Act (WHCRA)*. Please refer to the CareSource Medical policy titled *Breast Reconstruction Post Mastectomy* for additional information.
- III. Breast Asymmetry: For medical necessity and criteria for surgery to correct breast asymmetry see CareSource Medical Policy *Breast Reconstructive Post Mastectomy*.



MM-0248

Effective Date: 08/01/2020


IV. Schnur Sliding Scale

- A. The Schnur Sliding Scale is an evaluation tool used to determine the appropriate volume of tissue to be removed relative to a patient's total body surface area (BSA).
 - 1. This estimation can be instrumental in determining whether breast reduction surgery is being planned for cosmetic reasons or as a medically necessary procedure. In a survey of plastic surgeons utilizing this scale, Schnur et al.(1991) determined that a member whose removed breast weight was above the 22nd percentile were likely to receive the procedure for medical reasons.
 - 2. The weight of tissue to be removed from each breast must be above the 22nd percentile on the Schnur Sliding Scale (Appendix A below) based on the individual's body surface area (BSA).
 - 3. The body surface area in meters squared (m²) is calculated using the Mosteller formula as follows:
 - a. Square root of height (inches) x weight (lbs) divided by 3131.

Appendix A: Schnur Sliding Scale

Body Surface Area and Minimum Requirement for Breast Tissue Removal		
Body Surface Area m ²	Grams per Breast of Minimum Breast Tissue to be Removed	
1.350-1.374	199	
1.375-1.399	208	
1.400-1.424	218	
1.425-1.449	227	
1.450-1.474	238	
1.475-1.499	249	
1.500-1.524	260	
1.525-1.549	272	
1.550-1.574	284	
1.575-1.599	297	
1.600-1.624	310	
1.625-1.649	324	
1.650-1.674	338	
1.675-1.699	354	
1.700-1.724	370	
1.725-1.749	386	
1.750-1.774	404	
1.775-1.799	422	





	Effective Bate: 00/01/2020
1.800-1.824	441
1.825-1.849	461
1.850-1.874	482
1.875-1.899	504
1.900-1.924	527
1.925-1.949	550
1.950-1.974	575
1.975-1.999	601
2.000-2.024	628
2.025-2.049	657
2.050-2.074	687
2.075-2.099	717
2.100-2.124	750
2.125-2.149	784
2.150-2.174	819
2.175-2.199	856
2.200-2.224	895
2.225-2.249	935
2.250-2.274	978
2.275-2.299	1022
2.300-2.324	1068
2.325-2.349	1117
2.350-2.374	1167
2.375-2.399	1219
2.400-2.424	1275
2.425-2.449	1333
2.450-2.474	1393
2.475-2.499	1455
2.500-2.524	1522
2.525-2.549	1590
2.550 or greater	1662



MM-0248

Effective Date: 08/01/2020

E. Conditions of Coverage

F. Related Policies/Rules
Breast Reconstruction Surgery MM-0800

G. Review/Revision History

	DATE	ACTION
Date Issued	11/01/2018	
Date Revised	04/15/2020	Annual Update: Update to MCG Health 23 rd Edition Clinical Indications; addition of Related Policies.
Date Effective	09/01/2020	
Date Archived	07/01/2021	

H. References

- 1. Hansen, Juliana MD, Chang, Shilliang MD (January 14, 2020) Overview of Breast Reduction. Retrieved on April 8, 2020 from UpToDate.com
- 2. Hayes Technology Assessment. Reduction Mammoplasty (January 18, 2008). Retrieved on April 8, 2020 from www.Hayes.com
- 3. Knox, JA, Nelson, DA (January 08, 2018). US National Library of Medicine, National Institute of Health. Objective Effects of Breast Reduction on Physical Fitness. Retrieved on April 8, 2020 from www.ncbi.nlm.nih.gov
- 4. Perez-Panzano E., Gascon-Catalan A. (December 2017). US National Library of Medicine, National Institute of Health. Reduction mammoplasty improves levels of anxiety, depression and body image satisfaction in patients with symptomatic macromastia in the short and long term. Retrieved on April 8, 2020 from www.ncbi.nlm.nih.gov
- 5. Zeiderman, Matthew R. MD, Kelishadi, Shahrooz Sean, MD. (October 27, 2017)
 - US National Library of Medicine, National Institute of Health. Reduction Mammoplasty: Intraoperative Weight Versus Pathology Weight and Its Implications. Retrieved on April 9, 2020 from www.ncbi.nlm.nih.gov
- 6. MCG Health Guidelines (23rd Ed., 2020). Reduction Mammaplasty (Mammoplasty).
 - Retrieved on April 8, 2020 from www.MCG Health.com
- 7. American Society of Plastic Surgeons (ASPS). Evidence-based Clinical Practice Guideline: Reduction Mammaplasty (2011). Retrieved on April 8, 2020 from www.plasticsurgery.org
- 8. The Centers for Medicare & Medicaid Services (CMS). Women's Health and Cancer Rights Act (WHCRA 1998). Retrieved on April 8, 2020 from www.cms.gov





MM-0248 Effective Date: 08/01/2020

This guideline contains custom content that has been modified from the standard care guidelines and has not been reviewed or approved by MCG Health, LLC.

The Medical Policy Statement detailed above has received due consideration as defined in the Medical Policy Statement Policy and is approved.

Independent medical review - March 2018

