



MEDICAL POLICY STATEMENT

Georgia Medicaid

Policy Name & Number	Date Effective
Sacroiliac Joint Fusion-GAMCD-MM-1040	08/01/2022-03/31/2023
Policy Type	
MEDICAL	

Medical Policy Statement prepared by CareSource and its affiliates are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other policies and procedures.

Medical Policy Statements prepared by CareSource and its affiliates do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage) for the service(s) referenced in the Medical Policy Statement. If there is a conflict between the Medical Policy Statement and the plan contract (i.e., Evidence of Coverage), then the plan contract (i.e., Evidence of Coverage) will be the controlling document used to make the determination. According to the rules of Mental Health Parity Addiction Equity Act (MHPAEA), coverage for the diagnosis and treatment of a behavioral health disorder will not be subject to any limitations that are less favorable than the limitations that apply to medical conditions as covered under this policy.

Table of Contents

A. Subject.....	2
B. Background.....	2
C. Definitions.....	2
D. Policy.....	3
E. Conditions of Coverage.....	5
F. Related Polices/Rules.....	5
G. Review/Revision History.....	5
H. References.....	5

A. Subject
Sacroiliac Joint Fusion

B. Background

The sacroiliac joints (SIJ) are formed by the connection of the sacrum and the right and left iliac bones. The sacrum is the triangular-shaped bone in the lower portion of the spine, below the lumbar spine. While most of the vertebrae of the spine are mobile, the sacrum is made up of five vertebrae that are fused together and do not move. The iliac bones are the two large bones that make up the pelvis. As a result, the SI joints connect the spine to the pelvis. The sacrum and the iliac bones (ileum) are held together by a collection of strong ligaments. There is relatively little motion at the SI joints - normally less than 4° of rotation and 2 mm of translation.

SIJ dysfunction occurs when there is abnormal movement or malalignment of the sacroiliac joint and is the main source of lower back pain in 15% to 30% of patients. The condition causes disability and pain, and may be caused by prior lumbar sacral fusion, trauma, inflammatory arthritis, sacral tumors, osteoarthritis, or pregnancy. Patients may present with low back, groin, and/or gluteal pain. SIJ pain is often similar to discogenic pain or radicular back pain. This can lead to misdiagnosis and treatment; imaging studies and physical exam are usually necessary to determine if pain is caused by SIJ dysfunction. For many individuals, conservative management and/or minimally invasive procedures can improve pain management. However, joint fusion may be warranted following trauma or failure of conservative management and less invasive procedures (e.g., SIJ injections).

Open SIJ fusion is typically performed when a large visual field is required (e.g., post-traumatic injury, tumor resection), while percutaneous SIJ fusion may be used for treatment of refractory chronic low back pain. Open SIJ fusion typically involves opening the SIJ, denuding of cartilage, and bone grafting. To stabilize the SIJ, the iliac crest bone and the sacrum are typically held together by plates or screws or an interbody fusion cage until the bones fuse. The minimally invasive procedure for SIJ fusion is performed by an orthopedic or neurologic surgeon in an inpatient or outpatient setting. The procedure typically ranges from 45 to 70 minutes and requires general endotracheal anesthesia, fluoroscopic guidance, and a small (approximately 3 mm) incision in the buttock region. Postoperatively, patients ambulate with a walker or crutches and follow a progressive regimen to develop flexibility and strength until they are fully ambulatory.

C. Definitions

- **Conservative Therapy** - A multimodality plan of care. Multimodality care plans include both active and inactive conservative therapies.
 - **Active Conservative Therapies** - Include physical therapy, occupational therapy, and/or a physician supervised home exercise program (HEP).
 - **Inactive Conservative Therapies** - Include rest, ice, heat, medical devices, transcutaneous electrical nerve stimulation (TENS) unit, and/or prescription medications.

- **Transcutaneous Electric Nerve Stimulator (TENS Unit)** - A durable medical equipment device dispensed by prescription. Its use, frequency, duration, and start dates must be documented in the medical record to be considered part of conservative therapy during the period of prior authorization request.

D. Policy

I. Sacroiliac Joint Fusion

A. CareSource considers open sacroiliac joint (SIJ) fusion medically necessary when **ALL** the following criteria are met:

1. **At least ONE** of the following applies:
 - a. Patient has post-traumatic injury of the SIJ (e.g., following pelvic ring fracture);
 - b. The procedure will be performed as an adjunctive treatment for sacroiliac joint infection (e.g., osteomyelitis, pyogenic sacroiliitis)/sepsis;
 - c. The procedure will be performed for management of sacral tumor (e.g., partial sacrectomy);
 - d. The procedure will be performed as part of a multisegmental long fusion construct for the correction of spinal deformity (e.g., idiopathic scoliosis, neuromuscular scoliosis);
 - e. Prior percutaneous SIJ fusion has failed;
2. Recent (within 6 months) plain x-rays and/or cross-sectional imaging (CT or MRI) demonstrate localized SIJ pathology.

B. CareSource considers percutaneous SIJ fusion/stabilization for the treatment of back pain medically necessary when **ALL** the following criteria are met:

1. Presence of non-radiating pain that is caudal to the lumbar spine (L5 vertebrae), localized over the posterior SIJ, and consistent with SIJ pain;
2. SIJ pain severe enough to interfere with activities of daily living;
3. Localized tenderness with palpation of the posterior SIJ in the absence of tenderness of similar severity elsewhere (e.g. greater trochanter, lumbar spine, coccyx) and other obvious sources for their pain do not exist.
4. The patient has undergone and failed a minimum 6 months of conservative therapy, including **ALL** the following:
 - a. **ACTIVE** conservative therapy as part of a multimodality comprehensive approach and is addressed in the patient's care plan with documentation in the medical record that includes **ONE** of the following:
 01. The patient has received **ACTIVE** conservative therapy lasting for 6 months or more within the past 12 months including **at least ONE** of the following:
 - (1).Physical therapy;
 - (2).Occupational therapy;
 - (3).A physician supervised home exercise program (HEP), including the following two requirements:
 - i. An exercise prescription and/or plan documented in the medical record;

II. Exclusions

A. Open SIJ fusion is considered not medically necessary for any other indication not outlined above, including (but not limited to) the following:

1. Mechanical low back syndrome.
2. Sacroiliac joint syndrome.
3. Degenerative sacroiliac joint.
4. Presence of neural compression as seen on an MRI or CT that correlates with the patient's symptoms or other more likely source for their pain (e.g., radicular pain).

A. Percutaneous SIJ fusion for SIJ pain is NOT indicated in the presence of:

1. Systemic arthropathy such as ankylosing spondylitis or rheumatoid arthritis.
2. Generalized pain behavior (e.g. somatoform disorder) or generalized pain disorder (e.g. fibromyalgia).
3. Infection, tumor, or fracture.
4. Acute, traumatic instability of the SIJ.
5. Presence of neural compression as seen on an MRI or CT that correlates with the patient's symptoms or other more likely source for their pain.

E. Conditions of Coverage

NA

F. Related Polices/Rules

NA

G. Review/Revision History

	DATE	ACTION
Date Issued	05/13/2020	New Policy
Date Revised	04/28/2021 04/12/2022	Annual Update: Removed PA language Annual Review: added open SIJ fusion indications, re-organized percutaneous criteria, added references
Date Effective	08/01/2022	
Date Archived	03/31/2023	This Policy is no longer active and has been archived. Please note that there could be other Policies that may have some of the same rules incorporated and CareSource reserves the right to follow CMS/State/NCCI guidelines without a formal documented Policy.

H. References

1. Chou R. (2021, June 10). Subacute and chronic low back pain: nonsurgical interventional treatment. UpToDate. Retrieved April 6, 2022 from www.uptodate.com.
2. Chou R. (2021, June 11). Subacute and chronic low back pain: surgical treatment. UpToDate. Retrieved April 6, 2022 from www.uptodate.com.

3. DePhillipo NN, Corenman DS, Strauch EL, Zalepa King LA. Sacroiliac pain: structural causes of pain referring to the SI joint region. Clin Spin Surg. 2019 Jul;32(6):E282-E288. doi:10.1097/BSD.0000000000000745.
4. Graham Smit A, Capobianco R, Cher D, et al. Open versus minimally invasive sacroiliac fusion: a multi-center comparison of perioperative measures and clinical outcomes. Ann Surg Innov Res. 2013;7(1):14. doi:10.1186/1750-1164-7-14.
5. Health Technology Assessment. (2020 September 3). Minimally invasive sacroiliac joint fusion using triangular titanium implants (iFuse Implant System, SI-Bone Inc.). Hayes. Retrieved April 5, 2022 from www.evidence.hayesinc.com.
6. Lorio MP. ISASS Policy 2016 update – minimally invasive sacroiliac joint fusion. Int J Spine Surg. 2016;10:26. doi:10.14444/3026.
7. Lorio M, Kube R, Araghi A. International Society for the Advancement of Spine Surgery Policy 2020 Update - Minimally invasive surgical sacroiliac joint fusion (for chronic sacroiliac joint pain): coverage indications, limitations, and medical necessity. Int J Spine Surg. 2020:7156. doi:10.14444/7156.
8. National Institute for Health and Care Excellence. Minimally invasive sacroiliac joint fusion surgery for chronic sacroiliac pain [IPG578]. 2017;1-9. Retrieved March 24, 2022 from www.nice.org.
9. Unoki E, Abe E, Murai H, et al. Fusion of multiple segments can increase the incidence of sacroiliac joint pain after lumbar or lumbosacral fusion. Spine. 2016;41(12):999–1005. doi:10.1097/BRS.0000000000001409.
10. Unoki E, Miyakoshi N, Abe E, et al. Sacroiliac joint pain after multiple-segment lumbar fusion: a long-term observational study - non-fused sacrum vs. fused sacrum. Spine Surg Relat Res. 2017;1(2):90–95. doi:10.22603/ssr.1.2016-0010.
11. Unoki E, Miyakoshi N, Abe E, et al. Sacropelvic fixation with S2 alar iliac screws may prevent sacroiliac joint pain after multisegment spinal fusion. Spine. 2019;44(17):E1024-E1030. doi:10.1097/BRS.0000000000003041.
12. Zaidi HA, Montoure AJ, Dickman CA. Surgical and clinical efficacy of sacroiliac joint fusion: a systematic review of the literature. J Neurosurg Spine. April 2015; 23(1): 59-66. doi:10.3171/2014.10.SPINE14516.

Independent medical review – May 2020

GA-MED-P-1271918

Issue Date 05/13/2020

Approved DCH 04/29/2022