

PHARMACY POLICY STATEMENT

Georgia Medicaid

DRUG NAME	Rituxan (rituximab)
BILLING CODE	J9312
BENEFIT TYPE	Medical
SITE OF SERVICE ALLOWED	Office/Outpatient
COVERAGE REQUIREMENTS	Prior Authorization Required (Non-Preferred Product) QUANTITY LIMIT—see “Dosage Allowed” sections
LIST OF DIAGNOSES CONSIDERED NOT MEDICALLY NECESSARY	Click Here

Rituxan (rituximab) is a **non-preferred** product and will only be considered for coverage under the **medical** benefit when the following criteria are met:

Members must be clinically diagnosed with one of the following disease states and meet their individual criteria as stated.

GRANULOMATOSIS WITH POLYANGIITIS (GPA) (WEGENER’S GRANULOMATOSIS) AND MICROSCOPIC POLYANGIITIS (MPA)

For **initial** authorization:

1. Member is 2 years old or older; AND
2. Medication must be prescribed by or in consultation with a nephrologist or rheumatologist; AND
3. Member has a confirmed diagnosis of severe GPA or MPA, **or** non-severe disease (non-organ threatening, non-life-threatening) refractory to glucocorticoids in combination with methotrexate; AND
4. Rituxan will be initiated in combination with glucocorticoids; AND
5. Member has at least ONE of the following:
 - a) Member’s disease remains active or has progressed despite at least a 3 month trial of glucocorticoids in combination with cyclophosphamide;
 - b) Further treatment with cyclophosphamide would exceed the maximum cumulative dose;
 - c) Cyclophosphamide is contraindicated or not tolerated by the member.
6. **Dosage allowed:** Please refer to the Dosing and Administration section of the package insert.

If member meets all the requirements listed above, the medication will be approved for 6 months.

For **reauthorization**:

1. Member tolerates infusions; AND
2. Chart notes demonstrate clinical improvement of disease signs and symptoms.

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

PEMPHIGUS VULGARIS (PV)

For **initial** authorization:

1. Member is 18 years old or older; AND
2. Must be prescribed by or in consultation with a dermatologist; AND
3. Member has a documented diagnosis of moderate to severe PV; AND
4. Rituxan will be initiated in combination with a corticosteroid taper (unless contraindicated); AND
5. Member has tried and failed or has contraindication to high dose corticosteroid (equivalent to 1.5mg/kg/day prednisone) and an adjuvant immunosuppressive agent such as azathioprine or mycophenolate mofetil.
6. **Dosage allowed:** Initial: Two 1000mg doses separated by 2 weeks; Maintenance: 500mg infusion at month 12 and every 6 months thereafter or based on clinical evaluation -- no sooner than 16 weeks following the previous infusion; Relapse: 1000mg infusion.

If member meets all the requirements listed above, the medication will be approved for 12 months.

For **reauthorization**:

1. Member tolerates infusions; AND
2. Chart notes demonstrate clinical improvement of signs and symptoms (e.g. healed lesions, fewer new lesions, etc.)

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

RHEUMATOID ARTHRITIS (RA)

For **initial** authorization:

1. Member is 18 years old or older; AND
2. Medication is being prescribed by or in consultation with a rheumatologist; AND
3. Member has a documented diagnosis of moderately- to severely- active RA; AND
4. Rituxan is being used in combination with methotrexate, unless unable to tolerate; AND
5. Member must have inadequate response or intolerance to one or more tumor necrosis factor (TNF) antagonists (e.g. adalimumab, etanercept, infliximab) for at least 3 months each. Note: TNF antagonists require prior authorization.
6. **Dosage allowed:** Two 1000mg doses separated by 2 weeks; subsequent courses repeated no sooner than every 16 weeks (every 24 weeks is typical).

If member meets all the requirements listed above, the medication will be approved for 6 months.

For **reauthorization**:

1. Member tolerates infusions; AND
2. Chart notes demonstrate improvement of RA signs and symptoms (e.g. fewer number of painful and swollen joints, achievement of remission, etc.)

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

NON-HODGKIN'S LYMPHOMA (NHL)

These requests must be submitted through [NantHealth/Eviti](#) portal.

CHRONIC LYMPHOCYTIC LEUKEMIA (CLL)

These requests must be submitted through [NantHealth/Eviti](#) portal.

CareSource considers Rituxan (rituximab) not medically necessary for the treatment of the diseases that are not listed in this document.

DATE	ACTION/DESCRIPTION
6/9/2020	New policy for Rituxan created.

References:

- Rituxan [package insert]. South San Francisco, CA: Genentech, Inc.; 2020.
- Ntatsaki E, Carruthers D, Chakravarty K, et al. BSR and BHPR guideline for the management of adults with ANCA-associated vasculitis. *Rheumatology* April 2014: ket445
- Stone JH, Merkel PA, Spiera R, et al. Rituximab versus cyclophosphamide for ANCA-associated vasculitis. *N Engl J Med* 2010; 363:221.
- Jones RB, Tervaert JW, Hauser T, et al. Rituximab versus cyclophosphamide in ANCA-associated renal vasculitis. *N Engl J Med* 2010; 363:211.
- Jones RB, Tervaert JW, Hauser T, et al. Rituximab versus cyclophosphamide in ANCA-associated renal vasculitis: 2-year results of a randomized trial. *Ann Rheum Dis* 2015; 74(6): 1178-1182.
- Latimer NR, Carroll C, Wong R, et al. Rituximab in combination with corticosteroids for the treatment of anti-neutrophil cytoplasmic antibody-associated vasculitis: a NICE single technology appraisal. *Pharmacoeconomics* 2014; 32(12): 1171-1183.
- Pagnoux C. Updates in ANCA-associated vasculitis. *Eur J Rheumatol* 2015
- Singh JA, Saag KG, Bridges Jr. SL, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care & Research* 2015: 1-25.
- Leandro MJ. Rituximab: Principles of use and adverse effects in rheumatoid arthritis. *UpToDate*. https://www.uptodate.com/contents/rituximab-principles-of-use-and-adverse-effects-in-rheumatoid-arthritis?search=rheumatoid%20arthritis%20treatment&topicRef=7966&source=see_link. Updated April 3, 2020. Accessed June 9, 2020.
- Finckh A, Ciurea A, Brulhart L, et al. Which subgroup of patients with rheumatoid arthritis benefits from switching to rituximab versus alternative anti-tumour necrosis factor (TNF) agents after previous failure of an anti-TNF agent? *Annals of the Rheumatic Diseases*. 2009;69(2):387-393. doi:10.1136/ard.2008.105064
- Solau-Gervais E, Prudhomme C, Philippe P, et al. Efficacy of rituximab in the treatment of rheumatoid arthritis. Influence of serologic status, coprescription of methotrexate and prior TNF-alpha inhibitors exposure. *Joint Bone Spine*. 2012;79(3):281-284. doi:10.1016/j.jbspin.2011.05.002
- Harrold LR, Reed GW, Magner R, et al. Comparative effectiveness and safety of rituximab versus subsequent anti-tumor necrosis factor therapy in patients with rheumatoid arthritis with prior exposure to anti-tumor necrosis factor therapies in the United States Corrona registry. *Arthritis Research & Therapy*. 2015;17(1). doi:10.1186/s13075-015-0776-1
- Chatzidionysiou K, Lie E, Nasonov E, et al. Highest clinical effectiveness of rituximab in autoantibody-positive patients with rheumatoid arthritis and in those for whom no more than one previous TNF antagonist has failed: pooled data from 10 European registries. *Annals of the Rheumatic Diseases*. 2011;70(9):1575-1580. doi:10.1136/ard.2010.148759
- Emery P, Gottenberg JE, Rubbert-Roth A, et al. Rituximab versus an alternative TNF inhibitor in patients with rheumatoid arthritis who failed to respond to a single previous TNF inhibitor: SWITCH-RA, a global, observational, comparative effectiveness study. *Annals of the Rheumatic Diseases*. 2014;74(6):979-984. doi:10.1136/annrheumdis-2013-203993
- Hertl M, Eming R. Management of refractory pemphigus vulgaris and pemphigus foliaceus. *UpToDate*. https://www.uptodate.com/contents/management-of-refractory-pemphigus-vulgaris-and-pemphigus-foliaceus?search=pemphigus%20vulgaris&source=search_result&selectedTitle=3~40&usage_type=default&display_rank=3. Updated March 5, 2020. Accessed June 11, 2020.
- Heelan K, Al-Mohammed F, Smith MJ, et al. Durable Remission of Pemphigus With a Fixed-Dose Rituximab Protocol. *JAMA Dermatology*. 2014;150(7):703. doi:10.1001/jamadermatol.2013.6739
- Murrell DF, Dick S, Ahmed A, et al. Consensus statement on definitions of disease, end points, and therapeutic response for pemphigus. *Journal of the American Academy of Dermatology*. 2008;58(6):1043-1046. doi:10.1016/j.jaad.2008.01.012

18. Agarwal A, Hall RP, Bañez LL, Cardones AR. Comparison of rituximab and conventional adjuvant therapy for pemphigus vulgaris: A retrospective analysis. *Plos One*. 2018;13(9). doi:10.1371/journal.pone.0198074
19. Merkel PA, Kaplan AA, Falk RJ. Granulomatosis with polyangiitis and microscopic polyangiitis: Initial immunosuppressive therapy. UpToDate. https://www.uptodate.com/contents/granulomatosis-with-polyangiitis-and-microscopic-polyangiitis-initial-immunosuppressive-therapy?search=Granulomatosis%20with%20polyangiitis%20and%20Microscopic%20Polyangiitis&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2. Updated January 23, 2019. Accessed June 11, 2020.
20. McGeoch L, Twilt M, Famarca L, et al. CanVasc Recommendations for the Management of Antineutrophil Cytoplasm Antibody-associated Vasculitides. *The Journal of Rheumatology*. 2015;43(1):97-120. doi:10.3899/jrheum.150376
21. Yates M, Watts R, Bajema I, et al. OP0053 Eular/ERA-EDTA Recommendations for The Management of Anca-Associated Vasculitis. *Annals of the Rheumatic Diseases*. 2016;75(Suppl 2). doi:10.1136/annrheumdis-2016-eular.1168
22. Terrier B, Pagnoux C, Perrodeau É, et al. Long-term efficacy of remission-maintenance regimens for ANCA-associated vasculitides. *Annals of the Rheumatic Diseases*. 2018;77(8):1150-1156. doi:10.1136/annrheumdis-2017-212768

Effective date: 07/20/2020

Revised date: 06/09/2020