

PHARMACY POLICY STATEMENT

Georgia Medicaid

DRUG NAME	Skyrizi (risankizumab-rzaa)
BILLING CODE	Must use valid NDC code
BENEFIT TYPE	Pharmacy
SITE OF SERVICE ALLOWED	Home/Office
COVERAGE REQUIREMENTS	Prior Authorization Required (Non-Preferred Product) Alternative preferred products include Cimzia, Cosentyx, Enbrel, Otezla and Siliq QUANTITY LIMIT— see Dosage allowed below
LIST OF DIAGNOSES CONSIDERED NOT MEDICALLY NECESSARY	Click Here

Skyrizi (risankizumab-rzaa) is a **non-preferred** product and will only be considered for coverage under the **pharmacy** benefit when the following criteria are met:

Members must be clinically diagnosed with one of the following disease states and meet their individual criteria as stated.

PLAQUE PSORIASIS (PsO)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Must have a documented negative TB test (i.e., tuberculosis skin test (PPD), an interferon-release assay (IGRA)) within 12 months prior to starting therapy; AND
3. Medication must be prescribed by a rheumatologist or dermatologist; AND
4. Member has PsO for 6 months or longer; AND
5. Member has PsO involves 10% or more of the member's body surface area (BSA); AND
6. Member's Psoriasis Area and Severity Index (PASI) score is greater than or equal to 12; AND
7. Member has tried and failed to respond to treatment with at least one of the following:
 - a) At least 12 weeks of photochemotherapy (i.e., psoralen plus ultraviolet A therapy);
 - b) At least 12 weeks of phototherapy (i.e., UVB light therapy, Excimer laser treatments) (tanning beds emit mostly UVA light and therefore would not meet this criteria);
 - c) At least a 4 week trial with topical antipsoriatic agents (i.e., anthralin, calcipotriene, coal tar, corticosteroids, tazarotene);
 AND
8. Member has tried and failed to respond to treatment with traditional first-line oral/systemic therapies (i.e., cyclosporine, methotrexate, acitretin) for at least a 12 week trial; AND
9. Member has tried and failed treatment with at least **two** of the following: Cimzia, Cosentyx, Enbrel, Otezla and Siliq. Treatment failure requires at least 12 weeks of therapy with each drug.
10. **Dosage allowed:** 150 mg (two 75 mg injections) administered by subcutaneous injection at week 0, week 4, and every 12 weeks thereafter.

If member meets all the requirements listed above, the medication will be approved for 12 months.

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease (e.g., documented member's PASI score improvement, etc.).

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

CareSource considers Skyrizi (risankizumab-rzaa) not medically necessary for the treatment of the following disease states based on a lack of robust clinical controlled trials showing superior efficacy compared to currently available treatments:

- Active infections
- Ankylosing spondylitis
- Asthma
- Cellulitis
- Crohn's Disease
- Dissecting scalp cellulitis
- For use in combination with other TNF-inhibitors (i.e., Humira, Kineret, Enbrel, Remicade)
- Giant-cell arteritis
- Infectious uveitis
- Juvenile idiopathic arthritis
- Lupus perino
- Osteoarthritis
- Psoriatic Arthritis
- Recurrent pregnancy loss
- Relapsing polychondritis
- Rheumatoid arthritis
- Sarcoidosis
- Sciatica
- Spondyloarthritis (other than ankylosing spondylitis)
- Takayasu's arteritis
- Ulcerative Colitis
- Vogt-Koyanagi

DATE	ACTION/DESCRIPTION
07/28/2019	New policy for Skyrizi created.

References:

1. Skyrizi [prescribing information]. North Chicago, IL: AbbVie Inc.; April 2019.
2. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6. Guidelines of care for the treatment of psoriasis and psoriatic arthritis: Case-based presentations and evidence-based conclusions. Journal of the American Academy of Dermatology, Volume 65, Issue 1, 137 – 174.
3. Hsu S, Papp KA, Lebwohl MG, et al. Consensus guidelines for the management of plaque psoriasis. Arch Dermatol. 2012 Jan;148(1):95-102.
4. ClinicalTrials.gov. Identifier: NCT02684370. BI 655066 (Risankizumab) Compared to Placebo and Active Comparator (Ustekinumab) in Patients With Moderate to Severe Chronic Plaque Psoriasis. Available at: <https://clinicaltrials.gov/ct2/show/NCT02684370?term=ULTIMMA-1&rank=1>.

5. ClinicalTrials.gov. Identifier: NCT02684357. BI 655066 Versus Placebo & Active Comparator (Ustekinumab) in Patients With Moderate to Severe Chronic Plaque Psoriasis. Available at: <https://clinicaltrials.gov/ct2/show/NCT02684357?term=ULTIMMA-2&rank=1>.
6. Gottlieb AB, et al. Safety observations in 12095 patients with psoriasis enrolled in an international registry (PSOLAR): experience with infliximab and other systemic and biologic therapies. *J Drugs Dermatol*. 2014 Dec;13(12):1441-8.
7. Sbidian E, et al. Systemic pharmacological treatments for chronic plaque psoriasis: a network meta-analysis. *Cochrane Database Syst Rev*. 2017;12:CD011535.
8. Nast A, et al. European S3-Guideline on the systemic treatment of psoriasis vulgaris - Update Apremilast and Secukinumab - EDF in cooperation with EADV and IPC. *J Eur Acad Dermatol Venereol*. 2017;31(12):1951.

Effective date: 09/26/2019

Revised date: 07/28/2019