

# PHARMACY POLICY STATEMENT

## Georgia Medicaid

<b>DRUG NAME</b>	<b>Tremfya (guselkumab)</b>
BENEFIT TYPE	Pharmacy or Medical
STATUS	Prior Authorization Required

Tremfya (guselkumab) is an anti-psoriatic agent, interleukin-23 inhibitor, and monoclonal antibody initially approved by the FDA in 2017. It is currently FDA approved for the treatment of moderate to severe plaque psoriasis in adults who are candidates for systemic therapy or phototherapy, psoriatic arthritis, ulcerative colitis and Crohn’s disease in adults.

Tremfya (guselkumab) will be considered for coverage when the following criteria are met:

### Plaque Psoriasis (PsO)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Medication must be prescribed by or in consultation with a dermatologist; AND
3. Member has clinical documentation of moderate to severe plaque psoriasis characterized by 3% or more of body surface area (BSA) or disease affecting sensitive areas (e.g., hands, feet, face, genitals, etc.); AND
4. Member has tried and failed to respond to treatment with at least **ONE** of the following:
  - a) At least 12 weeks of photochemotherapy (i.e., psoralen plus ultraviolet A therapy);
  - b) At least 12 weeks of phototherapy (i.e., UVB light therapy, Excimer laser treatments);
  - c) At least a 4 week trial with topical antipsoriatic agents (i.e., anthralin, calcipotriene, coal tar, corticosteroids, tazarotene, tacrolimus, pimecrolimus); AND
5. Member has tried and failed, or unable to tolerate a systemic conventional DMARD (i.e., cyclosporine, methotrexate, acitretin) for at least 12 weeks; AND
6. Member has tried and failed at least **TWO** preferred biologic DMARDs for at least 3 months each; AND
7. Member has had a negative tuberculosis test within the past 12 months.
8. **Dosage allowed/Quantity limit:** 100 mg administered by subcutaneous injection at Week 0, Week 4 and every 8 weeks thereafter. Quantity limit: 1 syringe every 8 weeks after loading doses.

***If all the above requirements are met, the medication will be approved for 12 months.***

For **reauthorization**:

1. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease such as documented member’s BSA improvement, etc.

***If all the above requirements are met, the medication will be approved for an additional 12 months.***

### Psoriatic Arthritis (PsA)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Medication must be prescribed by or in consultation with a rheumatologist or dermatologist; AND
3. Member has a documented diagnosis of active psoriatic arthritis (PsA); AND
4. Member has met a 4-week trial of an NSAID taken at maximally tolerated doses AND a 3-month trial of a conventional DMARD agent (e.g., methotrexate, sulfasalazine, cyclosporine, etc.) unless **ONE** of the following situations is met:
  - a) Conventional DMARD is **NOT** required for:
    - i) Concomitant axial disease (i.e., involving sacroiliac joint and spine) or enthesitis; OR
  - b) NSAID and conventional DMARD are **NOT** required for:
    - i) Severe PsA (defined as having at least one of the following: erosive disease, active PsA at many sites including dactylitis or enthesitis, elevated levels of ESR or CRP, joint deformities, or major impairment in quality of life); AND
5. Member has tried and failed at least **TWO** preferred biologic DMARDs for at least 3 months each; AND
6. Member has had a negative tuberculosis test within the past 12 months.
7. **Dosage allowed/Quantity limit:** 100 mg administered by subcutaneous injection at Week 0, Week 4 and every 8 weeks thereafter. Quantity limit: 1 syringe every 8 weeks after loading doses.

***If all the above requirements are met, the medication will be approved for 12 months.***

For **reauthorization**:

1. Chart notes have been provided that show improvement of signs and symptoms of disease such as decreased joint swelling and pain, improved skin appearance, improved quality of life, etc.

***If all the above requirements are met, the medication will be approved for an additional 12 months.***

## Ulcerative Colitis (UC)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Medication must be prescribed by or in consultation with a gastroenterologist; AND
3. Member has a documented diagnosis of moderate to severely active UC; AND
4. Member must have a documented trial and inadequate response with **ONE** of the following:
  - a) 6-mercaptopurine or azathioprine;
  - b) Corticosteroid (e.g., budesonide, prednisone, methylprednisolone);
  - c) 5-aminosalicylate (e.g., Asacol HD, Lialda, Pentasa, Delzicol, mesalamine, etc.); AND
5. Member has had a negative tuberculosis test within the past 12 months.
6. **Dosage allowed/Quantity limit:**
  - a) 200 mg administered by intravenous infusion at Week 0, Week 4, and Week 8 then 100 mg administered by subcutaneous injection at Week 16, and every 8 weeks thereafter, or 200 mg administered by subcutaneous injection at Week 12, and every 4 weeks thereafter.
  - b) Quantity limit:
    - i) 100 mg/mL: 1 syringe/injector per 56 days after loading doses.
    - ii) 200 mg/mL: 1 pen/syringe per 28 days after loading doses.

***If all the above requirements are met, the medication will be approved for 4 months.***

For **reauthorization**:

1. Chart notes have been provided showing improvement of signs and symptoms of disease such as clinical remission, reduced rectal bleeding, decreased stool frequency, or endoscopic-histologic mucosal healing.

***If all the above requirements are met, the medication will be approved for an additional 12 months.***

## Crohn's Disease (CD)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Medication must be prescribed by or in consultation with a gastroenterologist; AND
3. Member has a documented diagnosis of moderate to severely active CD; AND
4. Member has had a documented trial and inadequate response, or intolerance to **ONE** of the following conventional therapies:
  - a) Corticosteroid;
  - b) 6-mercaptopurine, azathioprine, or methotrexate; OR
5. Provider attests member has severe disease that requires immediate use of an advanced therapy (biologic, JAK inhibitor, etc.) agent such as penetrating or fistulizing disease, multiple resections, etc.; AND
6. Member has baseline liver function tests completed or scheduled; AND
7. Member has had a negative tuberculosis test within the past 12 months.
8. **Dosage allowed/Quantity limit:**
  - a) Initial dose of 200 mg administered by intravenous infusion over at least one hour at Week 0, Week 4, and Week 8 or 400 mg administered by subcutaneous injection at Week 0, Week 4, and Week 8.
  - b) Maintenance dose of 100 mg administered by subcutaneous injection at Week 16, and every 8 weeks thereafter, or 200 mg administered by subcutaneous injection at Week 12, and every 4 weeks thereafter. Quantity limit: for the 100 mg/mL, one patient-controlled injector, pen or syringe per 56 days. For the 200 mg/mL, one pen or syringe per 28 days.

***If all the above requirements are met, the medication will be approved for 4 months.***

For **reauthorization**:

1. Chart notes have been provided that show improvement of signs and symptoms of disease such as improved endoscopic response, fewer flare-ups of symptoms, improved quality of life, etc.

***If all the above requirements are met, the medication will be approved for an additional 12 months.***

**CareSource considers Tremfya (guselkumab) not medically necessary for the treatment of conditions that are not listed in this document. For any other indication, please refer to the Off-Label policy.**

DATE	ACTION/DESCRIPTION
10/19/2017	New policy for Tremfya created.
02/26/2019	Humira was removed from criteria; Cimzia, Cosentyx, Otezla and Siliq added to trial agents list. Initial authorization length increased to 12 months. TB test allowed to be done within 12 months prior to initiation of therapy; chest x-ray option removed. Reauthorization criteria on documented member's PASI score improvement incorporated into general chart noted documentation requirements.
09/23/2020	New indication Psoriatic Arthritis added. For PsO Removed rheumatologist from prescriber. Removed PsO for 6 months or longer. Changed BSA to 3% or sensitive area involvement. Removed PASI score requirement. Removed repeated TB test for reauthorization.
02/22/2022	Transferred policy to new format; Removed initial criteria from reauthorization; Updated wording for biologic DMARDs; Clarified reauth criteria
05/15/2024	Added medical benefit option.

<b>10/01/2024</b>	Added ulcerative colitis indication including quantity limit. Simplified tuberculosis test requirement.
<b>04/15/2025</b>	Updated references; added Crohn's disease indication
<b>08/19/2025</b>	Updated references. UC: removed trial durations CD: removed trial option of biologic with CD indication and added alternative option of provider attestation that member has severe disease that requires immediate use of advanced therapy PsO and PsA: replaced "non-biologic" with "conventional"

References:

1. Tremfya [package insert]. Horsham, PA: Janssen Biotech, Inc.; 2025.
2. Elmets CA, Korman NJ, Prater EF, et al. Joint AAD-NPF Guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures [published online ahead of print, 2020 Jul 30]. *J Am Acad Dermatol*. 2020;S0190-9622(20)32288-X.
3. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol*. 2020;82(6):1445-1486.
4. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019;80(4):1029-1072.
5. Elmets CA, Lim HW, Stoff B, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management and treatment of psoriasis with phototherapy [published correction appears in *J Am Acad Dermatol*. 2020 Mar;82(3):780]. *J Am Acad Dermatol*. 2019;81(3):775-804.
6. Menter A, Cordoro KM, Davis DM, et al. Joint AAD-NPF Guidelines of care for the management and treatment of psoriasis in pediatric patients. *J Am Acad Dermatol* 2020;82:161-201
7. Deodhar A, Helliwell PS, Boehncke WH, et al. Guselkumab in patients with active psoriatic arthritis who were biologic-naïve or had previously received TNF $\alpha$  inhibitor treatment (DISCOVER-1): a double-blind, randomised, placebo-controlled phase 3 trial [published correction appears in *Lancet*. 2020 Apr 4;395(10230):1114. doi: 10.1016/S0140-6736(20)30680-2]. *Lancet*. 2020;395(10230):1115-1125. doi:10.1016/S0140-6736(20)30265-8
8. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the treatment of Psoriatic Arthritis. *Arthritis Rheumatol*. 2019 Jan;71(1):5-32.
9. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021 [published correction appears in *Nat Rev Rheumatol*. 2022 Dec;18(12):734. doi: 10.1038/s41584-022-00861-w]. *Nat Rev Rheumatol*. 2022;18(8):465-479. doi:10.1038/s41584-022-00798-0
10. Gossec L, Kerschbaumer A, Ferreira RJO, et al. EULAR recommendations for the management of psoriatic arthritis with pharmacological therapies: 2023 update. *Ann Rheum Dis*. 2024;83(6):706-719. Published 2024 May 15. doi:10.1136/ard-2024-225531
11. Gordon H, Minozzi S, Kopylov U, et al. ECCO Guidelines on Therapeutics in Crohn's Disease: Medical Treatment. *J Crohns Colitis*. 2024;18(10):1531-1555. doi:10.1093/ecco-jcc/jjae091
12. Feuerstein JD, Ho EY, Shmidt E, et al. AGA Clinical Practice Guidelines on the Medical Management of Moderate to Severe Luminal and Perianal Fistulizing Crohn's Disease. *Gastroenterology*. 2021;160(7):2496-2508. doi:10.1053/j.gastro.2021.04.022
13. Lichtenstein GR, Loftus EV, Afzali A, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol*. 2025;120(6):1225-1264. Published 2025 Jun 3. doi:10.14309/ajg.0000000000003465
14. Singh S, Loftus EV Jr, Limketkai BN, et al. AGA Living Clinical Practice Guideline on Pharmacological Management of Moderate-to-Severe Ulcerative Colitis. *Gastroenterology*. 2024;167(7):1307-1343. doi:10.1053/j.gastro.2024.10.001
15. Rubin D, Ananthakrishnan A, Siegel C, et al. ACG Clinical Guideline Update: Ulcerative Colitis in Adults. *The American Journal of Gastroenterology* 120(6):p 1187-1224, June 2025. | DOI: 10.14309/ajg.00000000000003



Effective date: 05/01/2026  
Revised date: 08/19/2025