

## PHARMACY POLICY STATEMENT

### Georgia Medicaid

<b>DRUG NAME</b>	Xolair (omalizumab)
<b>BILLING CODE</b>	J2357 (1 unit = 5 mg)
<b>BENEFIT TYPE</b>	Medical
<b>SITE OF SERVICE ALLOWED</b>	Office/Outpatient Hospital
<b>COVERAGE REQUIREMENTS</b>	Prior Authorization Required (Preferred Product) QUANTITY LIMIT – see dosage allowed
<b>LIST OF DIAGNOSES CONSIDERED NOT MEDICALLY NECESSARY</b>	<a href="#">Click Here</a>

**Xolair (omalizumab) is a preferred product and will only be considered for coverage under the medical benefit when the following criteria are met:**

Members must be clinically diagnosed with one of the following disease states and meet their individual criteria as stated.

### CHRONIC RHINOSINUSITIS WITH NASAL POLYPOSIS (CRSwNP)

For initial authorization:

1. Member is 18 years old or older; AND
2. Medication must be prescribed by or in consultation with allergist, immunologist, or otorhinolaryngologist (ENT); AND
3. Member has a diagnosis of bilateral CRSwNP for more than 12 weeks; AND
4. Chart notes must show documentation of bilateral nasal polyps by direct examination, endoscopy, or sinus CT scan; AND
5. Member has symptoms of chronic rhinosinusitis after at least a 4-week trial with an intranasal corticosteroid (e.g., mometasone, fluticasone) in combination with nasal saline irrigation AND one of the following:
  - a) Prior sinonasal surgery;
  - b) Systemic corticosteroids (unless not tolerated or contraindicated); AND
6. Medication is used as an add-on maintenance treatment in combination with intranasal corticosteroid, unless not tolerated or contraindicated; AND
7. The member's weight (kg) and baseline serum IgE level (IU/mL) is documented in chart notes; AND
8. Member does not have allergic fungal rhinosinusitis (AFRS).
9. Dosage allowed: 75 mg to 600 mg subQ every 2 or 4 weeks based on serum total IgE level (IU/mL) measure before the start of treatment and by body weight (kg). See the dose determination chart in package insert.

If member meets all the requirements listed above, the medication will be approved for 6 months.

For reauthorization:

1. Member must be in compliance with all other initial criteria; AND
2. Medication is to be used as add-on maintenance therapy in combination with intranasal corticosteroids, unless not tolerated or contraindicated; AND
3. Chart notes have been provided showing improvement of signs and symptoms such as reduced post-nasal drip, reduced nasal polyp size, and/or reduced nasal congestion symptoms.

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

## CHRONIC IDIOPATHIC URTICARIA (CIU)

For initial authorization:

1. Member must be 12 years of age or older; AND
2. Medication must be prescribed by or under the recommendation of a dermatologist, allergist, or immunologist; AND
3. Member has a diagnosis of Chronic Idiopathic Urticaria that has been continuously or intermittently present for at least 6 weeks; AND
4. Member has trialed and failed at least one of the following for no less than 14 days:
  - a) A second generation H1 antihistamine (i.e. loratadine, cetirizine, fexofenadine) at 2-4 times the FDA-approved dosage;
  - b) Two second generation H1 antihistamines in combination;
  - c) A second generation H1 antihistamine plus a leukotriene receptor antagonist (i.e. montelukast, zafirlukast);
  - d) A second generation H1 antihistamine plus a first generation H1 antihistamine (i.e. diphenhydramine, hydroxyzine, chlorpheniramine);
  - e) A second generation H1 antihistamine plus an H2 antagonist (i.e. famotidine, cimetidine, ranitidine).
5. Dosage allowed: 150 or 300 mg by subcutaneous injection every 4 weeks.

If member meets all the requirements listed above, the medication will be approved for 16 weeks.

For reauthorization:

1. Member must be in compliance with all other initial criteria; AND
2. Chart notes have been provided to support a positive clinical response (i.e. reduction in itch severity and/or hive count).

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

## MODERATE TO SEVERE PERSISTENT ASTHMA

For initial authorization:

1. Member must be 6 years of age or older; AND
2. Medication must be prescribed by or on in consultation with a pulmonologist, immunologist or allergist; AND
3. Member has allergy testing performed, as indicated by:
  - a) Positive skin testing for perennial aeroallergen; AND/OR
  - b) Reactivity to at least one aeroallergen documented by elevated serum IgE level; AND
4. Member has a weight documented and a baseline plasma immunoglobulin E (IgE) level of 30 IU/mL or higher; AND
5. Member has at least two documented severe asthma exacerbations requiring oral corticosteroids (OCS), or at least one requiring hospitalization, within last year; AND
6. Member's asthma has been inadequately controlled after 3 months of conventional treatment of medium to high doses of inhaled corticosteroids (ICS) and long acting beta 2-agonists (LABA); AND
7. Medication is being used as add-on maintenance treatment to conventional therapies for asthma (i.e. ICS, LABA, etc.); AND
8. Medication is not used in conjunction with any other biologic therapy for asthma.
9. Dosage allowed: 75 to 375 mg by subcutaneous injection every 2 or 4 weeks.

If member meets all the requirements listed above, the medication will be approved for 16 weeks.

**For reauthorization:**

1. Medication is not being used as monotherapy for asthma; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has demonstrated improvement during 16 weeks of medication therapy:
  - a) Decreased frequency of emergency department visit or hospitalizations due to asthma exacerbations; OR
  - b) Improved functional ability (i.e. decreased effect of asthma on ability to exercise, function in school or at work, or quality of sleep); OR
  - c) Decreased utilization of rescue medications or oral corticosteroids.

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

CareSource considers Xolair (omalizumab) not medically necessary for the treatment of the diseases that are not listed in this document.

DATE	ACTION/DESCRIPTION
05/18/2017	New policy for Xolair created. For CIU urticaria activity score, trial of oral corticosteroids and trial length added.
01/12/2021	<b>Persistent Asthma:</b> added documented weight; added exacerbation requirement (two requiring OCS or one requiring hospitalization within the last year); FEV1 removed; ICS + LTRA removed; added not to be used with other asthma biologics. <b>CIU:</b> added immunologist; documented urticaria activity and itch severity scores removed; trial of oral corticosteroids removed; added trial option of a 2 <sup>nd</sup> generation H1 antihistamine 2-4x FDA approved dosage; added examples of trial drugs for reference. New indication <b>Nasal Polyps</b> added.

**References:**

1. Xolair [package insert]. South San Francisco, CA: GenetechUSA, Inc; December, 2020.
2. Buhl R. Omalizumab (Xolair) improves quality of life in adult patients with allergic asthma: A review. *Respir Med.* 2003;97(2):123-129.
3. Finn A, Gross G, van Bavel J, et al. Omalizumab improves asthma-related quality of life in patients with severe allergic asthma. *J Allergy Clin Immunol.* 2003;111(2):278-284.
4. Bang LM, Plosker GL. Omalizumab: A review of its use in the management of allergic asthma. *Treat Respir Med.* 2004;3(3):183-199.
5. Difficult-To-Treat & Severe Asthma in Adolescent and Adult Patients: Diagnosis and Management. Global Initiative For Asthma (GINA); Apr. 2019. Available at: <https://ginasthma.org/wp-content/uploads/2018/11/GINA-SA-FINAL-wms.pdf>.
6. 2020 Focused Updates To The Asthma Management Guidelines. National Institute of Health; Dec 2020. Available at: <https://www.nhlbi.nih.gov/health-topics/asthma-management-guidelines-2020-updates>.
7. Bernstein JA, Lang DM, Khan DA, et al. The diagnosis and management of acute and chronic urticaria: 2014 update. *J Allergy Clin Immunol.* 2014;133(5):1270-1277. doi:10.1016/j.jaci.2014.02.036
8. Zuberbier T, Aberer W, Asero R, et al. The EAACI/GA<sup>2</sup>LEN/EDF/WAO guideline for the definition, classification, diagnosis and management of urticaria. *Allergy.* 2018; 73(7): 1355-1566. doi.org/10.1111/all.13397.
9. Gevaert P, Omachi TA, Corren J, et al. Efficacy and safety of omalizumab in nasal polyposis: 2 randomized phase 3 trials. *J Allergy Clin Immunol.* 2020;146(3):595-605.
10. Stevens WW, Schleimer RP, Kern RC. Chronic Rhinosinusitis with Nasal Polyps. *J Allergy Clin Immunol Pract.* 2016;4(4):565-572. doi:10.1016/j.jaip.2016.04.012.
11. Peters AT, Spector S, Hsu J, et al. Diagnosis and management of rhinosinusitis: a practice parameter update. *Ann Allergy Asthma Immunol.* 2014;113(4):347-85. DOI: 10.1016/j.anai.2014.07.025.
12. Sedaghat AR. Chronic rhinosinusitis. *Am Fam Physician.* 2017;96(8):500-506.
13. Laidlaw TM, Buchheit KM. Biologics in chronic rhinosinusitis with nasal polyposis. *Ann Allergy Asthma Immunol.* 2020;124(4):326-332. doi:10.1016/j.anai.2019.12.001.



14. Orlandi RR, Kingdom TT, Hwang PH, et al. International consensus statement of allergy and rhinology: rhinosinusitis. *Int Forum Allergy Rhinol.* 2016 Feb;6 Suppl 1:S22-209. doi: 10.1002/alr.21695.
15. Fokkens WJ, Lund V, Bachert C, et al. EUFOREA consensus on biologics for CRSwNP with or without asthma. *Allergy.* 2019 Dec;74(12):2312-2319. doi: 10.1111/all.13875.

Effective date: 07/01/2021

Revised date: 01/12/2021