



## MEDICAL POLICY STATEMENT INDIANA MEDICAID

Policy Name		Policy Number	Date Effective
Drug Testing		MM-0126	01/01/2022-09/30/2022
Policy Type			
MEDICAL	Administrative	Pharmacy	Reimbursement


Medical Policy Statement prepared by CareSource and its affiliates are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other policies and procedures.

Medical Policy Statements prepared by CareSource and its affiliates do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage) for the service(s) referenced in the Medical Policy Statement. If there is a conflict between the Medical Policy Statement and the plan contract (i.e., Evidence of Coverage), then the plan contract (i.e., Evidence of Coverage) will be the controlling document used to make the determination.

According to the rules of Mental Health Parity Addiction Equity Act (MHPAEA), coverage for the diagnosis and treatment of a behavioral health disorder will not be subject to any limitations that are less favorable than the limitations that apply to medical conditions as covered under this policy.

### Table of Contents

A. Subject.....	2
B. Background.....	2
C. Definitions.....	2
D. Policy.....	3
E. Conditions of Coverage.....	8
F. Related Policies/Rules.....	8
G. Review/Revision History.....	8
H. References.....	9



## A. Subject Drug Testing

### B. Background

Drug testing is a part of medical care during the initial assessment, ongoing monitoring, and recovery phase for members with substance use disorder (SUD); for members who are at risk for abuse/misuse or diversion of drugs; and/or for other medical conditions. The drug test guides a provider in diagnosing and planning the member's care when prescription medications or illegal drugs are of concern.

Drug testing is one component of a comprehensive clinical approach during the initial assessment, stabilization, maintenance and recovery phase for members with a substance use disorder (SUD). It is also used to screen members periodically that are prescribed chronic opioid therapy (COT) for pain based on a risk score.

For substance-related disorders, drug testing may help the provider compare a member's reported drug(s) of choice with the test results to verify subjective information. The assessment process including initial drug testing will aid the treatment provider to individualize the plan for drug testing for a member.

Drug testing may help determine if a member is adhering to prescription medication, reveal nonprescribed drugs or illicit drugs, or provide evidence to suggest diversion. Providers requesting drug testing should have proficiency in drug test interpretation and understand what they are ordering.

Urine is the most common specimen to monitor drug use. There are two main types of urine drug testing (UDT): presumptive and confirmatory. Drug testing is sometimes also referred to as toxicology testing.

### C. Definitions

- **Presumptive/Qualitative test** - The testing of a substance or mixture to determine its chemical constituents, also known as qualitative testing.
- **Confirmatory/Quantitative test** - A test that determines the amount of a substance per unit volume or unit weight, also known as quantitative or definitive testing.
- **Random drug test** - A laboratory drug test administered at an irregular interval that is not known in advance by the member.
- **Relapse** - When a person with addiction returns to use after a period of sobriety.
- **Early and Periodic Screening, Diagnostic and Treatment (EPSDT)** - This benefit provides comprehensive and preventive health care services for children under age 21 who are enrolled in Medicaid.
- **Aberrant behavior** - Member's behaviors that may indicate medication/drug abuse or misuse such as losing prescriptions, early refill requests, or multiple prescribers for controlled substances on the Indiana Prescription Drug Monitoring Program
- **Independent laboratory** - A laboratory certified to perform diagnostic and/or clinical tests independent of an institution or a provider's office.



- **Participating/non-participating** - Participating means in-network and contracted with CareSource. "Non-participating," means out-of-network, not contracted with CareSource.
- **Opioid treatment program (OTP)** - Program or qualified provider delivering opioid treatment to members with an opioid agonist treatment medication.
- **Residential services** - Includes individual and group therapy, medication training and support, case management, drug testing, and peer recovery supports.
- **Clinical Laboratory Improvement Amendments (CLIA)** - The Centers for Medicare & Medicaid Services (CMS) regulates programs that test human specimens to ensure accurate, reliable and timely patient test results regardless of where a test is performed and includes physician offices.
- **Chronic opioid therapy** - Refers to the use of opioids to treat chronic pain more than three months or past the time of normal tissue healing.
- **Diversion** - Unlawful channeling to regulated pharmaceuticals from legal sources to the illicit marketplace and includes transferring drugs to people they were not prescribed for.

#### D. Policy

##### I. Prior Authorization (PA)


- A. Prior authorization (PA) is required for all definitive drug panels with 15 or more drug classes (HCPCS code G0482 & G0483). PA will also be required for definitive testing performed beyond 16 cumulative units, per member, per calendar year, and for presumptive testing performed for 53 or more cumulative units, per member, per calendar year.
- B. PA is not required for the first 52 units of presumptive UDT, nor for the first 16 units of definitive UDT of under 15 drug classes (HCPCS code G0480, G0481).

##### II. UDT order

- A. An order for UDT must include, at a minimum, all of the following:
  - 1. List the type of test to be performed (presumptive or confirmatory).
  - 2. Include all medications currently prescribed to the member.
  - 3. Drug and drug class to be tested.
  - 4. Clinical indication.
  - 5. Be signed and dated by a qualified provider.
  - 6. UDT order must specifically match the number, level and complexity of the testing components performed.
- B. Copies of test results alone without the proper provider's order for the test are not sufficient documentation of medical necessity to support a claim.

##### III. Provider Documentation

- A. Provider must maintain a complete legible medical record for the member and include the following:
  - 1. Complete member name and identification on each page of record.
  - 2. Identification of the provider responsible for providing member care.
  - 3. Appropriate indication for UDT.
  - 4. How the UDT result will guide the plan of treatment.
  - 5. CPT code that accurately describes the service performed.




- B. Provider documentation must support medical necessity of UDT.
  - 1. All components of a UDT panel must be supported by medical necessity.
  - 2. A panel of drugs may be performed as part of an initial assessment to develop a monitoring plan as long as it is supported with medical necessity. A panel of drugs should only be conducted based on an individualized treatment plan noting the need for confirmatory test with greater than 15 drug tests. These tests are rarely indicated for routine UDT.
  - 3. Provider documentation needs to make a clear case for medical necessity for the level of testing being requested, it may include but is not limited to:
    - a. Phase of treatment (e.g. assessment, early recovery, induction, stabilization, maintenance).
    - b. Current level of care (e.g. use of ASAM levels).
    - c. Member drug(s) of choice.
    - d. Days since last drug test with unexpected results.
    - e. Current prescribed drugs including over-the-counter drugs and illicit drugs that have had unexpected results in recent tests.
    - f. Member current active symptoms that led to the request.
    - g. Provider actions taken on recent unexpected test results and member response to that action.
    - h. The clinical documentation shows that the member is contesting the result of an unexpected presumptive test.
    - i. The test is not being requested for third party reasons, or as a condition to stay in sober housing or residential facility (see additional information below).
    - j. Results of any pill counts performed by treatment team.
- C. Must submit appropriate clinical documentation with PA request to determine appropriate medical necessity.

IV. Presumptive drug testing:

- A. May be reported with CPT codes 80305- 80307.
- B. These codes differ based on the level of complexity of the testing methodology.
- C. Only one code from this code range may be reported per date of service.
- D. Providers performing validity testing on urine specimens utilized for drug testing shall not separately bill for validity testing of the specimen.
  - 1. For example, if a laboratory performs a urinary pH, specific gravity, creatinine, nitrates, oxidants, or other tests to confirm that a urine specimen is not adulterated, this testing is not separately billed.

V. Per recommendations from DMHA, IHCP providers are encouraged to use presumptive drug testing methods as these are clinically appropriate for detecting nearly all prescription opioids, benzodiazepines and illicit drugs.

- A. The use of definitive testing should instead be based on the need to detect specific opioids that cannot be identified on presumptive UDTs, or in the presence of unexpected UDT results.
  - 1. For example, rapid testing may be problematic for the accurate detection of amphetamines, and therefore confirmatory testing may be necessary for unanticipated results.



- B. Providers may also use confirmatory testing to assess for drug metabolites which may help identify if the member has been consistently taking prescribed medications as intended.
  - C. Providers should not test for substances for which results would not affect patient management.
  - D. The provider that is billing for UDT has the responsibility of ensuring that services were billed in accordance with these requirements.
- VI. UDT continues to not be covered in any of the following circumstances:
- A. Unnecessarily frequent drug testing without consideration for a specific drug's window of detection
  - B. Testing for the same drug with both a blood or saliva test and a urine specimen simultaneously (multiple tests seeking the same outcome)
  - C. Testing for legal intervention or employment
- VII. Laboratory
- A. Drug testing conducted for CareSource members by non-participating labs or facilities is not billable to and will not be reimbursed by CareSource, even if such tests were ordered by a participating provider.
  - B. CareSource may require documentation of FDA-approved complexity level for instrumented equipment, and/or Clinical Laboratory Improvement Amendments (CLIA) Certificate of Registration, Compliance, or Accreditation as a high complexity lab.
  - C. Laboratories must maintain hard copy documentation of lab results with copies of the order for the drug test and any required PA.

VIII. Clinical Indications

Testing should be individualized to the specific member. Analytes tested should be ordered based on the member's drug(s) of choice. Periodically, drugs that are commonly used may be rotated into the random test schedule. Regionally prevalent drugs may be periodically rotated into the random test schedule. The rationale for which tests is not meant to include all drugs all of the time, rather the drugs most likely to be seen in the individual. This information helps the provider focus the testing to likelihood it would be used. Testing should be at the lowest level to inform the provider that an intervention is needed based on the individual history of the member.

Drug testing is ideally performed on a random unannounced schedule with a specific time frame to produce a specimen. ASAM recommends a random-interval schedule to a fixed-interval schedule as it eliminates known non-testing periods. Testing every day, at every visits, on the same day of the week or at the same time is not random (ASAM reference). Providers should understand windows of detection time to determine frequency of testing. Too short of an interval may raise an issue of presumption of renewed when the same drug that was recently tested is still within its detection window. Drug testing does not have to be associated with an office visit when patients are called to do random tests.



Providers should be aware of the potential for cross-reactivity when using presumptive tests. Per SAMHSA, cross reactivity has a positive side. For example a confirmatory test for a specific opioid analyte will miss other opioids a member may be taking. Therefore an opioid screen is preferred over a specific test when looking for opiate type drugs.

When testing for alcohol, SAMHSA also states that a breathalyzer gives an estimate of blood alcohol level. This method is simple to use, inexpensive, gives instant results, and is noninvasive.

A. Drug Testing in Addiction Treatment

1. UDT frequency is expected to be more frequent when medically necessary early in treatment or when tapering. UDT frequency is expected to decrease as member stabilizes. Prior to Initiation or in the Induction Phase (early recovery).
  - a. Obtain history as well as a medical and psychological assessment.
  - b. Review approximate time frame of drug detected in urine.
  - c. Identify questions seeking to answer as well as treatment plan based on potential UDT results.
  - d. Obtain an individualized baseline UDT based on member's unique clinical presentation, prescribed medications, member's self-reported drugs of choice, and regional drug trends.
  - e. At least weekly (cite ASMA consensus guidelines).
  - f. Discuss results with member.
  - g. Agree on plan of care that includes treatment interventions and goals.
  - h. This phase includes members that have relapsed.
2. Maintenance phase.
  - a. At least once per month.
3. Intensive outpatient.
  - a. At least weekly.
4. Substance use disorder residential program.
  - a. At least monthly.
5. Stable recovery.
  - a. Drug testing may be done less frequently if in stable recovery.
6. For members taking long-action naltrexone.
  - a. At least monthly.

B. Drug testing in an Opioid Treatment Program (OTP)

1. In maintenance treatment, federal regulations governing OTP require initial toxicology plus 8 random UDT screens per year per member.
2. In short-term detoxification treatment, one initial UDT per member.
3. In long-term detoxification treatment initial and monthly random UDT per member.

C. Chronic Pain Management

1. Prior to or when initiating treatment
  - a. Complete an assessment for risk of substance abuse using a validated risk assessment screening tool such as Screener and Opioid Assessment for Patient with Pain-Revisited (SOAPP-R) or the Opioid Risk Tool (ORT).





- b. Review the state prescription drug monitoring program data (PDMP).
  - c. Obtain baseline UDT screening.
  - d. Discuss results with member.
  - e. Agree on plan of care that includes treatment goals, educating on risks and benefits, and strategies to mitigate risks.
  - f. Combine evidence-based non-pharmacologic and non-opioid pharmacologic therapy as necessary.
2. Ongoing monitoring of treatment is determined by level of risk for substance use:
- a. Low risk – UDT once a year.
  - b. Moderate risk – UDT twice a year.
  - c. High risk – UDT up to 4 times a year.
  - d. UDT when member shows aberrant drug-related behavior.
  - e. Review PDMP data every 1-3 months.
  - f. Evaluate benefits and harms of treatment at least every 3 months.

IX. Blood Testing

- A. Blood drug testing is considered medically necessary when it is in the emergency room setting.

X. Testing that is not medically necessary

- A. CareSource considers the following as not medically necessary for either presumptive or confirmatory testing:

1. Testing that is not individualized such as:
  - a. Reflexive testing.
  - b. Routine orders.
  - c. Standard orders.
  - d. Preprinted orders.
  - e. Requesting all tests that a machine is capable of doing solely because a result may be positive.
  - f. Large, arbitrary panels.
  - g. Universal testing.
  - h. Conduct additional testing as needed.

- B. Testing required by third parties such as:

1. Testing ordered by a court or other medico-legal purpose such as child custody.
2. Testing for pre-employment or random testing that is a requirement of employment.
3. Physician's health programs (recovery for physicians, dentists, veterinarians, pharmacists, etc.).
4. School entry or testing for athletics.
5. Testing required for military service.
6. Testing in residential facility, partial hospital, or sober living as a condition to remain in that community.
7. Testing with another pay source that is primary such as a county, state or federal agency.
8. Testing for marriage license.



9. Forensic.
  10. Testing for other admin purposes.
  11. Routine physical/medical examination.
- C. Testing for validity of specimen
1. It is included in the payment for the test and will not be reimbursed separately.
- D. Blood drug testing when completed outside of the emergency room.
- E. Hair, saliva, or other body fluid testing for controlled substance monitoring.  
Testing for the same drug with both a blood or saliva test and a urine specimen simultaneously (multiple tests seeking the same outcome).
- F. Any type of drug testing not addressed in this policy.
- G. Routine nonspecific or wholesale orders including routine drug panels.
- H. Routine use of confirmatory testing following a negative presumptive expected result.
- I. Custom Profiles, standing orders, drug screen panel, custom panel, blanket orders, reflex testing or conduct additional testing as needed orders.
- J. A confirmatory test prior to discussing results of presumptive test with member.
- K. Testing paid for by another primary source (e.g. county, state or federal agency).

**NOTE:** Compliance with the provisions in this policy may be monitored and addressed through post payment data analysis, subsequent medical review audits, recovery of overpayments identified, and provider prepay review.

E. Conditions of Coverage

NA


F. Related Policies/Rules

NA

G. Review/Revision History

DATE		ACTION
<b>Date Issued</b>	12/31/2017	
<b>Date Revised</b>	11/01/2019	Updated clinical indications, quantity limits, and prior authorization requirements
	01/15/2020	Updated quantity limits and PA requirements
	09/02/2020	Updated V.
	09/29/2021	Updated quantity limits and prior authorization requirements
<b>Date Effective</b>	01/01/2022	Updated quantity limits. Updated reference dates. Approved at PGC.
<b>Date Archived</b>	09/30/2022	This Policy is no longer active and has been archived. Please note that there could be other Policies that may have some of the same rules incorporated and CareSource reserves the right to follow CMS/State/NCCI guidelines without a formal documented Policy.





## H. References

1. A. Jaffe, S. Molnar, N. Williams, E. Wong, T. Todd, C. Caputo, J. Tolentino and S. Ye. (2016, January). Review and recommendations for drug testing in substance use treatment contexts. *Journal of Reward Deficiency Syndrome and Addiction Science*. 2(1): 28-45. doi: 10.17756/jrdsas.2016-025
2. A. Rzetelny, B. Zeller, N. Miller, K. E. City, K. L. Kirsh and S. D. Passik. (2016) Counselors' clinical use of definitive drug testing results in their work with substance-use patients: A qualitative study. *International Journal of Mental Health and Addiction*. (14) 64-80. doi: 10.1007/s11469-015-9569-7
3. American Society of Addiction Medicine. (2017, May/June). Appropriate use of drug testing in clinical addiction medicine. 11(3) 163-173. doi: 10.1097/ADM.0000000000000323
4. Andersson, H. W., Wenaas, M., & Nordfjærn, T. (2019). Relapse after inpatient substance use treatment: A prospective cohort study among users of illicit substances. *Addictive Behaviors*, (90)222-228. doi:10.1016/j.addbeh.2018.11.008
5. American Society of Addiction Medicine (2010, October). Public Policy Statement on Drug Testing as a Component of Addiction Treatment and Monitoring Programs and in other Clinical Settings. Retrieved August 9, 2021 from [www.asam.org](http://www.asam.org)
6. Dowell, D., Haegerich, T. M., & Chou, R. (2016, March). CDC Guideline for Prescribing Opioids for Chronic Pain – United States, 2016. Retrieved August 9, 2021 from [www.cdc.gov](http://www.cdc.gov)
7. eCFR-Code of Federal Regulations. 42 Code of Federal Regulations (CFR) Part 8. (n.d.). Retrieved August 9, 2021 from [www.ecfr.gov](http://www.ecfr.gov)
8. Gourlay, D. L., Heit, H. H., & Caplan, Y. H. (2015, August 31). Urine Drug Testing in Clinical Practice The Art and Science of Patient Care (Edition 6). PharmaCom Group Inc./Center for Independent Healthcare Education
9. Jarvis, M, Williams, J, Hurford, M, Lindsay, D, Lincoln, P, Giles, L, Luongo, P, Safarian, T. (2017, April 5). Appropriate Use of Drug Testing in Clinical Addiction Medication. *Journal of Addiction Medicine*. Retrieved August 9, 2021 from [www.dca.ca.gov](http://www.dca.ca.gov)
10. Medicare Learning Network. (2020, May). CLIA Program and Medicare Laboratory Services. Retrieved August 12, 2020 from [www.cms.gov](http://www.cms.gov)
11. National Academies of Sciences, Engineering, and Medicine. 2017. Pain management and the opioid epidemic: Balancing societal and individual benefits and risks of prescription opioid use. Washington, DC: The National Academies Press. doi: <https://doi.org/10.17226/24781>.
12. Owen, G, Burton, A, Schade, C, Passik, S. (2012) Urine Drug Testing: Current Recommendations and Best Practices. *Pain Physician Journal*. 15, ES119-ES133. Retrieved August 9, 2021 from [www.painphysicianjournal](http://www.painphysicianjournal)
13. Reisfield, MD, G. M., Webb, PhD, F. J., Bertholf, PhD, R. L., Sloan, MD, P. A., & Wilson, MD, G. R. (2007). Family physicians' proficiency in urine drug test interpretation. *Journal of Opioid Management*, 3(6), 333. doi:10.5055/jom.2007.0022
14. Substance Abuse and Mental Health Services Administration. Clinical Drug Testing in Primary Care. Technical Assistance Publication (TAP) 32. HHS Publication No. (SMA) 12-4668. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2012.



15. Stanos, S. P. (2017, October 10). President's Message. National Academies of Sciences, Engineering, and Medicine (NASEM). Pain Medicine, 18(10), 1835-1836. doi:10.1093/pm/pnx224
16. U.S. Department of Veterans Affairs. (2017, February). VA/DoD Clinical Practice Guideline for Opioid Therapy for Chronic Pain. Retrieved August 9, 2021 from [www.va.gov](http://www.va.gov)
17. Agency Medical Directors Group. (2010). Interagency Guideline on Opioid Dosing for Chronic Non-cancer Pain. Retrieved August 9, 2021 from [www.agencymeddirectors.wa.gov](http://www.agencymeddirectors.wa.gov)
18. Indiana Health Coverage Programs Provider Reference Module. (2019, November 21). Laboratory Services. Retrieved on August 9, 2021 from [www.in.gov](http://www.in.gov)
19. Indiana Health Coverage Programs. (2018, August 30). IHCP Bulletin BT201846 Retrieved on August 9, 2021 from [www.provider.indianamedicaid.com](http://www.provider.indianamedicaid.com)

**The Medical Policy Statement detailed above has received due consideration as defined in the Medical Policy Statement Policy and is approved.**

IN-MED-P-931858

Issue Date 12/31/2017

Approved OMPP 10/25/2021