

## PHARMACY POLICY STATEMENT

### Indiana Medicaid

DRUG NAME	Skyrizi (risankizumab-rzaa)
BILLING CODE	Must use valid NDC code
BENEFIT TYPE	Pharmacy
SITE OF SERVICE ALLOWED	Home/Office
COVERAGE REQUIREMENTS	Prior Authorization Required (Non-Preferred Product) Alternative preferred products include Cimzia, Cosentyx, Enbrel, Otezla and Siliq QUANTITY LIMIT— 2 syringes every 12 weeks after loading doses
LIST OF DIAGNOSES CONSIDERED <b>NOT</b> MEDICALLY NECESSARY	<a href="#">Click Here</a>

Skyrizi (risankizumab-rzaa) is a **non-preferred** product and will only be considered for coverage under the **pharmacy** benefit when the following criteria are met:

Members must be clinically diagnosed with one of the following disease states and meet their individual criteria as stated.

### PLAQUE PSORIASIS (PsO)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Medication must be prescribed by or in consultation with a dermatologist; AND
3. Member has clinical documentation of moderate to severe plaque psoriasis characterized by 3% or more of body surface area (BSA) or disease affecting sensitive areas (e.g., hands, feet, face, genitals, etc.); AND
4. Member must have a documented negative TB test (i.e., tuberculosis skin test (PPD), interferon-gamma release assay (IGRA)) within 12 months prior to starting therapy; AND
5. Member has tried and failed to respond to treatment with at least **one** of the following:
  - a) At least 12 weeks of photochemotherapy (i.e., psoralen plus ultraviolet A therapy);
  - b) At least 12 weeks of phototherapy (i.e., UVB light therapy, Excimer laser treatments);
  - c) At least a 4 week trial with topical antipsoriatic agents (i.e., anthralin, calcipotriene, coal tar, corticosteroids, tazarotene, tacrolimus, pimecrolimus); AND
6. Member has tried and failed, or unable to tolerate a systemic non-biologic DMARD (i.e., cyclosporine, methotrexate, acitretin) for at least 12 weeks; AND
7. Member has tried and failed treatment with at least **two** of the following: Cimzia, Cosentyx, Enbrel, Otezla and Siliq. Treatment failure requires at least 12 weeks of therapy with each drug.
8. **Dosage allowed:** 150 mg (two 75 mg injections) administered by subcutaneous injection at week 0, week 4, and every 12 weeks thereafter.

***If member meets all the requirements listed above, the medication will be approved for 12 months.***

For **reauthorization**:

1. Member must be in compliance with all other initial criteria; AND
2. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease (e.g., documented member's BSA improvement, etc.).

***If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.***

**CareSource considers Skyrizi (risankizumab-rzaa) not medically necessary for the treatment of the diseases that are not listed in this document.**

DATE	ACTION/DESCRIPTION
07/28/2019	New policy for Skyrizi created.
11/18/2020	Removed rheumatologist from prescriber requirement. Removed PsO 6 months or longer. Changed BSA to 3% or sensitive areas. Removed PASI score. Removed repeat TB for reauth. Updated references.

References:

1. Skyrizi [prescribing information]. North Chicago, IL: AbbVie Inc.; March 2020.
2. Elmets CA, Korman NJ, Prater EF, et al. Joint AAD-NPF Guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures [published online ahead of print, 2020 Jul 30]. *J Am Acad Dermatol*. 2020;S0190-9622(20)32288-X.
3. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol*. 2020;82(6):1445-1486.
4. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019;80(4):1029-1072.
5. Elmets CA, Lim HW, Stoff B, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management and treatment of psoriasis with phototherapy [published correction appears in *J Am Acad Dermatol*. 2020 Mar;82(3):780]. *J Am Acad Dermatol*. 2019;81(3):775-804.
6. Menter A, Cordoro KM, Davis DM, et al. Joint AAD-NPF Guidelines of care for the management and treatment of psoriasis in pediatric patients. *J Am Acad Dermatol* 2020;82:161-201.
7. ClinicalTrials.gov. Identifier: NCT02684370. BI 655066 (Risankizumab) Compared to Placebo and Active Comparator (Ustekinumab) in Patients With Moderate to Severe Chronic Plaque Psoriasis. Available at: <https://clinicaltrials.gov/ct2/show/NCT02684370?term=ULTIMMA-1&rank=1>.
8. ClinicalTrials.gov. Identifier: NCT02684357. BI 655066 Versus Placebo & Active Comparator (Ustekinumab) in Patients With Moderate to Severe Chronic Plaque Psoriasis. Available at: <https://clinicaltrials.gov/ct2/show/NCT02684357?term=ULTIMMA-2&rank=1>.

Effective date: 04/01/2021

Revised date: 11/18/2020