

PHARMACY POLICY STATEMENT Kentucky Medicaid	
DRUG NAME	Remicade (infliximab)
BILLING CODE	J1745 (1 unit = 10 mg or 1 x 100 mg vial = 10 units)
BENEFIT TYPE	Medical
SITE OF SERVICE ALLOWED	Office/Outpatient
COVERAGE REQUIREMENTS	Prior Authorization Required (Non-Preferred Product) Alternative preferred products include Actemra, Cimzia, Cosentyx, Enbrel, Kevzara, Olumiant, Otezla, Siliq and Xeljanz QUANTITY LIMIT—1200 mg (120 units per dose)
LIST OF DIAGNOSES CONSIDERED NOT MEDICALLY NECESSARY	Click Here

Remicade (infliximab) is a **non-preferred** product and will only be considered for coverage under the **medical** benefit when the following criteria are met:

Members must be clinically diagnosed with one of the following disease states and meet their individual criteria as stated.

ANKYLOSING SPONDYLITIS (AS)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Must have a documented negative TB test (i.e., tuberculosis skin test (PPD), an interferon-release assay (IGRA)) within 12 months prior to starting therapy; AND
3. Medication must be prescribed by a rheumatologist; AND
4. Member has had back pain for 3 months or more that began before the age of 50; AND
5. Current imaging results show an inflammation of one or both of the sacroiliac joints; AND
6. Member shows at least **one** of the following signs or symptoms of Spondyloarthritis:
 - a) Arthritis;
 - b) Elevated serum C-reactive protein;
 - c) Inflammation at the tendon, ligament or joint capsule insertions;
 - d) Positive HLA-B27 test;
 - e) Limited chest expansion;
 - f) Morning stiffness for 1 hour or more; AND
7. Member meets at least **one** of the following scenarios:
 - a) Member has Axial (spinal) disease;
 - b) Member has peripheral arthritis without axial involvement and has tried and failed treatment with methotrexate or sulfasalazine. Treatment failure requires at least 30 days of therapy without an adequate response;
 - c) Member has tried and failed to respond to treatment with at least **two** prescription NSAIDs taken at the maximum recommended dosages. Treatment failure requires at least 30 days of therapy with each NSAID without an adequate response; AND

8. Member must have tried and failed treatment with at least **two** of the following: Enbrel, Cimzia and Cosentyx. Treatment failure requires at least for 30 days of therapy with each drug.
9. **Dosage allowed:** 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks. Prior to any dosages or dosing frequencies greater than listed, medical necessity documentation must be supplied to justify coverage.

If member meets all the requirements listed above, the medication will be approved for 12 months.

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease.

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

CROHN'S DISEASE (CD)

For **initial** authorization:

1. Member is 6-17 years of age with moderately to severely active CD as defined by Pediatric Crohn's Disease Activity Index (PCDAI) greater than 30 OR member is 18 years of age or older with moderately to severely active non-fistulizing CD as defined by Crohn's Disease Activity Index (CDAI) greater than 220 and less than 400; AND
2. Member has had a trial and inadequate response to at least 30 days of **one** of the following:
 - a) 6-mercaptopurine;
 - b) Azathioprine;
 - c) Methotrexate;
 - d) Corticosteroid(s); OR
3. Member is 18 years of age or older with fistulizing CD; AND
4. Must have a documented negative TB test (i.e., tuberculosis skin test (PPD), an interferon-release assay (IGRA)) within 12 months prior to starting therapy; AND
5. Medication must be prescribed by a gastroenterologist.
6. **Dosage allowed:** 5 mg/kg at 0, 2, and 6 weeks, followed by 5 mg/kg every 8 weeks thereafter. If no response by week 14, consider discontinuing therapy.

If member meets all the requirements listed above, the medication will be approved for 12 months.

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease; AND
4. Documented member's PCDAI or CDAI score improvement.

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

PLAQUE PSORIASIS (PsO)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Must have a documented negative TB test (i.e., tuberculosis skin test (PPD), an interferon-release assay (IGRA)) within 12 months prior to starting therapy; AND
3. Medication must be prescribed by a dermatologist or rheumatologist; AND
4. Member has PsO for 6 months or longer; AND
5. Member is not going to receive systemic therapy or phototherapy while on Remicade; AND
6. Member's plaque psoriasis involving 10% or more of the body surface area (BSA) or 5% or more of BSA if psoriasis involves sensitive areas (hands, feet, face, or genitals); AND
7. Member's Psoriasis Area and Severity Index (PASI) greater than or equal to 12; AND
8. Member has tried and failed to respond to treatment with at least **one** of the following:
 - a) At least 30 days of photochemotherapy (i.e., psoralen plus ultraviolet A therapy);
 - b) At least 30 days of phototherapy (i.e., UVB light therapy, Excimer laser treatments (tanning beds emit mostly UVA light and therefore would not meet this criteria)).
 - c) At least a 30-day trial with topical antipsoriatic agents (i.e., anthralin, calcipotriene, coal tar, corticosteroids, tazarotene); AND
9. Member has tried and failed to respond to treatment with traditional first-line oral/systemic therapies (i.e., cyclosporine, methotrexate, acitretin) for at least 30 days; AND
10. Member has tried and failed treatment with at least **two** of the following: Cimzia, Cosentyx, Enbrel, Otezla and Siliq. Treatment failure requires at least for 30 days of therapy with each drug.
11. **Dosage allowed:** 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks. Prior to any dosages or dosing frequencies greater than listed, medical necessity documentation must be supplied to justify coverage.

If member meets all the requirements listed above, the medication will be approved for 12 months.

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease (e.g., documented member's PASI score improvement, etc.).

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

PSORIATIC ARTHRITIS (PsA)

For **initial** authorization:

1. Member must be 18 years of age or older; AND
2. Must have a documented negative TB test (i.e., tuberculosis skin test (PPD), an interferon-release assay (IGRA)) within 12 months prior to starting therapy; AND
3. Medication must be prescribed by a rheumatologist or dermatologist; AND
4. Member meets at least **one** of the following scenarios:
 - a) Member has predominantly axial disease (i.e., sacroiliitis or spondylitis) as indicated by radiographic evidence;
 - b) Member has shown symptoms of predominantly axial disease (i.e., sacroiliitis or spondylitis) for more than 3 months (i.e., limited spinal range of motion, spinal morning stiffness for more than 30 minutes) and has tried and failed to respond to treatment with at least 2 prescription NSAIDs taken

at the maximum recommended dosages. Treatment failure requires at least 30 days of therapy with each NSAID without an adequate response;

- c) Member has predominately non-axial disease (e.g., peripheral synovitis or dactylitis or nail involvement) and has tried and failed to respond to treatment with at least 30-day trial of methotrexate and NSAID taken at the maximum recommended dosages (if unable to tolerate or has contraindication to methotrexate than 30-day trial of sulfasalazine or azathioprine or cyclosporine); AND
5. Member must have tried and failed treatment with at least **two** of the following: Enbrel, Cimzia, Cosentyx, Otezla and Xeljanz. Treatment failure requires at least for 30 days of therapy with each drug.
6. **Dosage allowed:** 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks. Prior to any dosages or dosing frequencies greater than listed, medical necessity documentation must be supplied to justify coverage.

If member meets all the requirements listed above, the medication will be approved for 12 months.

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease.

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

RHEUMATOID ARTHRITIS (RA)

For **initial** authorization:

1. Member must be 18 years of age or older with moderate to severe active RA; AND
2. Must have a documented negative TB test (i.e., tuberculosis skin test (PPD), an interferon-release assay (IGRA)) within 12 months prior to starting therapy; AND
3. Medication must be prescribed by a rheumatologist; AND
4. Medication must be used in combination with methotrexate, or if intolerant to methotrexate, another immunosuppressant (i.e., azathioprine, hydroxychloroquine, cyclosporine, etc.); AND
5. Member must have tried and failed treatment with at least **two** non-biologic DMARDS OR must have a contraindication to all non-biologic DMARDS. Treatment trial duration with each non-biologic DMARD agent must have been at least 30 days (non-biologic DMARDS include: methotrexate, hydroxychloroquine, sulfasalazine, azathioprine, cyclosporine and leflunomide); AND
6. Member has tried and failed treatment with at least **two** of the following: Actemra, Cimzia, Enbrel, Kevzara, Olumiant and Xeljanz. Treatment failure requires at least for 30 days of therapy with each drug.
7. **Dosage allowed:** 3 mg/kg at 0, 2 and 6 weeks, then every 8 weeks. Prior to any dosages or dosing frequencies greater than listed, medical necessity documentation must be supplied to justify coverage.

If member meets all the requirements listed above, the medication will be approved for 12 months.

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease.

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

ULCERATIVE COLITIS (UC)

For **initial** authorization:

1. Member is 6-17 years of age with moderate to severe active UC as defined by Pediatric Ulcerative Colitis Activity Index (PUCAI) of 35 or greater OR member is 18 years of age or older with moderately to severely active UC as defined by Mayo score of 6 or greater with an endoscopy subscore of 2 or 3; AND
2. Medication must be prescribed by a gastroenterologist; AND
3. Must have a documented negative TB test (i.e., tuberculosis skin test (PPD), an interferon-release assay (IGRA)) within 12 months prior to starting therapy; AND
4. Member must have tried and failed with at least 30-day treatment with **one** or more of the following:
 - a) 6-mercaptopurine;
 - b) Azathioprine;
 - c) Methotrexate;
 - d) Oral corticosteroids;
 - e) Salicylates.
5. **Dosage allowed:** 5 mg/kg at 0, 2, and 6 weeks, followed by 5 mg/kg every 8 weeks thereafter.

If member meets all the requirements listed above, the medication will be approved for 12 months.

For **reauthorization**:

1. Must have been retested for TB with a negative result within the past 12 months; AND
2. Member must be in compliance with all other initial criteria; AND
3. Chart notes have been provided that show the member has shown improvement of signs and symptoms of disease.

If member meets all the reauthorization requirements above, the medication will be approved for an additional 12 months.

CareSource considers Remicade (infliximab) not medically necessary for the treatment of the following disease states based on a lack of robust clinical controlled trials showing superior efficacy compared to currently available treatments:

- Amyloid angiopathy
- Asthma
- Behcet's disease
- Birdshot retinochoroidopathy
- Bronchiolitis obliterans
- Central nervous system amyloidosis

- Chemotherapy induced enterocolitis (not due to Yervoy or Opdivo)
- Chronic immune-mediated myelitis
- Chronic obstructive pulmonary disease
- Cogan's syndrome
- Corneal ulcer
- Cranial nerve palsy
- Cystoid macular degeneration
- Disc herniation-induced sciatica
- Discoid lupus erythematosus
- Eczema
- Eosinophilic fasciitis
- Graft-versus-host-disease
- Granuloma annulare
- Granulomatous angiitis
- Granulomatous mastitis
- Hepatitis C genotype 1
- IgG4-related disease
- Infectious uveitis
- Iritis
- Juvenile idiopathic arthritis
- Kawasaki syndrome
- Localized scleroderma/morphea
- Membranous glomerulopathy
- Microscopic colitis
- Multifocal osteomyelitis (e.g., (chronic recurrent multifocal osteomyelitis (CRMO))
- Neurosarcoidosis
- Nodular scleritis
- Panniculitis
- Polyarteritis nodosa
- Polymyositis
- Prevention of post-operative recurrence of Crohn's disease
- Rejection following small bowel transplantation
- Relapsing polychondritis
- Scleroderma
- Sjogren's syndrome
- Still's disease
- Systemic lupus erythematosus
- Takayasu arteritis
- Tolosa-Hunt syndrome
- Tubulo-interstitial nephritis with uveitis (TINU) syndrome
- Wegener's granulomatosis/Wegener's peripheral neuropathy.

DATE	ACTION/DESCRIPTION
05/16/2017	New policy for Remicade created. Polices SRx-0041, SRx-0042, and SRx-0043 archived. For diagnosis of AS: trial of Humira and Enbrel requirement was added. For CD: Pediatric Crohn's Disease Activity Index (PCDAI) and Crohn's Disease Activity Index (CDAI) were requirements added; trial of Humira was added. For diagnosis of PsO: immunosuppressive drug criterion was separated from phototherapies and topical agents' trials; Psoriasis Area and Severity Index (PASI) score requirement was added; trials of Humira and Enbrel were added. For PsA: trials of Humira and Enbrel were added. For RA: non-biologic DMARDS were listed and criterion was added to use drug in combination with methotrexate, or if intolerant to methotrexate, use another immunosuppressant; trials of Humira and Enbrel were added. For UC: requirement for moderate to severe UC was revised, Pediatric Ulcerative Colitis Activity Index (PUCAI) was added. Trial of Humira required for member ≥ 18 y.o. List of diagnoses considered not medically necessary was added.
02/26/2019	Humira removed from trial criteria. Actemra, Cimzia, Cosentyx, Enbrel, Kevzara, Olumiant, Otezla, Siliq and Xeljanz added to trial agents list. TB test allowed to be done within 12 months prior to initiation of therapy; chest x-ray option removed. Symptoms of back pain for AS extended till before age of 50. Other drugs options allowed for PsA if there is an intolerance or contraindication to methotrexate.

References:

1. Remicade [prescribing information]. Horsham, PA; Janssen Biotech, Inc.: January, 2015.
2. Remicade. Lexi-Drugs. Lexicomp. Wolters Kluwer Health, Inc. Riverwoods, IL. Available at: <http://online.lexi.com>. Accessed February 28, 2017.
3. Lofberg R. Treatment of fistulas in Crohn's disease with infliximab. *Gut*. 1999;45(5):642-643.
4. Lichtenstein GR, Hanauer SB, Sandborn WJ, Practice Parameters Committee of American College of Gastroenterology. Management of Crohn's disease in adults. *American Journal of Gastroenterology* 2009;104(2):465-83; quiz 464, 484. DOI: 10.1038/ajg.2008.168.
5. Terdiman JP, Gruss CB, Heidelbaugh JJ, Sultan S, Falck-Ytter YT; AGA Institute Clinical Practice and Quality Management Committee. American Gastroenterological Association Institute guideline on the use of thiopurines, methotrexate, and anti-TNF-alpha biologic drugs for the induction and maintenance of remission in inflammatory Crohn's disease. *Gastroenterology*. 2013 Dec;145(6):1459-63.
6. Sandborn, W., Binion, D., Persley, K., Atreja, A., & Kosinski, L. (2014). AGA Institute Guidelines for the Identification, Assessment and Initial Medical Treatment in Crohn's Disease: Clinical Decision Support Tool. AGA Institute. Retrieved August 14, 2015, from www.gastro.org/IBDcarepathway.
7. Foundation for Sarcoidosis Research. <http://www.stopsarcoidosis.org/wp-content/uploads/2013/03/FSR-Physicians-Protocol1.pdf>.
8. How Is Sarcoidosis Treated? National Heart, Lung, and Blood Institute. Updated: June 14, 2013. Available at: <https://www.nhlbi.nih.gov/health/health-topics/topics/sarc/treatment>. Accessed February 28, 2017.
9. Ricart E, Sandborn WJ. Infliximab for the treatment of fistulas in patients with Crohn's disease. *Gastroenterology*. 1999;117(5):1247-1248.
10. Sands BE, Anderson FH, Bernstein CN et al. A randomized controlled trial of infliximab maintenance therapy for fistulizing Crohn's disease (ACCENT II). *N Engl J Med*. 2004;350:876-885.
11. American College of Rheumatology Ad Hoc Committee on Clinical Guidelines. Guidelines for the management of rheumatoid arthritis: *Arthritis Rheum*. 1996;39(5):713-723.
12. Hsu S, Papp KA, Lebwohl MG, et al. Consensus guidelines for the management of plaque psoriasis. *Arch Dermatol*. 2012 Jan;148(1):95-102.
13. American Gastroenterological Association. Identification, assessment and initial medical treatment in Cohn's disease. AGA institute. 2014. <http://www.gastro.org/IBDcarepathway>. Accessed April 20, 2017.



14. Terdiman JP, Gruss CB, Heidelbaugh JJ, et al. American Gastroenterological Association Institute Guideline on the Use of Thiopurines, Methotrexate, and Anti-TNF- α Biologic Drugs for the Induction and Maintenance of Remission in Inflammatory Crohn's Disease. *Gastroenterology* 2013; 145:1459-1463.

Effective date: 04/01/2019

Revised date: 02/26/2019