



MEDICAL POLICY STATEMENT

Nevada Medicaid

Policy Name & Number	Date Effective
Peroral Endoscopic Myotomy-NV MCD-MM-1803	09/01/2026
Policy Type	
MEDICAL	

Medical Policy Statements are developed from a review of the available evidence-based, clinical information based on and supported by clinical guidelines from medical specialties, peer-reviewed medical and scientific studies, nationally recognized utilization and technology assessment guidelines, and other medical management industry standards to aid in determining the medical necessity of a service. Medically necessary services are defined in the plan-specific Evidence of Coverage, Member Handbooks, Provider Manuals, Medical Policy Statements, and/or other plan policies and procedures.

Medical Policy Statements do not guarantee an authorization or payment of services. Please refer to the plan contract (e.g., Evidence of Coverage, Member Handbook) for the service(s) referenced in the Medical Policy Statement. Except as otherwise required by law, if there is a conflict between the Medical Policy Statement and the plan contract and/or provider agreement, then the plan contract and/or provider agreement will be the controlling document. Reasonable discretion may be used in interpreting and applying this Policy Statement to services provided in a particular case, and the policy may be modified at any time. Medical Policy Statements may be superseded by federal and state laws and regulations, as applicable. In the case of a discrepancy between the Policy Statement's effective date and any applicable legal or regulatory requirement, the law or regulation will govern.

Coverage for mental health and substance use disorder (MH/SUD) benefits are provided at the same level as coverage for medical and surgical (M/S) benefits, as required by the Mental Health Parity and Addiction Equity Act (MHPAEA). MH/SUD benefits are not subject to limits or requirements that are more restrictive than those that apply to M/S benefits.

Table of Contents

A. Subject	2
B. Background	2
C. Definitions.....	2
D. Policy	3
E. Conditions of Coverage	4
F. Related Policies/Rules	4
G. Review/Revision History	4
H. References	4

A. Subject

Peroral Endoscopic Myotomy

B. Background

Achalasia is a rare motility disorder that results from the degeneration of ganglion cells in the wall of the esophagus leading to failed relaxation of the lower esophageal sphincter and loss of esophageal peristalsis; the cause is unknown. As the sphincter becomes paralyzed and dilated over time, difficulty may occur passing food and liquid to the stomach. Symptoms include chest pain, heart burn, and difficulty belching. Potential complications are progressive dilation of the esophagus and increased risk of esophageal cancer. Although achalasia cannot be cured, symptoms may be controlled with treatment such as oral medications, dilation or stretching of the esophagus, open or laparoscopic surgery, endoscopic surgery, or injection of muscle-relaxing medicines (such as botulinum toxin) directly into the esophagus.

While laparoscopic Heller myotomy (LHM) is the most common surgical procedure for the treatment of achalasia, peroral endoscopic myotomy (POEM) is a newer, less invasive surgical alternative. The POEM procedure is performed through a submucosal tunnel layer of the esophagus and proximal stomach endoscopically without the need for incisions in the chest or abdomen. Since POEM does not require external incisions, this approach aims to reduce procedure-related pain and return patients to regular activities sooner than surgical modalities.

A review of clinical data suggests that the overall efficacy of POEM is similar to surgical myotomy and better than pneumatic dilation. However, patients who undergo POEM show an increased risk of gastroesophageal reflux disease (GERD) and erosive esophagitis which should be discussed.

C. Definitions

- **Achalasia** – A rare disorder making it difficult for food and liquid to pass from the esophagus to the stomach. In achalasia, nerve cells in the esophagus degenerate and as a result, the lower end of the esophagus, the lower esophageal sphincter (LES), fails to open to allow food into the stomach, leading to complications (eg, coughing, choking, aspiration pneumonia, ulceration, and weight loss). There are three different types of achalasia:
 - **Type I (Classic Achalasia)** – Characterized by minimal esophageal pressurization, this type is characterized by the incomplete relaxation of the LES, a lack of mobility in terms of contraction and relaxation, and a small amount of pressure built up in the esophagus.
 - **Type II (Pan-esophageal Pressurization)** – Indicated by esophageal compression, this type is more severe with more massive compression in the esophagus, often caused by the failure to relax and the build-up of pressure in the esophagus, typically from food.
 - **Type III (Spastic Achalasia)** – Characterized by spasms that result in sudden, abnormal squeezing of the esophagus and the LES, this type is the most severe and can also elicit the most severe symptoms (eg, severe chest pains that may

mimic those of a heart attack and spasms severe enough to wake a person from sleep).

- **Eckardt Score** – Standardized, scored assessment used to evaluate symptoms, stages, and efficacy of achalasia treatment. It attributes points (0 to 3 points) for four symptoms of the disease (dysphagia, regurgitation, chest pain, and weight loss) with scores ranging from 0 to 12.
- **Gastroesophageal Reflux Disease (GERD)** – A chronic disorder that occurs when stomach bile or acid flows into the esophagus and irritates the lining.
- **Laparoscopic Heller Myotomy (LHM)** – A minimally invasive, surgical procedure. The procedure involves making cuts in the opening of the lower esophageal sphincter (LES) to allow food to pass more easily.
- **Pneumatic Balloon Dilation (PD)** – An endoscopic therapy for achalasia. An air-filled cylinder-shaped balloon disrupts the muscle fibers of the lower esophageal sphincter, which is too tight in patients with achalasia.

D. Policy

- I. CareSource considers the POEM procedure to be medically necessary when all the following clinical criteria are met:
 - A. Member has a diagnosis of primary achalasia, type I, II, or III.
 - B. POEM is being proposed after the member has tried and failed conventional therapy, including pneumatic dilation or is not a surgical candidate for Heller myotomy.
 - C. Member has an Eckardt score of greater than or equal to 3.
 - D. Member has no history of previous open surgery of the stomach or esophagus.
- II. Request for members 18 years or younger will undergo individualized medical necessity review based on disease severity and available evidence.
- III. POEM for any other indication is considered experimental, investigational, and unproven.
- IV. The following conditions may represent contraindications and, if present, warrant individualized clinical assessment:
 - A. severe erosive esophagitis
 - B. significant coagulation disorders
 - C. liver cirrhosis with portal hypertension
 - D. severe pulmonary disease
 - E. esophageal malignancy
 - F. prior therapy that may compromise the integrity of the esophageal mucosa or lead to submucosal fibrosis, including recent esophageal surgery, radiation, endoscopic mucosal resection, or radiofrequency ablation
- V. Previous therapies for achalasia (eg, PD, botulinum toxin injection, or LHM) are not contraindications to POEM.

VI. Members receiving POEM should be made aware there is a high risk in developing GERD and should be advised as such prior to undergoing the procedure.

E. Conditions of Coverage
NA

F. Related Policies/Rules
NA

G. Review/Revision History

DATE		ACTION
Date Issued	08/27/2025	New market, approved at Committee.
Date Revised	06/17/2026	Annual review; background, definitions, and references updated; revised Sec D. II. and D. IV. Approved at Committee.
Date Effective	09/01/2026	
Date Archived		

H. References

1. Aiolfi A, Bona D, Riva CG, et al. Systematic review and bayesian network meta-analysis comparing laparoscopic Heller myotomy, pneumatic dilatation, and peroral endoscopic myotomy for esophageal achalasia. *J Laparoendosc Adv Surg Tech A*. 2020;30(2):147-155. doi:10.1089/lap.2019.0432
2. Familiari P, de Andreis FB, Landi R, et al. Long versus short peroral endoscopic myotomy for the treatment of achalasia: results of a non-inferiority randomized controlled trial. *Gut*. 2023;72(8):1442-1450. doi:10.1136/gutjnl-2021-325579
3. Health Technology Assessment: Peroral Endoscopic Myotomy for Treatment of Esophageal Achalasia. Hayes; 2019. Reviewed March 7, 2023. Accessed May 15, 2026. www.evidence.hayesinc.com
4. Huang Z, Cui Y, Li Y, et al. Peroral endoscopic myotomy for patients with achalasia with previous Heller myotomy: a systematic review and meta-analysis. *Gastrintest Endosc*. 2021;93(1):47-56.e5. doi:10.1016/j.gie.2020.05.056
5. Khashab MA, Vela MF, Thosani N, et al. ASGE guideline on the management of achalasia. *Gastrointest Endosc*. 2020;91(2):213-227. doi:10.1016/j.gie.2019.04.231
6. Khashab MA, Kumbhari V, Tieu AH, et al. Peroral endoscopic myotomy achieves similar clinical response but incurs lesser charges compared to robotic Heller myotomy. *Saudi J Gastroenterol*. 2017;23(2):91-96. doi:10.4103/1319-3767.203360
7. Kohn GP, Dirks RC, Ansari MT, et al. SAGES guidelines for the use of peroral endoscopic myotomy (POEM) for the treatment of achalasia. *Surg Endosc*. 2021;35(5):1931-1948. doi:10.1007/s00464-020-08282-0
8. Medicaid services manual: section 603. Division of Health Care Financing and Policy. Accessed May 15, 2026. www.dhcfp.nv.gov
9. Meng F, Li P, Wang Y, et al. Peroral endoscopic myotomy compared with pneumatic dilation for newly diagnosed achalasia. *Surg Endosc*. 2017;31(11):4665-4672. doi:10.1007/s00464-017-5530-0

10. Patel DA, Lappas BM, Vaezi MF. An overview of achalasia and its subtypes. *Gastroenterol Hepatol*. 2017;13(7):411-421. Accessed May 15, 2026. www.ncbi.nlm.nih.gov
11. Schneider AM, Louie BE, Warren HF, et al. A matched comparison of per oral endoscopic myotomy to laparoscopic Heller myotomy in the treatment of achalasia. *J Gastrointest Surg*. 2016;20(11):1789-1796. doi:10.1007/s11605-016-3232-x
12. Spechler SJ. Achalasia: pathogenesis, clinical manifestations, and diagnosis. UpToDate. Updated July 3, 2024. Accessed May 15, 2026. www.uptodate.com
13. Tan S, Zhong C, Ren Y, et al. Efficacy and safety of peroral endoscopic myotomy in achalasia patients with failed previous intervention: a systematic review and meta-analysis. *Gut Liver*. 2021;15(2):153-167. doi:10.5009/gnl19234
14. Vaezi MF, Pandolfino JE, Yadlapati RH, et al. ACG clinical guidelines: diagnosis and management of achalasia: diagnosis and management. *Am J Gastroenterol*. 2020;115(9):1393-1411. doi:10.14309/ajg.0000000000000731
15. Vespa E, Pellegatta G, Chandrasekar VT, et al. Long-term outcomes of peroral endoscopic myotomy for achalasia: a systematic review and meta-analysis. *Endoscopy*. 2023;55(2):167-175. doi:10.1055/a-1894-0147