



MEDICAL POLICY STATEMENT OHIO MEDICAID

Original Issue Date	Next Annual Review		Effective Date
06/21/2004	05/01/2019		05/01/2018
Policy Name			Policy Number
Obesity Surgery			MM-0026
Policy Type			
MEDICAL	Administrative	Pharmacy	Reimbursement

Medical Policy Statements prepared by CSMG Co. and its affiliates (including CareSource) are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other policies and procedures.

Medical Policy Statements prepared by CSMG Co. and its affiliates (including CareSource) do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage) for the service(s) referenced in the Medical Policy Statement. If there is a conflict between the Medical Policy Statement and the plan contract (i.e., Evidence of Coverage), then the plan contract (i.e., Evidence of Coverage) will be the controlling document used to make the determination.

Contents of Policy

<u>MEDICAL POLICY STATEMENT</u>	1
<u>TABLE OF CONTENTS</u>	1
<u>A. SUBJECT</u>	2
<u>B. BACKGROUND</u>	2
<u>C. DEFINITIONS</u>	3
<u>D. POLICY</u>	3
<u>E. CONDITIONS OF COVERAGE</u>	4
<u>F. RELATED POLICIES/RULES</u>	4
<u>G. REVIEW/REVISION HISTORY</u>	4
<u>H. REFERENCES</u>	4



A. SUBJECT

Obesity Surgery

B. BACKGROUND

Obesity is a major health threat in the United States facing a large proportion of adults and children. The Centers for Disease Control and Prevention (CDC) estimate that over 37% of adults in the United States, older than the age of 20 are obese (2016). Only tobacco has a higher modifiable risk factor in adult mortality. Obesity will become the number one modifiable risk factor in adult mortality if it continues to trend at the current rate. Obesity related health problems include: hypertension, Type II diabetes, hyperlipidemia, atherosclerosis, heart disease and stroke, diseases of the gallbladder, osteoarthritis, sleep apnea and certain cancers. The early implementation of healthy life styles remains the initial direction at impacting the rising epidemic of obesity. Obesity surgery must not be experimental or investigational, must meet current standard of care guidelines, and any device utilized must be FDA approved.

Bariatric surgery has been shown to have a positive effect on psychosocial functioning, however 20% of patients fail to achieve significant weight loss. A National Institutes of Health (NIH) consensus panel concluded that patients contemplating bariatric surgery should undergo pre-surgery psychological evaluation along with monitoring and addressing of psychological and behavioral factors pre- and post-surgery.

Obesity is clinically defined using the body mass index (BMI). BMI is the most common measure used to measure relative weight in comparison in children and adults.

The National Heart, Lung and Blood Institute (NHLBI) classify the ranges of BMI in adults as follows (NHLBI, 1998):

- <18.5 - Underweight
- 18.5 to 24.9 kg/m² - Normal
- 25-29.9 kg/m² - Overweight
- 30-34.9 kg/m² - Obesity Class I
- 35-39.9 kg/m² - Obesity Class II
- greater than 40 kg/m² - Extreme Obesity Class III

Additionally groups of individuals who have a BMI greater than 50 have been identified as super – obese.

Professional Societies

The following professional societies' recommendations are derived from the latest guidelines and scientific based literature available.

American Diabetes Association (ADA)

The ADA recommends bariatric surgery should be a recommended option to treat type 2 diabetes in appropriate surgical candidates with a body mass index (BMI) of 40 or greater regardless of glycemic control, and patients with a BMI of 35.0 to 39.9 with inadequately controlled hyperglycemia despite lifestyle and optimal medical therapy. Surgery should also be considered as an option to treat type 2 diabetic patients with a BMI of 30.0 to 34.9 and inadequately controlled hyperglycemia despite optimal treatment with medications. (2016)

American Society for Metabolic and Bariatric Surgery (ASMBS), American Association of Clinical Endocrinologists (AACE) and The Obesity Society (TOS)

The ASMBS in a joint commission with the AACE and the TOS recommends bariatric surgery be offered for patients with a BMI of 40 or greater and without a coexisting medical condition and for whom the surgery would not be considered high risk and for patients with a BMI of 35 or greater and 1 or more co-morbidities (2013).



Society of American Gastrointestinal and Endoscopic Surgeons (SAGES)

The Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) recommends bariatric surgery for the following (2008):

- BMI 35 to 40 with obesity-related co-morbid medical conditions
- BMI \geq 40 without co-morbidity if the weight adversely affects the patient
- Demonstration that dietary attempts at weight control have been ineffective
- Patients are motivated and well-informed
- Patients are free of psychological disease

National Institute of Diabetes and Digestive and Kidney Diseases (NIDDKD)

The National Institutes of Diabetes and Digestive and Kidney Diseases recommend bariatric surgery for the following (2011):

- BMI of greater than 40
- BMI of greater than 35 with serious comorbidity including:
 - Type 2 Diabetes
 - Heart Disease
 - Severe Sleep Apnea

C. DEFINITIONS

- **Body Mass Index (BMI):** The National Heart, Lung and Blood Institute defines BMI as a measure of body fat based on height and weight that applies to adult men and women.

D. POLICY

- I. The surgery should be considered medically necessary if **ALL** of the following conditions are met:
 - A. Prior authorization requests and supporting clinical information for Obesity Surgery have been submitted by the surgeon intending to perform the procedure. Further, supporting clinical information may be necessary from the PCP and/or other practitioners.
 - B. The patient is at least 20 years of age. Members less than 20 years old will be considered only after sustained efforts to lose weight have been made unsuccessfully and are considered medically necessary under the CareSource Obesity Surgery Adolescent policy.
 - C. The BMI (Body Mass Index) and associated conditions suggest surgery is the most prudent treatment:
 1. Patient has a BMI of 40 Obesity Class III or greater
 2. Patient has a BMI of 35 Obesity Class II or greater and at least two of the following co-morbid conditions related to obesity:
 - 2.1 *Obesity hypoventilation*
 - 2.2 *Obstructive sleep disorder* diagnosis, not otherwise well-controlled by standard therapies.
 - 2.3 Poorly controlled hypertension on multi-drug therapy
 - 2.4 Inadequately controlled diabetes despite optimal medical therapy (as outlined by the ADA).
 - 2.5 Coronary Artery Disease
 - 2.6 Pulmonary HTN
 3. The co-morbid condition is either poorly controlled on appropriate medical therapy and would likely improve with weight reduction OR by virtue of family history and existing clinical conditions, the patient would remain high risk for short term co-morbid complications without the surgery.

D. Correctable cause for obesity not identified (e.g., endocrine disorder)



- E. Written clinical documentation and supporting information from the attending surgeon or relevant specialist(s) have been submitted and includes all of the following:
 - 1. Letter of medical necessity supporting letter of medical necessity from the PCP or appropriate specialist.
 - 2. Evidence that there has been at least a 6 month documented physician supervised trial of diet and exercise within the last 24 months.
 - 3. A description of a multi-disciplinary approach, including, patient demonstrated preparedness, regarding management of the patient during the pre-operative, and an extended post-operative period.
 - 4. Documentation from the treating psychiatrist, and/or independently-licensed psychologist illustrating the patient has been evaluated from a psychological standpoint within the past 6 months including:
 - 4.1 Psychiatric diagnoses
 - 4.2 Relative contraindications such as:
 - a. Current acute or inadequately managed mental illness, particularly active psychosis and recent or active suicidal ideation
 - b. Active substance abuse or dependence
 - c. Inability to participate in informed decision making
 - d. An unwillingness or inability to comply with pre and post-surgical protocols
 - 4.3 Readiness for change: Surgery candidates must be able to tolerate the risks of surgery and comply with a postoperative regimen that requires long-term medical surveillance and lifelong changes in diet and activity. Adherence to behavioral regimens, medication and medical appointments may provide insight into member's ability to be successful after surgery.
 - 4.4 Patient expectations relative to the surgical and post-surgical process: The psychiatric assessment is used to clarify expectations regarding surgery and to identify psychosocial issues that may require attention over time (e.g. changes in behaviors and relationships)
 - 4.5 Post-surgical behavioral monitoring: Psychiatric assessment should outline the post-operative plan of care for any behavioral health issues that have been identified.
- II. Inadequate weight loss due to individual noncompliance with postoperative nutrition and exercise recommendations is not a medically necessary indication for revision or conversion surgery and is not covered.

E. CONDITIONS OF COVERAGE

HCPCS
CPT

AUTHORIZATION PERIOD

F. RELATED POLICIES/RULES

N/A

G. REVIEW/REVISION HISTORY

	DATES	ACTION
Date Issued	09/21/2004	New Policy.
Date Revised	10/17/2017	Annual update
Date Effective	05/01/2018	

H. REFERENCES

1. AAP Updates Recommendations on Obesity Prevention: It's Never Too Early to Begin Living a Healthy Lifestyle. (n.d.). Retrieved June 20, 2016, from <https://www.aap.org/en-us/about-the->



aap/aap-press-room/pages/AAP-Updates-Recommendations-on-Obesity-Prevention-It's-Never-Too-Early-to-Begin-Living-a-Healthy-Lifestyle.aspx

2. American Society for Metabolic and Bariatric Surgery <https://asmbs.org/resources/clinical-practice-guidelines-for-the-perioperative-nutritional-metabolic-and-nonsurgical-support-of-the-bariatric-surgery-patient>
3. Bariatric Surgery | NIDDK. (n.d.). Retrieved from <https://www.niddk.nih.gov/health-information/weight-management/bariatric-surgery#clinicaltrials>
4. Bariatric procedures for the management of severe obesity: Descriptions. (n.d.). Retrieved June 22, 2016, from <http://www.uptodate.com/contents/bariatric-procedures-for-the-management-of-severe-obesity-descriptions>
5. Buchwald, H., Avidor, Y., Braunwald, E., Jensen, M. D., Pories, W., Fahrbach, K., & Schoelles, K. (2004). Bariatric Surgery. *JAMA*, 292(14), 1724. doi:10.1001/jama.292.14.1724
6. CDC. (2016, April 27). Obesity and overweight. Retrieved July 15, 2016, from <http://www.cdc.gov/nchs/fastats/obesity-overweight.htm>
7. Chapman, A. (2004). Laparoscopic adjustable gastric banding in the treatment of obesity: A systematic literature review. *Surgery*, 135(3), 326-351. doi:10.1016/s0039-6060(03)00392-1
8. Decision Memo for Bariatric Surgery for the Treatment of Morbid Obesity (CAG-00250R). U.S. Dept. of Health and Human Services, Centers for Medicare and Medicaid Services, February 21, 2006.
9. Kalarchian, M. (2007). Psychiatric Disorders Among Bariatric Surgery Candidates: Relationship to Obesity and Functional Health Status. *American Journal of Psychiatry*, 164(2), 328. doi:10.1176/appi.ajp.164.2.328
10. Late complications of bariatric surgical operations. (n.d.). Retrieved June 22, 2016, from https://www.uptodate.com/contents/late-complications-of-bariatric-surgical-operations?source=search_result
11. Marcus, M. D., Kalarchian, M. A., & Courcoulas, A. P. (2009). Psychiatric evaluation and follow-up of Bariatric surgery patients. *American Journal of Psychiatry*, 166(3), 285–291. doi:10.1176/appi.ajp.2008.08091327
12. Milliman Care Guidelines (MCG): Ambulatory Care Guidelines, 21st Ed., 2017.
13. New Diabetes Guidelines Include Recommendations for Bariatric Surgery. (n.d.). Retrieved June 21, 2016, from <https://www.hayesinc.com/subscribers/displaySubscriberArticle.do?>
14. **The Practical Guide to Identification and Treatment of Overweight and Obesity in Adults.** (n.d.). Retrieved June 21, 2016, from http://www.nhlbi.nih.gov/files/docs/guidelines/prctgd_c.pdf
15. **Repeat Bariatric Surgery for Patients Who Have Not Reached Weight-loss Goals after Previous Surgery.** (n.d.). Retrieved June 22, 2016, from <https://www.ecri.org/components/Hotline/Pages/14173.aspx>
16. Shekelle, P. G. (n.d.). Mental Health Assessment and Psychological Interventions for bariatric surgery. Retrieved from <https://www.hsrd.research.va.gov/publications/esp/bariatric-REPORT.pdf>
17. Updated Guidelines for Bariatric Surgery. (n.d.). Retrieved June 21, 2016, from <https://www.hayesinc.com/subscribers/displaySubscriberArticle.do?>

This guideline contains custom content that has been modified from the standard care guidelines and has not been reviewed or approved by MCG Health, LLC.

The Medical Policy Statement detailed above has received due consideration as defined in the Medical Policy Statement Policy and is approved.