



MEDICAL POLICY STATEMENT OHIO MEDICAID

Original Issue Date	Next Annual Review	Effective Date	
05/15/2009	05/01/2019	05/01/2018	
Policy Name		Policy Number	
Obesity Surgery in Adolescents		MM-0027	
Policy Type			
MEDICAL	Administrative	Pharmacy	Reimbursement

Medical Policy Statements prepared by CSMG Co. and its affiliates (including CareSource) are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other policies and procedures.

Medical Policy Statements prepared by CSMG Co. and its affiliates (including CareSource) do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage) for the service(s) referenced in the Medical Policy Statement. If there is a conflict between the Medical Policy Statement and the plan contract (i.e., Evidence of Coverage), then the plan contract (i.e., Evidence of Coverage) will be the controlling document used to make the determination.

Contents of Policy

<u>MEDICAL POLICY STATEMENT</u>	1
<u>TABLE OF CONTENTS</u>	1
<u>A. SUBJECT</u>	2
<u>B. BACKGROUND</u>	2
<u>C. DEFINITIONS</u>	3
<u>D. POLICY</u>	3
<u>E. CONDITIONS OF COVERAGE</u>	4
<u>F. RELATED POLICIES/RULES</u>	4
<u>G. REVIEW/REVISION HISTORY</u>	4
<u>H. REFERENCES</u>	4



A. SUBJECT

Obesity Surgery in Adolescents

B. BACKGROUND

Childhood obesity continues to be a serious health problem in the United States. The Centers for Disease Control (CDC) estimate the prevalence of obesity to be approximately 17% affecting 12.7 million children and adolescents (2015). Severely obese children and adolescents are at risk for developing serious comorbidities, including obstructive sleep apnea, diabetes, hypertension, cardiac hypertrophy, and nonalcoholic fatty liver disease (NAFLD). They may also develop depression and suffer from impaired quality of life. The early implementation of healthy life styles remains the initial direction at impacting the rising epidemic of obesity.

Professional Societies

The following professional societies' recommendations are derived from the latest guidelines and scientific based literature available.

American Academy of Pediatrics

The American Academy of Pediatrics (AAP) recommends patients being considered for bariatric surgery "be physically mature, have a BMI of 50 or greater or 40 or greater with significant comorbidities, have experienced failure of a formal 6 month weight loss program and be capable of adhering to the long-term lifestyle changes required after surgery." In addition, adolescents "should be referred to specialized centers with a multidisciplinary bariatric team capable of providing long-term follow-up care. (2004)

National Institute of Diabetes, Digestive and Kidney Diseases (NIDDK)

The National Institute of Diabetes, Digestive and Kidney Diseases (NIDDK) (2011) suggest surgery for youth candidates only when they have tried for at least 6 months to lose weight with no success and met the following criteria:

- Extreme obesity (BMI of 40 or greater)
- Reached adult height (usually 13 or older for girls and 15 or older for boys)
- Serious health problems linked to weight (DM Type II, Sleep Apnea) that may improve with surgery
- A thorough assessment by health care providers documenting the patient's and the patient's parent's emotional preparedness and ability to make lifestyle changes
- Referral to a special youth bariatric surgery center

Growth and Development in Adolescents

For adolescents, physical development and maturation may be determined utilizing the gender specific growth chart and BMI chart

Estimated adult height may also be calculated utilizing the Mid-Parental height calculation:

Boy

- In: $(\text{Father's Height} + \text{Mother's Height} + 5) / 2$
- Cm: $(\text{Father's Height} + \text{Mother's Height} + 13) / 2$

Girl

- In: $(\text{Father's Height} - 5 + \text{Mother's Height}) / 2$
- Cm: $(\text{Father's Height} - 13 + \text{Mother's Height}) / 2$



Tanner Stage	Male	Female	Pubic Hair (Male and Female)
I	Prepubertal	Prepubertal	Prepubertal (can see velus hair similar to abdominal wall)
II	Enlargement of scrotum and testes; scrotum skin reddens and changes in texture	Breast bud stage with elevation of breast and papilla; enlargement of areola	Sparse growth of long, slightly pigmented hair, straight or curled, at base of penis or along labia
III	Enlargement of penis (length at first); further growth of testes	Further enlargement of breast and areola; no separation of their contour	Darker, coarser and more curled hair, spreading sparsely over junction of pubes
IV	Increased size of penis with growth in breadth and development of glans; testes and scrotum larger, scrotum skin darker	Areola and papilla form a secondary mound above level of breast	Hair adult in type, but covering smaller area than in adult; no spread to medial surface of thighs
V	Adult genitalia	Mature stage: projection of papilla only, related to recession of areola	Adult in type and quantity, with horizontal distribution ("feminine")

C. DEFINITIONS

- **Mid-Parental Height:** (see above) This calculation provides a target mean and range for the genetic potential of a child based upon the biologic parents' heights. This calculation *alone* is not sufficient to predict final height; it only calculates a *reference* range for assessing growth
- **Body Mass Index For Age Percentile: (BMI):** The CDC defines BMI as a person's weight in kilograms divided by the square of height in meters. BMI is age and sex related for children and teens and is often referred to as BMI-for-age.

D. POLICY

- I. Obesity surgery for adolescents may be considered medically necessary if the following are met:
 - A. An adolescent candidate is morbidly obese with a BMI of 50 or greater
 - B. An adolescent candidate is morbidly obese with a BMI of 40 or greater with one of the following comorbidities expected to resolve with surgery:
 1. DM Type II
 2. Obstructive Sleep Apnea
 3. Heart disease
 4. Poorly controlled Hypertension
 - C. The youth will have attained a majority skeletal maturity (equal to or greater than 13 years for girls and equal to or greater than 15 years of age for boys).
 - D. The potential candidate for bariatric surgery should be referred to specified centers with multi-disciplinary weight management teams that have expertise in meeting the needs of adolescents, including the immediate availability of critical care services, psychology, nutrition, and physical activity instruction.
 - E. Written clinical documentation and supporting information for the attending surgeon must include:
 1. Letter of medical necessity from the PCP or appropriate specialist



2. Evidence that there have been adequate conservative attempts at weight loss for at least a 6 month period
3. A description of a multi-disciplinary approach to preparing and managing the patient in the pre-operative periods, peri-operative periods and through an extended post-operative period
4. Documentation illustrating the patient has been evaluated from a psychological standpoint within the past 6 months by the treating behavioral health provider including:
 - 4.1 Evidence that any co-existing psychiatric condition is stable
 - 4.2 Evidence the patient can withstand the rigors of surgery
 - 4.3 Evidence the patient can adhere with long-term follow-up care
 - 4.4 Assessment, listing of diagnoses, and treatment plan must be provided.
- F. The intended procedure must not be experimental or investigational, must meet current standard of care guidelines, and any device utilized must be FDA approved.
- G. The patient should not have a current or planned pregnancy within 12 to 18 months of surgery.

E. CONDITIONS OF COVERAGE

HCPCS

CPT

AUTHORIZATION PERIOD


F. RELATED POLICIES/RULES

G. REVIEW/REVISION HISTORY

DATES		ACTION
Date Issued	05/15/2009	New Policy.
Date Revised	10/28/2017	Revisions include Tanner Stages, Growth Charts and BMI charts
Date Effective	05/01/2018	

H. REFERENCES

1. AAP Updates Recommendations on Obesity Prevention: It's Never Too Early to Begin Living a Healthy Lifestyle. (n.d.). Retrieved October 3, 2017, from <https://www.aap.org/en-us/about-the-aap/aap-press-room/pages/AAP-Updates-Recommendations-on-Obesity-Prevention-It's-Never-Too-Early-to-Begin-Living-a-Healthy-Lifestyle.aspx?>
2. Bariatric surgery for severe obesity. (2016, July 27). Retrieved October 3, 2017, from <https://www.niddk.nih.gov/health-information/health-topics/weight-control/bariatric-surgery-severe-obesity/Pages/bariatric-surgery-for-severe-obesity.aspx#b>
3. Childhood Obesity Facts. (2015). Retrieved October 3, 2017, from <http://www.cdc.gov/obesity/data/childhood.html>
4. CDC. (2015, May 15). About child & teen BMI. Retrieved October 3, 2017, from http://www.cdc.gov/healthyweight/assessing/bmi/childrens_bmi/about_childrens_bmi.html
5. Garcia, V., & DeMaria, E. (2006). Adolescent Bariatric Surgery: Treatment Delayed, Treatment Denied, a Crisis Invited. *Obesity Surgery*, 16(1), 1-4. doi:10.1381/096089206775222195
6. Guidelines for Clinical Application of Laparoscopic Bariatric Surgery - A SAGES Publication. (n.d.). Retrieved from <https://www.sages.org/publications/guidelines/guidelines-for-clinical-application-of-laparoscopic-bariatric-surgery/>
7. Hayes Medical Technology Directory June 7, 2007, Pediatric Bariatric Surgery for Morbid Obesity.
8. Inge, T. H. (2004). Bariatric Surgery for Severely Overweight Adolescents: Concerns and Recommendations. *PEDIATRICS*, 114(1), 217-223. doi:10.1542/peds.114.1.217



9. Krebs, N. F., Himes, J. H., Jacobson, D., Nicklas, T. A., Guilday, P., & Styne, D. (2007). Assessment of Child and Adolescent Overweight and Obesity. *PEDIATRICS*, 120(Supplement), S193-S228. doi:10.1542/peds.2007-2329d
10. Ludwig, D. S. (2007). Childhood Obesity — The Shape of Things to Come. *New England Journal of Medicine*, 357(23), 2325-2327. doi:10.1056/nejmp0706538
11. Michalsky, M., Kramer, R. E., Fullmer, M. A., Polfuss, M., Porter, R., Ward-Begnoche, W., ... Reichard, K. W. (2011). Developing Criteria for Pediatric/Adolescent Bariatric Surgery Programs. *Pediatrics*, 128(Supplement 2), S65-S70. doi:10.1542/peds.2011-0480f
12. Recommendations for Treatment of Child and Adolescent Overweight and Obesity | SUPPLEMENT ARTICLES | *Pediatrics*. (2007, December). Retrieved from http://pediatrics.aappublications.org/content/120/Supplement_4/S254
13. Sugerman, H. (2003). Bariatric Surgery for Severely Obese Adolescents. *Journal of Gastrointestinal Surgery*, 7(1), 102-108. doi:10.1016/s1091-255x(02)00125-7
14. Up to Date (n.d.). Retrieved October 3, 2017, from https://www.uptodate.com/contents/surgical-management-of-severe-obesity-in-adolescents?source=search_result
15. Xanthakos, S. A., Daniels, S. R., & Inge, T. H. (2006). Bariatric surgery in adolescents: an update. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/?term=Bariatric+Surgery+in+Adolescents%3A++An+Update.++Adolescence+Medical+Clin+2006%3A+589-612>

The Medical Policy Statement detailed above has received due consideration as defined in the Medical Policy Statement Policy and is approved.

Archiving