



MEDICAL POLICY STATEMENT

Ohio Medicaid

Policy Name & Number	Date Effective
Sacroiliac Joint Procedures-OH MCD-MM-0010	12/01/2023
Policy Type	
MEDICAL	

Medical Policy Statement prepared by CareSource and its affiliates are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other policies and procedures.

Medical Policy Statements prepared by CareSource and its affiliates do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage) for the service(s) referenced in the Medical Policy Statement. If there is a conflict between the Medical Policy Statement and the plan contract (i.e., Evidence of Coverage), then the plan contract (i.e., Evidence of Coverage) will be the controlling document used to make the determination. According to the rules of Mental Health Parity Addiction Equity Act (MHPAEA), coverage for the diagnosis and treatment of a behavioral health disorder will not be subject to any limitations that are less favorable than the limitations that apply to medical conditions as covered under this policy.

Table of Contents

A. Subject.....	2
B. Background.....	2
C. Definitions	2
D. Policy	3
E. Conditions of Coverage.....	5
F. Related Polices/Rules	5
G. Review/Revision History.....	5
H. References.....	6

A. Subject**Sacroiliac Joint Procedures****B. Background**

Nearly 84% of adults experience back pain during their lifetime. Long-term outcomes are largely favorable for most patients, but a small percentage of patients' symptoms are persistent. Persistent pain is categorized as subacute when lasting between four and twelve weeks and chronic when persisting for at least three months.

Up to 10% to 25% of patients with persistent low back pain may have a component of pain related to sacroiliac joints (SIJ). Comprehensive pain management care plans are most effective in managing a patient's chronic pain. These plans focus on a person-centered approach and incorporate conservative treatment with other modalities. These multidisciplinary treatments include promoting patient self-management and aim to reduce the impact of pain on a patient's daily life, even if the pain cannot be relieved completely. In addition to conservative therapy, additional treatment options may include nonpharmacologic or pharmacologic treatments, nonsurgical interventions, and surgical interventions. Interventional procedures for the management of pain unresponsive to conservative treatment should be provided only by physicians qualified to deliver these health services.

Sacroiliac joint injections using local anesthetic and/or corticosteroid medication have been shown to be effective for diagnostic purposes but provide limited short-term relief from pain resulting from SIJ dysfunction. Long-term use has not been adequately studied to establish standards of care. Radiofrequency ablation (RFA) is another treatment method, which uses heat to destroy nerves. RFA for the treatment of low back pain has inconsistent results in the peer-reviewed medical literature with limited follow-up. However, clinical experience suggests that some patients obtain more significant relief from these procedures, making it reasonable to offer SIJ injections and/or RFA when conservative management has failed.

C. Definitions

- **Conservative Therapy** - A multimodality plan of care including both active and inactive conservative therapies.
 - **Active Conservative Therapies** - Actions or activities that strengthen supporting muscle groups and target key spinal structures, including physical therapy, occupational therapy, physician supervised home exercise program (HEP), and/or chiropractic care.
 - **HEP** - A six-week program requiring an exercise prescription and/or plan and a follow-up documented in the medical record after completion, or documentation of the inability to complete the HEP due to a stated physical reason (i.e., increased pain, inability to physically perform exercises). Patient inconvenience or noncompliance without explanation does not constitute an inability to complete.

- **Inactive Conservative Therapies** - Passive activities that aid in treating symptoms associated with pain, including rest, ice, heat, medical devices, TENS unit, and/or prescription medications.
 - **Transcutaneous Electrical Nerve Stimulator (TENS)** - A device that utilizes electrical current delivered through electrodes placed on the surface of the skin to decrease the patient's perception of pain by inhibiting the transmission of afferent pain nerve impulses and/or stimulating the release of endorphins. Its use, frequency, duration, and start dates must be documented in the medical record to be considered part of conservative therapy during the period of prior authorization request.
- **Radiofrequency Ablation (RFA)** - Minimally invasive treatment modality that percutaneously introduces an electrode under fluoroscopic guidance to thermocoagulate medial branches of the dorsal spinal nerves.
- **Sacroiliac Joint (SIJ) Injections** - Corticosteroid and local anesthetic therapeutic injections into the SIJ to treat pain that hasn't responded to conservative therapies.
- **Functional Disability** - Acquired difficulty in performing basic everyday tasks or more complex tasks needed for independent living.

D. Policy

I. Sacroiliac Joint Injections

- A. Diagnostic Injections: CareSource considers up to 2 diagnostic sacroiliac joint injections for the treatment of chronic low back pain medically necessary when **ALL** the following criteria are met:
 - 1. Somatic or nonradicular low back and lower extremity pain experienced for at least 3 months;
 - 2. Pain and tenderness are located in sacroiliac joint region and cause functional disability;
 - 3. Positive response to at least one SIJ pain provocation test (e.g., distraction, compression, thigh thrust, Gaenslen's, Patrick's test/FABER test, sacral thrust);
 - 4. Failure of conservative therapy, as evidenced by **ALL** the following:
 - a. Documentation in the medical record of at least 6 weeks of active conservative therapy (e.g., Physical therapy; occupational therapy; physician supervised home exercise program (HEP); acupuncture; chiropractic care) within the past 6 months OR inability to complete active conservative therapy due to contraindication, increased pain, or intolerance;
 - b. Documentation in the medical record of at least 6 weeks of inactive conservative therapy (e.g., rest, ice, heat, TENS use, pharmacotherapy [e.g., non-steroidal anti-inflammatory drugs, acetaminophen]) within the past 6 months;
 - 5. If a second diagnostic injection is requested, at least 1 week has passed since the initial injection.
- B. Therapeutic injections: CareSource considers therapeutic sacroiliac joint injections medically necessary when **ALL** the following criteria are met:

1. Previous SIJ injection led to significant functional pain relief as measured by a decrease in pain level (demonstrated on a qualitative pain scale), decrease in pain medications, and/or increase in physical function;
 2. Member experiences return of pain or deterioration in function;
 3. Injection is to be used in conjunction with other non-invasive treatment methods;
 4. Injections should not be repeated more frequently than every 2 months;
 5. No more than 4 injections total (diagnostic and therapeutic), have been administered at the same site the last 12 months.
- C. Exclusions/Limitations:
1. Codes 64451 and 27096 are considered the same procedure and may not be billed together. Only one code will be reimbursed.
 2. Image guidance and/or injection of contrast is included in sacroiliac injection procedures and may not be billed separately.
 3. If neural blockade is applied for different regions or different sides, injections are performed at least one week apart.
 4. Pain management literature highlighting controlled studies of SI joint pain management has not demonstrated injections of the SI joint to be effective as a long-term management modality. Long-term continuation may be subject to medical necessity review.
 5. Monitored anesthesia and conscious sedation are not medically necessary.
 6. The use of SIJ injections for the treatment of pain as a result of Herpes Zoster is not indicated due to insufficient evidence demonstrating efficacy in the peer-reviewed published literature.
- II. Radiofrequency Ablation of the SIJ
- A. Initial Radiofrequency Ablation of the SIJ
1. Radiofrequency ablation is considered medically necessary when **ALL** the following have been met in the last 6 months:
 - a. The clinical criteria above for failed conservative therapy (I.A.4.a. and I.A.4.b.) has been met; and
 - b. One diagnostic injection per joint to evaluate pain and attain therapeutic effect has been performed, with significant functional pain relief as measured by a decrease in pain level (demonstrated on a qualitative pain scale), decrease in pain medications, and/or increase in physical function.
- B. Repeat Radiofrequency Ablation of the SIJ
1. Conservative therapy and diagnostic injections are not required if there has been a reduction in pain for at least 12 months or more from the initial RFA within the last 36 months.
 2. When there has not been a repeat RFA in the last 36 months, a diagnostic injection is required.
 3. A maximum of 1 radiofrequency ablation for SIJ pain per side per rolling 12 months is considered medically necessary.
- C. Exclusions/Limitations
1. Codes 64451 and 27096 are considered the same procedure and may not be billed together. Only one code will be reimbursed.

2. The use of cooled RFA for SI joint-mediated low back pain is considered not medically necessary due to insufficient evidence demonstrating efficacy in the peer-reviewed published literature.
3. Pain management literature highlighting controlled studies of SI joint pain management has not demonstrated the effectiveness of RFA as a long-term management modality. Long-term continuation may be subject to medical necessity review.

III. Implantable Spinal Cord Stimulators

Patients with indwelling implanted spinal cord stimulators or pain pumps should have a device interrogation report submitted with medical records for a prior authorization request for proposed interventional pain injections. If a device is not functioning properly, an escalation in pain may warrant evaluation and management of the implanted device.

E. Conditions of Coverage

NA

F. Related Policies/Rules

NA

G. Review/Revision History

	DATE	ACTION
Date Issued	02/10/2015	New Policy
Date Revised	07/28/2015 07/26/2016 02/08/2018 03/06/2019 10/8/2019 05/13/2020 04/28/2021 03/29/2022 03/01/2023	Removed Herpes Criteria Annual Update: Removed start and end dates. Addition of PA clarification and documentation requirements. Revision of injection frequency. Removed joint fusion denial language. This service will be addressed in the Sacroiliac Joint Fusion policy. Annual Update: Added clinical criteria for coverage of radiofrequency ablation of the SI Joint. Added coding information. Annual Update: Removed PA language. Annual Review: Updated background and references, re-organized criteria into initial and subsequent procedures, type of procedure. Annual Review: restructured conservative management and clinical criteria, added provocation tests
Date Effective	12/01/2023	
Date Archived		

The MEDICAL Policy Statement detailed above has received due consideration as defined in the MEDICAL Policy Statement Policy and is approved.

H. References

1. Chou R, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med.* 2007 Oct;147(7):478-491. Retrieved January 20, 2023 from www.ncbi.nlm.nih.gov.
2. Chou R, et al. Nonpharmacologic therapies for low back pain: a systematic review for an American College of Physicians Clinical Practice Guideline. *Ann Intern Med.* 2017 Apr;166(7):493-505. Retrieved January 20, 2023 from www.ncbi.nlm.nih.gov.
3. Chou R. (2021, June). Subacute and chronic low back pain: nonsurgical interventional treatment. UpToDate. Retrieved January 20, 2023 from www.uptodate.com.
4. Hansen H, et al. A systematic evaluation of the therapeutic effectiveness of sacroiliac joint interventions. *Pain Phys.* 2012 May-Jun;15(3):E247-E278. Retrieved January 20, 2023 from www.painphysicianjournal.com.
5. Jamjoom AM, Saeedi RJ, Jamjoom AB. Placebo effect of sham spine procedures in chronic low back pain: a systematic review. *J Pain Res.* 2021 Sep;14:3057-3065. Retrieved January 20, 2023 from www.ncbi.nlm.nih.gov.
6. Lee DW, et al. Latest evidence-based application for radiofrequency neurotomy (LEARN): best practice guidelines from the American Society of Pain and Neuroscience (ASPN). *J Pain Res.* 2021;14:2807-2831. doi:10.2147/JPR.S235665.
7. Maas ET, et al. Radiofrequency denervation for chronic low back pain. *Cochrane Database Syst Rev.* 2015 Oct;2015(10):CD008572. doi:10.1002/14651858.CD008572.pub.
8. Manchikanti L, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part I: introduction and general considerations. *Pain Phys.* 2013 Apr;16(2 Suppl):S1-S48. Retrieved January 20, 2023 from www.ncbi.nlm.nih.gov.
9. Manchikanti L, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Phys.* 2013 Apr;16(2 Suppl):S49-S283. Retrieved January 20, 2023 from www.ncbi.nlm.nih.gov.
10. Manchikanti L, et al. Comprehensive evidence-based guidelines for facet joint interventions in the management of chronic spinal pain: American Society of Interventional Pain Physicians (ASIPP) Guidelines Facet Joint Interventions 2020 Guidelines. *Pain Phys.* 2020 May;23(3S):S1-S127. Retrieved January 20, 2023 from www.painphysicianjournal.com.
11. Ohio Administrative Code. (2022 March 1). Rule 5160-4-12. Immunizations, injections and infusions (including trigger-point injections), skin substitutes, and provider-administered pharmaceuticals. Retrieved January 30, 2023 from www.codes.ohio.gov.
12. Simopoulos T, et al. A systematic evaluation of prevalence and diagnostic accuracy of sacroiliac joint interventions. *Pain Phys.* 2012 May-Jun;15(3):E305-E344. Retrieved January 20, 2023 from www.ncbi.nlm.nih.gov.
13. Stelzer W, et al. Use of cooled radiofrequency lateral branch neurotomy for the treatment of sacroiliac joint-mediated low back pain: a large case series. *Pain Med.* 2013 Jan;14(1):29-35. Retrieved January 20, 2023 from www.ncbi.nlm.nih.gov.

14. Stevens AC, Carroll DD, Courtney-Long EA, et al. Adults with one or more functional disabilities – United States, 2011-2014. *MMWR Morb Mortal Wkly Rep* 2016;65:1021-1025. doi:10.15585/mmwr.mm6538a1

Independent medical review – March 2019

Approved by ODM 8/17/2023