

MEDICAL POLICY STATEMENT		
Effective Date	Next Annual Review Date	Last Review / Revision Date
5/2009	7/2015	7/2014
Author John Bloom, MD		



CSMG Medical Policy Statements are derived from literature based and supported clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services are those health care services or supplies which are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased or new morbidity, impairment of function, dysfunction of a body organ or part or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative and are not provided mainly for the convenience of the member or provider.

A. SUBJECT

Obesity Surgery in Adolescents

B. BACKGROUND

This document is to establish a policy to review the medical necessity for coverage of obesity surgery for adolescents.

The prevalence of extreme obesity in U.S. children and adolescents has been estimated at 4%. There is no standard treatment for morbid obesity. Low-calorie diets, exercise, behavioral modification regimens and medical treatment have generally been unsuccessful in long-term weight management for morbidly obese young people.

The early implementation of healthy life styles remains the initial direction at impacting the rising epidemic of obesity. There are the rare morbidly obese children who are candidates for bariatric surgery. The intended procedure must not be experimental or investigational, must meet current standard of care guidelines, and any devices utilized must be FDA approved.

C. POLICY

The surgery may be considered medically necessary if the following are met:

1. Adolescent candidates for bariatric surgery should be morbidly obese (defined by the World Health Organization as a BMI of equal or greater than 40).
2. These youth will have attained a majority skeletal maturity (equal or greater than 13 years for girls and equal or greater than 15 years of age for boys).
3. Adolescents have experienced failure of 6 continuous months of organized weight loss attempts and have met certain anthropometric, medical (R/O endogenous causes) and psychological evaluation of both patient and family.
4. Potential candidate for bariatric surgery should be referred to specified centers with multi-disciplinary weight management teams* that have expertise in meeting the needs of adolescents.

5. Cooperation, compliance and understanding by both the patient and their family are essential. This included both the pre-operative period as well as post-operative.
6. These adolescents will require a regular, prolonged post-operative follow up.
7. Written clinical documentation and supporting information for the attending surgeon must include:
 - a. Letter of medical necessity.
 - b. Evidence that there have been adequate conservative attempts at weight loss
 - c. Summary (support to meeting policy standard #1) of all co-morbid condition with their conservative treatment.
 - d. A description of a multi-disciplinary approach to preparing and managing the patient in the pre-operative periods, peri-operative periods and through an extended post-operative period.
 - e. Evidence the patient has been evaluated from a psychological standpoint.
 - f. Evidence the patient has the support and understanding of the family.
 - g. Supporting letter of medical necessity from the PCP.

The team should include specialists with expertise in adolescent obesity evaluation and management, psychology, nutrition, and physical activity instruction.

If there is no LCD or NCD present reference the CSMG Policy for coverage.

D. REVIEW / REVISION HISTORY

Date Issued: 5/2009

Date Revised: 5/2009

Date Reviewed: 5/2009, 7/1/2011, 3/2012, 7/2013, 7/2014

E. REFERENCES

1. Assessment of Child and Adolescent Overweight and Obesity: Dec 2007, A Supplement to PEDIATRICS
2. Inge TH, Krebs NP, Garcia VF et al Bariatric Surgery for Severely Overweight Adolescents: Concerns and Recommendations. Pediatrics 2004; 114:217-223
3. Sugeran HJ, DeNaruam EJ et al. Bariatric Surgery for Severely Obese Adolescents. J Gastrintest Surg. 2003; 1:102-108.
4. Xanthakos SA, Daniels SR, Inge TH. Bariatric Surgery in Adolescents: An Update. Adolescence Medical Clin 2006: 589-612.
5. Ludwig DH Childhood Obesity-The shape of things to come. NEJM 2007;357: 2325-2327.
6. Garcia VF, DeMaria EJ. Adolescent Bariatric surgery: Treatment delayed, Treatment denied, A Crisis Invited. Obesity Surgery 2006; 16: 1-3.
7. Hayes Medical Technology Directory June 7, 2007, Pediatric Bariatric Surgery for Morbid Obesity.

The Medical Policy Statement detailed above has received due consideration as defined in the Medical Policy Statement Policy and is approved.