

UTILIZATION MANAGEMENT MEDICAL POLICY

POLICY: Oncology (Intravesical) – Anktiva Utilization Management Medical Policy

- Anktiva® (nogapendekin alfa inbakicept-pmln intravesical solution – ImmunityBio)

REVIEW DATE: 04/08/2026

OVERVIEW

Anktiva, an interleukin-15 (IL-15) receptor agonist, is indicated with Bacillus Calmette-Guerin (BCG) for the treatment of **BCG-unresponsive non-muscle invasive bladder cancer (NMIBC)** in adults with carcinoma in situ with or without papillary tumors.¹

Dosing Information

Anktiva is for intravesical use only.¹ For induction therapy, the recommended dose is 400 mcg administered intravesically with BCG once weekly for 6 weeks. A second induction course can be administered if the patient did not achieve a complete response at Month 3. For maintenance therapy, the recommended dose is 400 mcg with BCG once weekly for 3 weeks at Months 4, 7, 10, 13, and 19. For patients with an ongoing complete response at Month 25, additional doses of 400 mcg plus BCG can be given once weekly for 3 weeks at Months 25, 31, and 37. Treatment can continue until disease persistence after the second course of induction therapy, disease recurrence or progression, unacceptable adverse events, or a maximum of 37 months.

Guidelines

The National Comprehensive Cancer Network (NCCN) **bladder cancer** clinical guidelines (version 1.2026 – March 16, 2026) recommend Anktiva in combination with BCG for the treatment of BCG-unresponsive, high-risk NMIBC with carcinoma in situ with or without Ta/T1 papillary tumors or Ta/T1 papillary tumors only without carcinoma in situ as initial treatment or for cytology-positive, imaging- and cystoscopy-negative, recurrent or persistent disease (category 2A).^{3,4}

POLICY STATEMENT

Prior Authorization is recommended for medical benefit coverage of Anktiva. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Extended approvals are allowed if the patient continues to meet the Criteria and Dosing. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Anktiva as well as the monitoring required for adverse events and long-term efficacy, approval requires Anktiva to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Anktiva is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Non-Muscle Invasive Bladder Cancer.** Approve for the duration noted if the patient meets ONE of the following (A or B):

A) **Initial Therapy:** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, and v):

Note: This allows enough time for a patient to complete two courses of induction therapy if needed.

i. Patient is ≥ 18 years of age; AND

ii. Patient has high risk Bacillus Calmette-Guerin (BCG)-unresponsive disease; AND

iii. Patient meets ONE of the following (a or b):

a) Patient has carcinoma in situ (CIS); OR

b) Patient has Ta/T1 papillary tumors without carcinoma in situ (CIS); AND

iv. Medication is used in combination with BCG; AND

v. Medication is prescribed by or in consultation with a urologist or an oncologist; OR

B) **Maintenance Therapy:** Approve for 3 months if the patient meets ALL of the following (i, ii, and iii):

i. Patient has an ongoing complete response defined as ONE of the following (a or b):

a) Patient has negative cystoscopy and meets ONE of the following [(1) or (2)]:

(1) Negative urine cytology; OR

(2) Malignant urine cytology if cancer found in the upper tract or prostatic urethra and random bladder biopsies are negative; OR

b) Patient has positive cystoscopy with biopsy-proven benign or low-grade Ta non-muscle invasive bladder cancer and negative urine cytology; AND

ii. Medication is used in combination with BCG; AND

iii. Medication is prescribed by or in consultation with a urologist or an oncologist.

Dosing. Approve the following dosing regimens (A or B):

A) **Induction Therapy:** Approve 400 mcg administered intravesically once a week for 6 weeks. A second course of induction therapy can be administered at Month 3 if a complete response was not achieved with the first course; OR

B) **Maintenance Therapy:** Approve 400 mcg administered intravesically once a week for 3 weeks at Months 4, 7, 10, 13, and 19. Additional course can be given at Months 25, 31, and 37.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Anktiva is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Anktiva intravesical solution [prescribing information]. Culver City, CA: ImmunityBio; April 2024.
2. Chamie K, Chang SS, Kramolowsky E, et al. IL-15 superagonist NAI in BCG-unresponsive non-muscle-invasive bladder cancer. *NEJM Evid.* 2023;2(1):EVIDoa2200167.
3. The NCCN Bladder Cancer Clinical Practice Guidelines in Oncology (version 1.2026 – March 16, 2026). © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on April 3, 2026.
4. The NCCN Drugs & Biologics Compendium. © 2026 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Search term: nogapendekin. Accessed on April 3, 2026.

HISTORY

| Type of Revision | Summary of Changes | Review Date |
|------------------|--|-------------|
| New Policy | -- | 05/08/2024 |
| Update | 04/04/2025: The policy name was changed from “Oncology (Other) - Anktiva UM Medical Policy” to “Oncology (Intravesical) - Anktiva UM Medical Policy”. | NA |
| Annual Revision | Non-Muscle Invasive Bladder Cancer: For the requirement that the patient has Bacillus Calmette-Guerin (BCG) unresponsive disease, added “high-risk” as a qualifier. For the requirement that the patient has carcinoma in situ, removed “with or without papillary tumors”. | 04/30/2025 |
| Annual Revision | Non-Muscle Invasive Bladder Cancer: An option for approval was added for a patient who has Ta/T1 papillary tumors without carcinoma in situ (CIS). | 04/08/2026 |