



MEDICAL POLICY STATEMENT

Arkansas PASSE

Policy Name & Number	Date Effective
Trigger Point Injections-AR PASSE-MM-1503	04/01/2026
Policy Type	
MEDICAL	

Medical Policy Statement prepared by CareSource and its affiliates are derived from literature based on and supported by clinical guidelines, nationally recognized utilization and technology assessment guidelines, other medical management industry standards, and published MCO clinical policy guidelines. Medically necessary services include, but are not limited to, those health care services or supplies that are proper and necessary for the diagnosis or treatment of disease, illness, or injury and without which the patient can be expected to suffer prolonged, increased or new morbidity, impairment of function, dysfunction of a body organ or part, or significant pain and discomfort. These services meet the standards of good medical practice in the local area, are the lowest cost alternative, and are not provided mainly for the convenience of the member or provider. Medically necessary services also include those services defined in any Evidence of Coverage documents, Medical Policy Statements, Provider Manuals, Member Handbooks, and/or other policies and procedures.

Medical Policy Statements prepared by CareSource and its affiliates do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage) for the service(s) referenced in the Medical Policy Statement. If there is a conflict between the Medical Policy Statement and the plan contract (i.e., Evidence of Coverage), then the plan contract (i.e., Evidence of Coverage) will be the controlling document used to make the determination. According to the rules of Mental Health Parity Addiction Equity Act (MHPAEA), coverage for the diagnosis and treatment of a behavioral health disorder will not be subject to any limitations that are less favorable than the limitations that apply to medical conditions as covered under this policy.

Table of Contents

A.	Subject	2
B.	Background	2
C.	Definitions	2
D.	Policy	2
E.	Conditions of Coverage	4
F.	Related Policies/Rules	4
G.	Review/Revision History	4
H.	References	5

A. Subject

Trigger Point Injections

B. Background

Myofascial trigger points are self-sustaining, hyper-irritative foci in any skeletal muscle, often occurring in response to strain produced by acute or chronic overload. There is no associated neurologic deficit, and the pain may be aggravated by hyperextension of the spine, standing, and walking. These trigger points produce a referred pain pattern characteristic for that individual muscle. Each pattern becomes part of a single muscle myofascial pain syndrome (MPS) which is responsive to appropriate treatment. To successfully treat chronic myofascial pain syndrome, each single muscle syndrome needs to be identified along with every perpetuating factor. The purpose of a trigger point injection (TPI) is to treat not only the symptom but also the cause through the injection of a single substance (eg, a local anesthetic) or a mixture of substances (eg, a corticosteroid with a local anesthetic) directly into the affected body part in order to alleviate inflammation and pain.

Interventional procedures for management of pain should be part of a comprehensive pain management care plan that incorporates an initial trial of conservative treatment utilizing appropriate medications, physical therapy modalities, and behavioral support, as needed.

Interventional procedures for the management of pain unresponsive to conservative treatment should be completed only by qualified healthcare providers acting within their scope of practice.

C. Definitions

- **Acute Pain** – Pain that lasts less than 4 weeks.
- **Physician Supervised Home Exercise Program (HEP)** – A 6-week program requiring an exercise prescription and/or plan and a follow-up documented in the medical record after completion, or documentation of the inability to complete the HEP due to a stated physical reason (ie, increased pain, inability to physically perform exercises). Patient inconvenience or noncompliance without explanation does not constitute an inability to complete.
- **Subacute Pain** – Pain that has lasted between 4 weeks and 12 weeks.
- **Trigger Point** – A hyper excitable area of the body, where the application of a stimulus will provoke pain to a greater degree than in the surrounding area.

D. Policy

I. Trigger Point Injections – Initial Injection

- A. Trigger point injections of anesthetic and/or corticosteroid for back pain, neck pain, or myofascial pain syndrome are considered medically necessary when pain has persisted despite appropriate medical management and **ALL** the following criteria are met:

The MEDICAL Policy Statement detailed above has received due consideration as defined in the MEDICAL Policy Statement Policy and is approved.

1. Member presents with new (acute or subacute) localized pain, occurring in the last 3 months.
 2. Member has been refractory or intolerant of conservative therapies for at least 1 month, including **at least 1** of the following:
 - a. bed rest
 - b. active exercise
 - c. ultrasound
 - d. range of motion
 - e. heating or cooling treatments
 - f. massage
 3. TPIs are being given as a part of an overall conservative management (usually short term) plan, including **at least 1** of the following:
 - a. physical therapy
 - b. occupational therapy
 - c. physician supervised home exercise program (HEP)
 - d. manipulative therapy
 4. Pharmacotherapies are being administered, including **at least 1** of the following:
 - a. non-steroidal anti-inflammatory drugs (NSAIDS)
 - b. muscle relaxants
 - c. non-narcotic analgesics
 - d. anti-depressants
 5. The member must have a diagnosis for which the trigger point injection is an appropriate treatment. **ALL** the following information must be documented in the patient's medical record:
 - a. proper evaluation including a patient history and physical examination leading to diagnosis of the trigger point
 - b. reason or reasons for selecting this therapeutic option
 - c. affected muscle or muscles
 - d. muscle or muscles injected and the number of injections
 - e. frequency of injections required
 - f. name of the medication used in the injection
 - g. results of any prior treatment
 - h. corroborating evidence that the injection is medically necessary
- B. Localization techniques to image or otherwise identify trigger point anatomic locations are not indicated and will not be covered for payment when associated with trigger point injection procedures.
- II. Trigger Point Injections – Subsequent Injections
- A. Trigger point injections should be repeated only if doing so is reasonable and medically necessary. No more than 8 dates of service per calendar year per member may be reimbursed.
 - B. Injections may be repeated only with documented positive results to the most recent trigger point injection of the same anatomic site. Documentation should include at least 50% improvement in pain, functioning, and activity tolerance.

- III. There is no laboratory or imaging test for establishing the diagnosis of trigger points. Diagnosis is dependent upon a detailed history and a thorough, directed examination. The following clinical features are present most consistently and are helpful in making the diagnosis:
- A. history of onset and its cause (injury, sprain, etc.)
 - B. distribution of pain
 - C. restriction of movement
 - D. mild muscle specific weakness
 - E. focal tenderness of a trigger point
 - F. palpable taut band of muscle in which trigger point is located
 - G. local taut response to snapping palpitation
 - H. reproduction of referred pain pattern upon most sustained mechanical stimulation of the trigger point

IV. Payment Information

- A. Certain procedure codes for trigger point injections specify the number of injection sites. For these codes, the unit of service is different from the number of injections given.
- B. Payment may be made for 1 unit of service of the appropriate procedure code reported on a claim for service rendered to a particular patient on a particular date.
- C. A trigger point injection is normally considered a stand-alone service. No additional payment will be made for an office visit on the same date of service unless there is an indication on the claim (eg, in the form of a modifier appended to the evaluation and management procedure code) that a separate evaluation and management service was performed.
- D. Dry needling is not an acceptable alternative to trigger point injections by an appropriately licensed clinician.

E. Conditions of Coverage

NA

F. Related Policies/Rules

NA

G. Review/Revision History

	DATE	ACTION
Date Issued	05/10/2023	New policy, approved at Committee.
Date Revised	02/28/2024	Annual review: updated references, approved at Committee
	02/12/2025	Review: updated references, approved at Committee
	01/14/2026	Review: updated references. Approved at Committee.
Date Effective	04/01/2026	
Date Archived		

The MEDICAL Policy Statement detailed above has received due consideration as defined in the MEDICAL Policy Statement Policy and is approved.

H. References

1. Appasamy M, Lam C, Alm J, et al. Trigger point injections. *Phys Med Rehabil Clin N Am*. 2022;33(2):307-333. doi:10.1016/j.pmr.2022.01.011
2. Chou R. Subacute and chronic low back pain: nonsurgical interventional treatment. UpToDate. Updated April 29, 2025. Accessed December 12, 2025. www.uptodate.com
3. Debrosse M, Sukhman S, Shah A, et al. Trigger point injection therapies for chronic myofascial neck and back pain: a systematic review. *Interventional Pain Medicine*. 2022; 1(3):100076. doi.org/10.1016/j.inpm.2022.100076
4. Hammi C, Schroader JD, Yeung B. Trigger point injection. *StatPearls*. StatPearls Publishing; 2024. Updated July 24, 2023. Accessed December 12, 2025. www.ncbi.nlm.nih.gov
5. Hamzoian H, Zograbyan V. Trigger point injections versus medical management for acute myofascial pain: a systematic review and meta-analysis. *Cureus*. 2023;15(8):e43424. doi:10.7759/cureus.43424
6. Isaac Z. Management of non-radicular neck pain in adults. UpToDate. Updated September 30, 2025. Accessed December 12, 2025. www.uptodate.com
7. Khoury AL, Keane H, Varghese F, et al. Trigger point injection for post-mastectomy pain: a simple intervention with high rate of long-term relief. *NPJ Breast Cancer*. 2021;7:123. doi:10.1038/s41523-021-00321-w
8. Meyer GW. Anterior cutaneous nerve entrapment syndrome. UpToDate. Updated February 14, 2025. Accessed December 12, 2025. www.uptodate.com
9. Moynihan LK, Elkadry E. Myofascial pelvic pain syndrome in females: treatment. UpToDate. Updated January 11, 2023. Accessed December 12, 2025. www.uptodate.com
10. Tu FF, As-Sanie S. Chronic pelvic pain in adult females: treatment. UpToDate. Updated September 4, 2025. Accessed December 12, 2025. www.uptodate.com

Independent Medical Review – January 2025