

## SPECIALTY GUIDELINE MANAGEMENT

### PROLIA (denosumab)

#### POLICY

##### I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

###### A. FDA-Approved Indications

1. Treatment of postmenopausal women with osteoporosis at high risk for fracture
2. Treatment to increase bone mass in men with osteoporosis at high risk for fracture
3. Treatment to increase bone mass in men at high risk for fracture receiving androgen deprivation therapy (ADT) for non-metastatic prostate cancer
4. Treatment to increase bone mass in women at high risk for fracture receiving adjuvant aromatase inhibitor therapy for breast cancer

###### B. Compendial Uses

Prevention or treatment of osteoporosis during androgen deprivation therapy for patients with high fracture risk

All other indications are considered experimental/investigational and are not a covered benefit.

##### II. CRITERIA FOR INITIAL APPROVAL

###### A. **Osteoporosis in Postmenopausal Women**

Authorization of 24 months may be granted to postmenopausal female members when ANY of the following criteria are met:

1. Member has a history of fragility fractures
2. Member has a pre-treatment T-score of  $\leq -2.5$  OR member has osteopenia with a high pre-treatment FRAX fracture probability (See Appendix B) and meets ANY of the following criteria:
  - a. Member has indicators of higher fracture risk (e.g., advanced age, frailty, glucocorticoid use, very low T-scores, or increased fall risk)
  - b. Member has failed prior treatment with or is intolerant to previous injectable osteoporosis therapy (e.g., zoledronic acid [Reclast], teriparatide [Forteo])
  - c. Member has had an oral bisphosphonate trial of at least 1-year duration or there is a clinical reason to avoid treatment with an oral bisphosphonate (See Appendix A)

###### B. **Osteoporosis in Men**

Authorization of 24 months may be granted to male members with osteoporosis when ANY of the following criteria are met:

1. Member has a history of an osteoporotic vertebral or hip fracture
2. Member has a pre-treatment T-score of  $\leq -2.5$
3. Member has osteopenia with a high pre-treatment FRAX fracture probability (See Appendix B)

###### C. **Breast Cancer**

Authorization of 24 months may be granted to members who are receiving adjuvant aromatase inhibitor therapy for breast cancer.

###### D. **Prostate Cancer**

Authorization of 24 months may be granted to members who are receiving androgen deprivation therapy for prostate cancer.

### III. CONTINUATION OF THERAPY

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

### IV. APPENDIX

#### Appendix A. Clinical reasons to avoid oral bisphosphonate therapy

- Esophageal abnormality that delays emptying such as stricture of achalasia
- Active upper gastrointestinal problem (e.g., dysphagia, gastritis, duodenitis, erosive esophagitis, ulcers)
- Inability to stand or sit upright for at least 30 to 60 minutes
- Inability to take at least 30 to 60 minutes before first food, drink, or medication of the day
- Renal insufficiency (creatinine clearance <30 mL/min)
- History of intolerance to an oral bisphosphonate

#### Appendix B. WHO Fracture Risk Assessment Tool

- High FRAX fracture probability: 10 year major osteoporotic fracture risk  $\geq 20\%$  or hip fracture risk  $\geq 3\%$ .
- 10-year probability; calculation tool available at: <http://www.shef.ac.uk/FRAX/tool.jsp>

### V. REFERENCES

1. Prolia [package insert]. Thousand Oaks, CA: Amgen Inc.; August 2016.
2. The NCCN Drugs & Biologics Compendium™ © 2016 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed October 23, 2016.
3. Bisphosphonates. *Drug Facts and Comparisons. Facts & Comparisons® eAnswers* [online]. 2015. Available from Wolters Kluwer Health, Inc. Accessed October 20, 2016.
4. Cosman F, de Beur SJ, LeBoff MS, et al. National Osteoporosis Foundation. Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int*. 2014;25(10): 2359-2381.
5. Jeremiah MP, Unwin BK, Greenwald MH, et al. Diagnosis and management of osteoporosis. *Am Fam Physician*. 2015;92(4):261-268.
6. Watts NB, Bilezikian JP, Camacho PM, et al. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the diagnosis and treatment of postmenopausal osteoporosis. *Endocr Pract*. 2016;22 (Suppl 4):1-42.
7. ACOG Practice Bulletin Number 129: Osteoporosis. *Obstet Gynecol*. 2012;120(3):718-734. – no change
8. National Institute for Health and Care Excellence. Osteoporosis Overview. Last updated September 20, 2016. Available at: <http://pathways.nice.org.uk/pathways/osteoporosis>. Accessed October 17, 2016.
9. Treatment to prevent osteoporotic fractures: an update. Department of Health and Human Services, Agency for Healthcare Research and Quality. 2012; Publication No. 12-EHC023-EF. Available at [www.effectivehealthcare.ahrq.gov/lbd.cfm](http://www.effectivehealthcare.ahrq.gov/lbd.cfm).
10. Watts NB, Adler RA, Bilezikian JP, et al. Osteoporosis in men : an Endocrine Society clinical practice guideline. *J Clin Endocr Metab*. 2012;97(6):1802-1822. – no change, guidelines coming out in 2017
11. Gralow JR, Biermann S, Farooki A, et al. NCCN Task Force Report: Bone Health in Cancer Care. *JNCCN*. 2013; 11(Suppl 3):S1-50
12. FRAX® WHO fracture risk assessment tool. © World Health Organization Collaborating Centre for Metabolic Bone Diseases: University of Sheffield, UK. Available at: <http://www.shef.ac.uk/FRAX>. Accessed October 7, 2015.