

## PHARMACY POLICY STATEMENT TRICARE

<b>DRUG NAME</b>	<b>Pombiliti (cicapglucosidase alfa-atga) and Opfolda (miglustat)</b>
<b>BENEFIT TYPE</b>	Medical, Pharmacy
<b>STATUS</b>	Prior Authorization Required

Pombiliti and Opfolda, approved by the FDA in 2023, are indicated to be used in combination for the treatment of adult patients with late-onset Pompe disease (lysosomal acid alpha-glucosidase [GAA] deficiency) weighing  $\geq 40$  kg and who are not improving on their current enzyme replacement therapy (ERT). Pombiliti provides an exogenous source of the deficient GAA enzyme, and Opfolda is an enzyme stabilizer.

Pompe disease is a rare, genetic lysosomal storage disorder that results in the buildup of glycogen in cell lysosomes causing serious and life-threatening muscle damage and weakness. It can be broadly classified as infantile onset within the first few months of life (IOPD) or late onset beyond infancy (LOPD). Classic IOPD is rapidly progressive with severe cardiomyopathy. Non-classic IOPD progresses slower with less severe cardiomyopathy. LOPD does not typically present with cardiomyopathy and has more variable symptoms, especially skeletal muscle weakness. Pombiliti and Opfolda are only indicated to treat LOPD.

Of note, miglustat is also the active ingredient in Zavesca, which is approved for Gaucher disease.

Miglustat may also be used off-label for the treatment of Niemann-Pick disease type C, as supported by guidelines. As of 2024, Miplyffa is FDA approved for use with miglustat for NPC.

Pombiliti (cicapglucosidase alfa-atga) and Opfolda (miglustat) will be considered for coverage when the following criteria are met:

### Pompe Disease

For initial authorization:

1. Member is at least 18 years of age; AND
2. Medication must be prescribed by or in consultation with a geneticist, neurologist, pulmonologist, or metabolic specialist; AND
3. Pombiliti and Opfolda are being prescribed in combination; AND
4. Member has a diagnosis of late onset Pompe disease confirmed by an enzyme activity assay showing GAA deficiency (2% to 40% of normal); AND
5. Molecular genetic testing shows pathogenic mutation of the GAA gene; AND
6. Member is not improving on their current enzyme replacement therapy (i.e., Lumizyme or Nexviazyme); AND
7. Member must show signs or symptoms (i.e., motor weakness, reduced respiratory parameters); AND
8. Documentation of baseline percent-predicted forced vital capacity (FVC) and 6-minute walk test (6MWT); AND
9. If female, attestation that the member is not pregnant.
10. **Dosage allowed/Quantity limit:** Starting 2 weeks after the last ERT dose:  
Pombiliti: 20 mg/kg every other week IV infusion



Opfolda: Orally every other week; 260 mg for patients weighing  $\geq 50$  kg or 195 mg for patients weighing  $\geq 40$  kg to  $< 50$  kg

**If all the above requirements are met, the medication will be approved for 12 months.**

For **reauthorization**:

1. Chart notes must document positive clinical response such as improved or stabilized respiratory muscle strength (i.e., forced vital capacity (FVC)) or functional endurance (e.g., 6-minute walk test).

**If all the above requirements are met, the medication will be approved for an additional 12 months.**

## Niemann-Pick disease type C (NPC); OPFOLDA ONLY

For **initial** authorization:

1. Medication must be prescribed by or in consultation with a neurologist, metabolic specialist, or geneticist; AND
2. Member has a diagnosis of NPC confirmed by one of the following:
  - a) Genetic testing that shows mutations in both alleles of the *NPC1* or *NPC2* gene, or
  - b) Mutation in one allele of *NPC1* or *NPC2*, AND either elevated plasma biomarkers (i.e., cholestanetriol or trihydroxy-cholanoyl-glycine and/or lysoSM-509 with normal or slightly elevated lysoSM) OR positive filipin testing; AND
3. Member has at least 1 neurological sign of disease; AND
4. Member does NOT have advanced neurological disease.

5. **Dosage allowed/Quantity limit:**  
Opfolda: 3 capsules (195 mg) three times a day (off-label).

**If all the above requirements are met, the medication will be approved for 12 months.**

For **reauthorization**:

1. Chart notes must document improvement or stabilization of neurological signs and symptoms compared to baseline.

**If all the above requirements are met, the medication will be approved for an additional 12 months.**

**TRICARE Prime® Demo by CareSource Military & Veterans™ considers Pombiliti (cipaglucosidase alfa-atga) and Opfolda (miglustat) not medically necessary for the treatment of conditions that are not listed in this document. For any other indication, please refer to the Off-Label policy.**

DATE	ACTION/DESCRIPTION
10/25/2023	New policy for Pombiliti and Opfolda created.
11/21/2024	Added criteria for diagnosis of NPC (approved in combination with Miplyffa; off-label as monotherapy or in combination with Aqneursa). Added pharmacy as benefit type.



## References:

1. Pombili [prescribing information]. Amicus Therapeutics US, LLC; 2023.
2. Opfolda [prescribing information]. Amicus Therapeutics US, LLC; 2024.
3. Schoser B, Roberts M, Byrne BJ, et al. Safety and efficacy of cipaglucosidase alfa plus miglustat versus alglucosidase alfa plus placebo in late-onset Pompe disease (PROPEL): an international, randomised, double-blind, parallel-group, phase 3 trial [published correction appears in Lancet Neurol. 2023 Oct;22(10):e11]. *Lancet Neurol.* 2021;20(12):1027-1037. doi:10.1016/S1474-4422(21)00331-8
4. Cupler EJ, Berger KI, Leshner RT, et al. Consensus treatment recommendations for late-onset Pompe disease. *Muscle Nerve.* 2012;45(3):319-333. doi:10.1002/mus.22329
5. Van der Ploeg AT, Kruijshaar ME, Toscano A, et al. European consensus for starting and stopping enzyme replacement therapy in adult patients with Pompe disease: a 10-year experience. *Eur J Neurol.* 2017;24(6):768-e31. doi:10.1111/ene.13285
6. Wang RY, Bodamer OA, Watson MS, Wilcox WR; ACMG Work Group on Diagnostic Confirmation of Lysosomal Storage Diseases. Lysosomal storage diseases: diagnostic confirmation and management of presymptomatic individuals. *Genet Med.* 2011;13(5):457-484. doi:10.1097/GIM.0b013e318211a7e1
7. Tarnopolsky M, Katzberg H, Petrof BJ, et al. Pompe Disease: Diagnosis and Management. Evidence-Based Guidelines from a Canadian Expert Panel. *Can J Neurol Sci.* 2016;43(4):472-485. doi:10.1017/cjn.2016.37
8. Geberhiwot T, Moro A, Dardis A, et al. Consensus clinical management guidelines for Niemann-Pick disease type C. *Orphanet J Rare Dis.* 2018;13(1):50. Published 2018 Apr 6. doi:10.1186/s13023-018-0785-7
9. Patterson MC, Clayton P, Gissen P, et al. Recommendations for the detection and diagnosis of Niemann-Pick disease type C: An update. *Neurol Clin Pract.* 2017;7(6):499-511. doi:10.1212/CPJ.0000000000000399
10. Patterson MC, Vecchio D, Prady H, Abel L, Wraith JE. Miglustat for treatment of Niemann-Pick C disease: a randomised controlled study. *Lancet Neurol.* 2007;6(9):765-772. doi:10.1016/S1474-4422(07)70194-1
11. Mengel E, Patterson MC, Da Riol RM, et al. Efficacy and safety of arimoclomol in Niemann-Pick disease type C: Results from a double-blind, randomised, placebo-controlled, multinational phase 2/3 trial of a novel treatment. *J Inherit Metab Dis.* 2021;44(6):1463-1480. doi:10.1002/jimd.12428

Effective date: 01/01/2026

Revised date: 11/21/2024